Punctuating DNA

The epigenome controls the differential expression of genes in a cell-, tissue-, or time-specific manner via inheritance without associated DNA sequence alterations. Epigenetic control involves several mechanisms, including DNA methylation, histone modifications and variants, and chromatin structural alterations. DNA methylation is the most stable type of currently identified epigenetic modification. Methylation of CpG islands within promoter regions is associated with transcriptional repression (silencing) of associated genes. Histone acetylation, phosphorylation, and methylation play roles in the positioning of nucleosomes on DNA, affecting gene expression by controlling access of the transcription machinery to genes.

Inappropriate epigenetic programming has been associated with human diseases, including cancer, neurodevelopmental disorders, aging-related conditions, and autoimmunity. Interaction of the epigenome with the environment, including pollutants, infections, stress, and nutrition can alter patterns of gene expression and may lead to late onset diseases. Although heritable, at least in somatic cells, agents capable of reversing epigenetic changes are being identified and used clinically [e.g., DNA demethylating drugs and histone deacetylase (HDAC) inhibitors].

A Role in Cancer

"It's a very exciting time now in the epigenetics of cancer biology," says Stephen Baylin, Professor of Oncology and Medicine, Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins University, Baltimore, MD. "The results with demethylating drugs and HDAC inhibitors in hematopoietic malignancies are astounding. It looks as if many more genes are epigenetically silenced in cancer than mutated, although the importance is not yet known." Candidate silenced genes include tumor suppressor

Deregulated cell cycle expression of DNA methyltransferase 1 (*DNMT1*), the main DNA methylation enzyme, leads to cellular transformation in NIH 3T3 cells. Ectopic expression of *DNMT1* induces entry into the S phase of the cell cycle. The 3' untranslated region (UTR) inhibits ectopic expression of *DNMT1* at the G0 phase of the cell cycle.

genes. DNA methylation in aberrant gene silencing and alterations affecting chromatin architecture are leading to more and more legitimate therapeutic targets for reversal of this silencing as a strategy for cancer prevention and/or therapy.

Moshe Szyf, Professor, Department of Pharmacology and Therapeutics, McGill University, Montreal, QC, Canada, refers to methylation as the "punctuation mark of DNA that over- and under-rides what genes are expressed and what not." His laboratory is interested in understanding the links between chromatin, DNA methylation, and cancer therapy, what mechanisms define DNA methylation patterns, and why these patterns are tightly correlated with chromatin structure. One question he'd like to answer is how to activate tumor suppressor genes without activating metastatic genes. For example, treating breast cancer cells with hypomethylating drugs increases their potential to metastasize. It's not yet known how to target hypomethylating drugs to specific sequences to activate desirable genes without activating undesirable ones. The cross-talk between chromatin and DNA methylation has therapeutic potential, Szyf observes. "If you change chromatin, you could change methylation. It's a dynamic world that offers potential. We're still scratching the surface."

Learning Process

Xinyu Zhao, Assistant Professor, Department of Neurosciences, University of New Mexico, Albuquerque, AZ, says, "epigenetics is pretty new for the neurosciences. Even 3 years ago I had problems finding reviewers for my papers." Her work focuses on identifying genetic and epigenetic factors that regulate postnatal neurogenesis and neural stem cell functions, including how DNA methylation functions in neurodevelopment. She is studying mouse models with mutations in methylated-CpG binding proteins (MeCP2 and MBD1) that exhibit phenotypes similar to human neurodevelopmental syndromes.

"There may be a limit to how far we can understand our own brain."

She believes a lot of neurodevelopmental and complex mental disorders result not from classic genetic mutations but from epigenetic changes. "It's very exciting, especially in the neuroscience field, now that we are past the genomic era. We know the blueprint, beyond that, what can we learn? However, there may be a limit to how far we can understand our own brain. Neurodevelopmental stud-

ies will require crossing fields, including stem cell research, epigenetics, and neurosciences. It's a learning process."

Karen Usdin, Chief, Gene Structure and Disease Section, National Institute of Diabetes, Digestive, and Kidney Diseases, National Institutes of Health, Bethesda, MD, studies Repeat Expansion Diseases, conditions with neurodevelopmental implications resulting from gene silencing. One of these diseases resulting from expansion is Friedreich ataxia (FA), caused by the expansion of a GAAoTTC repeat in the frataxin gene, leading to reduced levels of frataxin messenger RNA (mRNA), which in turn causes sensory motor neuron degeneration, diabetes, and cardiomyopathy. The second disease involves expansion of a CGGoCCG repeat in the 5' untranslated region of the fragile X mental retardation 1 (*FMR1*) gene, and the consequences depend upon the number of repeats in the expanded allele. Premutation alleles with 59 to 200 repeats are at risk of fragile X-associated tremor and ataxia syndrome (FXTAS). Female carriers of this allele are also at risk for fragile X-associated premature ovarian failure and can transmit an *FMR1* allele with greatly expanded repeats to offspring.

Carriers with full mutation alleles, those with over 200 repeats, are at very high risk of fragile X syndrome (FXS), a developmental disorder, characterized by moderate to severe mental retardation. Usdin's group is looking at a mouse model of premutation FXS, with 120 CGG-CGG repeats in the murine *Fmr1* gene.

Questions Usdin is trying to answer include how changes in an intron or 5' untranslated region change gene expression and how the number of repeats affects pathology. They are also exploring the feasibility of gene modification or reactivation as potential treatments for this disorder. Although other mechanisms are involved in FA and FXS, Usdin believes epigenetic changes play a role. "Chromatin modification is involved in FA, but little is known about events leading to epigenetic modification. Much of our thinking is informed by the cancer epigenetic field," she says. Observing that much about histone is not known, she feels it is one missing link in understanding gene silencing and epigenetic changes in signaling. For example demethylation drugs don't fully reverse gene silencing, and these drugs don't necessarily distinguish between normal and aberrant demethylation.

Toward an Epigenome Project

Clearly the Human Genome Project provided information that has opened up many new areas of research, but it has not answered the question of differential gene expression in the absence of DNA sequence changes. Benjamin Tycko, Associate Professor of Pathology, Columbia University, New York, NY, points out that the advantage of the Human Genome Project is that there really is a single human genome, with only a small amount of inter-individual polymorphism. Although an expensive endeavor, it was straightforward. However, there is no single human epigenome. As Szyf comments, the

The methylation single nucleotide polymorphism (MSNP) method for high-throughput genetic and epigenetic profiling in human tissues: two germ cell tumors (GCT), a breast carcinoma (breast CA), and normal placenta, showing different DNA methylation patterns. Left panel, class 1 SNP loci (insensitive to CpG methylation but reporting DNA copy number); right panel, class 2 SNP loci (sensitive to CpG methylation); both on Affymetrix 250K *NspI* SNP arrays, with SNP loci arranged vertically by position on human chromosomes. Red indicates increased and blue indicates decreased signal intensity relative to diploid control DNA. With *HpaII* predigestion class 1 SNP loci largely unaffected, some class 2 SNP loci reduced intensity (hypomethylation); with methylation-insensitive isoschizomer *MspI* nearly all class 2 loci strongly reduced in intensity (control), validating the MSNP strategy. GCT from male patients show gain of chromosome arm 12p (asterisks; red signal) and reduced X chromosome copy number (blue).

Spectral karyotyping (SKY) of a MBD1^{-/-} (methyl-CpG binding protein null) adult mouse neural stem/progenitor cell, which has increased genomic instability. There are 3 chromosomes 10 and 18 and an extra fragment of chromosome 2 (white arrow).

entire epigenome should include every cell type at every time point. The cancer epigenome differs from the normal epigenome, and each different cancer epigenome differs from the others. "Worse than that," Tycko says, "look at normal tissues, like neurons, liver cells, lymphocytes. In one person these cells all have different epigenomes. One approach which will be pursued is to pick a tissue and start somewhere to try to define a reference epigenome. How useful that will be remains to be seen, but it will go forward."

In that vein, the American Association for Cancer Research (AACR) has been organizing workshops and actively promoting plans to further a Human Epigenome Project. In Europe, work has already begun, and in the U.S., efforts in nonhuman models have been underway. Other countries, including China, Korea, and Japan, have also either begun work or have expressed an interest.

Joseph Costello, Associate Professor, University of California, San Francisco, CA, concurs that an epigenome project is complicated because the epigenome is a moving target. Not only are there many epigenomes in different cell types at different times in development, but all are subject to environmental influences. This, he says, is not just a problem, but provides the excitement of the project and will provide a framework for studying environmental interactions with the genome. "We are not determined entirely by genetics. How diet and exercise affect different people different ways is determined partly by epigenetics," he observes.

"We are not determined entirely by genetics."

An epigenomic analysis of several tissue types could provide a framework to suggest which specific genes should be analyzed in depth in both human populations and in other species that are related evolutionarily.

"The most exciting parts of science, the big steps, often come unexpectedly," Costello points out. One concept that is arising from the understanding that cancer has both genetic and epigenetic causes is the study of combination therapy targeting both genetic and epigenetic changes.

Sequencing is currently too expensive to answer many epigenetic questions, although bisulfite sequencing is being used to analyze DNA methylation patterns. New, high-throughput technologies are being developed, including modified bisulfite sequencing, matrix-assisted laser desorption/

ionization mass spectrometry (MALDI-MS), and methylation single nucleotide polymorphism (MSNP) analysis. Tycko and colleagues have adapted SNP arrays from Affymetrix (Santa Clara, CA, USA) to analyze genetics and epigenetics in one high-throughput MSNP system, which has been validated by comparing results with combined bisulfite restriction analysis and sequencing cloned PCR products from bisulfite-converted DNA. MSNP, Tycko says, provides reasonably dense coverage of most CpG sites.

Interesting analyses will be to compare cells from normal bone marrow and from patients with acute leukemia before and after treatment with one of the demethylating drugs [e.g., decitabine (5-aza-2'-deoxycytidine) or azacitidine]. Szyf would like to see a comparison of epigenomes exposed to different environments or as individuals move from one environment to another. Tycko concludes that with current technology, "epigenetic projects can't characterize the thousands of different epigenomes. High-throughput low cost methods are essential."

- Lynne Lederman is a freelance medical writer in Mamaroneck, NY.