The Rise of High Throughput

The realization that gene, protein, and metabolite space is vast has led to the development of novel high-throughput methodologies aimed directly at interrogating biomolecules and their interactions at previously impossible depths. In this special focus section, the editors of *BioTechniques* explore the development of high-throughput applications in five areas of life science (page 27), while a primary research article by Kapetis et al. details an enhanced software platform capable of automated cross-platform microarray data analysis (page 33). It is our hope that through the articles in this section, readers will obtain a deeper understanding of high-throughput methodologies and their evolution.

BioTechniques 53:25 (*BioTechniques Focus: The Rise of High Throughput* July 2012)
doi 10.2144/000113894

INTRODUCTION

- 25 **The Rise of High-Throughput**
The Editors, BioTechniques
doi 10.2144/0001137894

SPECIAL NEWS FEATURE

- 27 **Tales on the Road to High-Throughput**
The Editors, BioTechniques
doi 10.2144/000113892

REPORT

- 33 **ADMA 2.13: A major update for automated cross-platform microarray data analysis**
Dimos Kapetis, Ferdinando Clarelli, Federico Vitulli, Nicole Kerlero de Rosbo, Ottavio Beretta, Maria Foti, Paola Ricciardi-Castagnoli, and Francesca Zolezzi
doi 10.2144/000113889

Tales on the Road to High-Throughput

The editors of *BioTechniques* examine five fields where high-throughput methodologies are making significant impacts.

Robotics: from microplates to microfluidics

Advances in robotics have been vital to the recent ascendance of high-throughput methodologies in modern biological research, enabling automation of various repetitive laboratory tasks that tend to be error-prone if done manually. Robotic instrumentation, such as plate handlers, liquid dispensers, and plate readers, allows processing and analysis of large numbers of samples around the clock for applications ranging from large-scale drug screens to next-generation sequencing. The key element for numerous high-throughput techniques has been the refinement of the multiwell microplate, as sample throughput is determined in large measure by the number of wells per microplate. While 384-well plates have largely supplanted the original 96-well plates and are now fairly standard, scientists working in ultra high-throughput applications have embraced even higher capacity 1536-well and 3456-well plates. A challenge arises,

however, as wells get smaller and more numerous: how do you pipette the required nanoliter (or smaller) volumes rapidly with acceptable precision? This challenge has been met in part with the development of acoustic droplet ejection systems that can precisely and gently dispense nanoliter or even picoliter volumes into the wells of high-density microplates without any physical contact with the transferred solution (1). However, a tradeoff does occur with these systems, as the advantages of higher sample throughput and reduced reagent costs can be offset by the expense of acoustic liquid dispensers and the need for humidity-controlled environments to reduce the significant problem of rapid evaporation when dealing with sub-microliter volumes. The cost of the robotic instrumentation required to carry out high-throughput screens using higher-density microplates makes such systems inaccessible to many academic investigators with budgets far

more modest than those of large pharmaceutical companies. In this regard, then, the future of affordable high-throughput screening appears to be increasingly focused on microchip platforms—microarrays or microfluidics—that can greatly reduce, if not eliminate, the need for expensive robotics. In particular, microfluidic chips can manipulate tiny volumes of fluid in closed microchannels that are only tens of micrometers wide and continuously carry out sequential assay steps within a single chip. This results in a high-throughput platform that is inexpensive to fabricate, reduces analysis times, is flexible to integrate with other systems, and thus proves ideal for more affordable high-throughput applications.

1. Ellson, R., M. Mutz, et al. 2003. *Journal of Laboratory Automation*. 8:29-34.

Cloning: the more, the merrier

A key step in high-throughput proteomics and functional genomics studies is the cloning of open reading frames (ORFs) into expression vectors. Traditional cloning methods utilizing restriction digestion and ligation are not well-suited to high-throughput applications given their inefficiency and the constraints imposed by restriction site selection. Recombination-based and ligation-independent cloning (LIC) methods have proven to be more amenable to a high-throughput format. Site-specific recombination (SSR) methods, such as the Gateway or Creator systems, allow cloning of PCR-amplified ORF fragments flanked by specific recombination sites into suitable entry vectors containing corresponding recombination sites. The ORFs in entry vectors can then be easily transferred between different vectors. LIC uses the controlled digestion of insert and vector to generate complementary single stranded ends, which are

ligated together when co-transformed into bacterial cells. Both the SSR and LIC methods, however, suffer the drawback of adding undesirable extraneous sequences to the ORF being cloned. Engler et al. (1) overcame this problem with their “Golden Gate” method, a modification of the original Gateway system that uses type II restriction enzymes in a single-tube, one-step reaction for the seamless transfer of an ORF from a Gateway entry vector to an expression vector. More recently, Quan and Tian (2) devised an efficient method for the simple and economical assembly of circular DNA called circular polymerase extension cloning (CPEC) that is compatible for high-throughput cloning. CPEC is a single-tube, one-step reaction based on overlap extension PCR that is directional, sequence-independent, ligase-free and allows seamless cloning. Regardless of which method is chosen to insert a target DNA sequence into a vector, a

major bottleneck in most high-throughput cloning workflows is the need to manually plate transformed bacteria onto selective media and pick colonies in order to obtain the desired clone. However, Yehezkel et al. (3) recently described a new method for the high-throughput cloning of bacteria in liquid medium using standard multiwell plates that is completely controlled by liquid handling robots and plate readers, eliminating the need to plate and pick bacterial colonies.

1. Engler, C., R. Kandzia, et al. 2008. *PLoS One* 3: e3647.

2. Quan, J., and J. Tian. 2011. *Nat Protoc*. 6:242-51.

3. Ben Yehezkel, T., S. Nagar, D. Mackrants, Z. Marx, G. Linshiz, et al. 2011. *BioTechniques*. 50:124-7.

Sequencing: the long and short of it

Sanger sequencing (1) easily ranks among the most influential methodologies devised in the past 50 years. Yet as with many modern techniques that become widely adopted, researchers have continued to automate the sequencing process. Although some automation was quickly implemented, it took a couple of decades until Applied BioSystems (ABI) introduced the first fully automated DNA sequencer. That instrument relied on capillary electrophoresis rather than traditional slab gels, thus allowing automated injection of samples. With further refinements, highly accurate read lengths of about 1000 bp could be obtained using capillary sequencing and this platform became the workhorse of the Human Genome Project (HGP). Upon completion of the HGP, many researchers shifted their focus from sequencing to resequencing. The result was the development of a new generation of massively parallel next-generation sequencing (NGS) platforms that offered dramatically greater numbers of sequence reads from a single experiment. The compromise, however, was a reduction in the possible read length that could be attained compared with capillary instruments. The first commercially introduced NGS system was the Roche 454 platform (2), capable of

400,000 200-300 bp reads at 99% accuracy within four hours. The methodology relies on emulsion PCR with single primers tethered to beads that are arrayed in picoliter scale wells for pyrosequencing. The next NGS system appeared on the scene in 2006. The first iteration of the Illumina Genome Analyzer took advantage of bridge PCR with alternating cycles of extension and denaturation, resulting in dense clusters of double stranded DNA in each channel of a flow cell where reversible dye terminators could be visualized in four channels. 2008 brought the arrival of the SOLiD sequencing platform (ABI) and the Polonator, which both rely on emulsion PCR with randomly immobilized smaller beads, creating significantly higher density sequencing arrays. The major difference for the SOLiD platform was that sequencing was driven by ligase, an approach that results in a reduced rate of errors. "Third-generation" single molecule sequencing technologies have recently become available. These systems include the HeliScope system from Helicos, the first single molecule sequencing technology available; Ion Torrent which detects sequence using electrical changes across a semiconductor chip (2011); and SMRT sequencing from Pacific Biosciences

(2010), which attaches a DNA polymerase to the bottom of a well and monitors the incorporation of fluorescently labeled nucleotides. With SMRT, simultaneous monitoring of polymerase kinetics offers the ability to detect methylation and other epigenomic markers for the first time in a sequencing system. In one day, this system allows the detection of up to 75,000 single molecule sequencing reactions in parallel, with read lengths greater than 3000 bp. The next step in massively parallel sequencing methods appears to be the continued refinement of single molecule sequencing, albeit with longer read lengths. In 2012, Oxford Nanopore Technologies announced the upcoming release of a nanopore-based system capable of 5.4-100 kb read lengths and a throughput of hundreds of kb per second.

1. Sanger F, S. Nicklen, and A.R. Coulson. 1977 Dec. *Proc Natl Acad Sci U S A*. 74(12):5463-7
2. Margulies, M., et al. 2005 Sep 15. *Nature*. 437(7057):376-80. Epub 2005 Jul 31.
3. Harris, T.D., et al. 2008 Apr 4. *Science*. 320(5872):106-9.

PCR: amplification by miniaturization

Since its introduction in 1983, PCR has become a staple in the lives of life scientists worldwide. It has been modified, automated, multiplexed, and monitored in real-time yielding countless variations on the original theme. By enabling the amplification and detection of specific nucleic acid sequences from complex samples, PCR facilitated rapid advancements in cloning and sequencing. But applications for template quantification lagged behind until 1992, when Higuchi et al. introduced real-time PCR. Fluorescent dyes and probes enabled researchers to monitor the quantity of product being formed during each PCR amplification cycle; using estimates of amplification efficiency, it became possible to calculate the number of molecules carrying a specific sequence present in any sample. The subsequent improvement of detection chemistries and instrumentation spurred an increase in multiplex PCR capabilities wherein multiple DNA targets could be

amplified in the same tube using combinations of different fluorophores. These advances reduced sample requirements and provided significant cost savings in large-scale genotyping, gene expression, environmental monitoring, forensic, and diagnostic applications. The continued demand for higher throughput has led to the parallelization and miniaturization of PCR reactions, including development of microfluidic-based PCR platforms. One such platform, Access Array (Fluidigm), became available in 2009, essentially enables simultaneous creation of 48 barcoded, target enriched, and amplicon tagged libraries for sequencing. That same year, Life Technologies commercialized the OpenArray, a high-density array of up to 3000 individual nanoliter PCR reactions contained on a plate the size of a microscope slide. More recently, RainDance Technologies, a pioneer in the use of droplet approaches for targeted resequencing, announced the RainDrop Digital PCR System, a high-

throughput platform capable of generating more than a billion picoliter reaction droplets, each containing all the components necessary for a single PCR amplification reaction in a single day. RainDrop and a similar platform, QX100 Droplet Digital PCR System, provide absolute template quantification by screening each droplet, a digital PCR solution which is ideal for studies of copy number variation, rare sequence and mutation detection, and gene expression analysis. As with other techniques, it is expected that demands for PCR throughput will increase, driving development of new and innovative methods and technologies to meet those needs, each of which will increase capacity, while reducing instrument size, reaction cost, and labor.

1. Higuchi, R., G. Dollinger, P.S. Walsh, and R. Griffith. 1992. *Bio-Technology* 10(4), 413-417

Protein analysis: tearing into the protein universe

The desire to clone, express, and purify proteins quickly, and in large number, led to the emergence of high-throughput protein analysis techniques starting in the early 2000s. While His- or GST-tagged proteins can be routinely purified for downstream analysis with some optimization of conditions, when purifying proteins en masse, the challenge becomes identifying expression and purification conditions that work for not just one protein, but for a group of related, or even unrelated, proteins. Early work in high-throughput protein expression showed both the possibilities and difficulties in high-throughput protein analysis. In 2004, Luan et al. cloned and expressed 10,167 *C. elegans* open reading frames in *E. coli*. The results showed the potential for mass expression of proteins—expression was reported for 4854 ORFs of which 1536 were soluble. Although a good start, the result also represented expression of less than 50% of the cloned ORFs in the study. Other high-throughput studies have demonstrated higher percentages of expression, albeit with less proteins. In 2003, Agaton et al. expressed 142 human proteins with a reported success rate of 76% paving the

way for large-scale antibody production efforts. These early efforts were followed by further improvements and optimization which have led researchers to integrate high-throughput protein expression and purification methodologies even more effectively into a number of downstream applications. Protein structure determination is crucial to understanding the protein universe, and one of the biggest proponents of high-throughput protein structure determination has been the NIH-funded Protein Structure Initiative (PSI). Started in 2000, PSI has been instrumental in advancing methodologies for protein expression, purification, and characterization—PSI efforts have resulted in more than 50,000 plasmid clones which are now available from the PSI: Biology Materials Repository at Arizona State University. To date, the PSI has solved several thousand protein structures, and the project has moved into Phase III, applying the tools and methods developed during the initial phases toward specific biomedical problems. Yet another application where refinement of high-throughput protein expression and purification technology is creating new possibilities is in the generation of antibodies. The Human

Protein Atlas project, an effort to generate renewable antibodies to human proteins, has expanded from the work of Agaton et al. in 2003 to the present where the project's protein expression and purification team has developed and implemented a fully automated system enabling purification of up to 60 cell lysates at a time. Given the expansive scope of the protein universe—which is estimated to number in the millions when you take into account protein modifications—the need for high-throughput approaches is clear. In the coming years, expect even more technical refinement, which should lead to a clearer understanding of basic protein biochemistry.

1. Luan, C-H., Qiu, S., Finley, S.B., et al. 2004. *Genome Research*, 14 (10b):2102-2110.

2. Agaton, C., Galli, J., et al. 2003. *Molecular & Cellular Proteomics*, 2:405-414.

BioTechniques 53:25-31 (July 2012)
doi 10.2144/000113892

To purchase reprints of this article, contact:
biotechniques@fosterprinting.com

BACHEM

Bachem. Leading beyond peptides

**From Venoms to
Drugs!**

melusine®

www.bachem.com/melusine

Melusine® kits are unexplored **Libraries of Peptides & Proteins**, pre-fractionated and ready for HTS lead discovery, distributed by Bachem.

Melusine® is a product of Atheris Laboratories, Geneva - Switzerland

Atheris
DIFFUSION