Benchmarks

come and may be chosen for the convenience of the investigator. We used primarily neonatal toe and ear punch tissues and chose a 30-min heating time. Note also that the control DNA (Figure 2, lane C) produced more nonspecific PCR product than the HotSHOT DNA. This was confirmed in another experiment that compared HotSHOT DNA to DNA prepared by other methods and found that HotSHOT produced the cleanest PCR products (data not shown).

The HotSHOT method has been used in our laboratories to amplify 39 different products, including microsatellites, up to 1.9 kb in length. Standard PCR conditions were used for all assays. No primer pairs failed with HotSHOT DNA. Although we usually run PCR soon after preparing the DNA, we have stored samples up to three months at 4°C and -20°C before running PCR

and have had good results. Reviewing our records of a specific assay run with DNA prepared by traditional extraction methods and the same assay run with HotSHOT DNA, we found that 26 out of 1079 reactions failed with DNA prepared by traditional methods and only 9 out of 1011 reactions failed with HotSHOT DNA (P-value of Fisher's exact test, 0.0095), indicating that HotSHOT DNA performed slightly better than DNA prepared by traditional methods. The same reagents have also been used to prepare DNA from bacterial colonies for PCR; no heating was required for this application.

In an additional experiment, we found that HotSHOT DNA is not suitable for genotyping transgenic mice by Southern analysis. The DNA is extensively sheared and its concentration is too low. It is also possible that the DNA was not efficiently digested by the restriction enzyme because of interfering substances in the DNA preparation or because the DNA did not reanneal efficiently after it was neutralized. Although HotSHOT DNA preparation is likely to be limited to PCR applications, the savings in time and reagents over traditional DNA preparation methods is substantial, and the quality of the results is as good or slightly better than DNA prepared by traditional methods.

REFERENCES

1. Cesarone, C., C. Bolognesi, and L. Santi. 1979. Improved microfluorometric DNA determination in biological material using 33258 Hoechst. *Anal. Biochem.* 102:188-197.
2. Rudbeck, L. and J. Dissing. 1998. Rapid, simple alkaline extraction of human genomic DNA from whole blood, buccal epithelial cells, semen and forensic stains for PCR. *BioTechniques* 25:588-592.
3. Strauss, W.M. 1998. Preparation of genomic DNA from mammalian tissue, p. 2.2.1-2.2.3. In Ausubel et al. (Eds.), *Current Protocols in Molecular Biology*. John Wiley & Sons, New York.

Address correspondence to Dr. Gary E. Truett, Developmental Genetics Laboratory, Pennington Biomedical Research Center, 6400 Perkins Road, Baton Rouge, LA 70808, USA. e-mail: truettge@pbrc.edu

Received 15 November 1999; accepted 10 March 2000.

G.E. Truett, P. Heeger¹, R.L. Mynatt, A.A. Truett, J.A. Walker and M.L. Warman¹

Pennington Biomedical
Research Center
Baton Rouge, LA

¹Case Western Reserve
University
Cleveland, OH, USA

Choice of Microcentrifuge Tubes Influences T Cell Proliferation Assay

BioTechniques 29:54-59 (July 2000)

Microcentrifuge tubes are widely used in life science laboratories, including those focusing on cell culture, molecular biology and microbiology. Typically holding 0.2, 0.5 or 1.5–1.6 mL, these tubes generally provide passive support for reactions or storage of liquids. Recently, we realized that this is not always the case. We routinely perform proliferation assays with T lymphocyte cell lines to assess substances and media factors for inhibitory activities (1) and have observed that some sets of samples all contained some inhibitory capability. Subsequent investigation suggested that one brand of microcentrifuge tube used for storage and dilution of examined substances affected T cell proliferation responses. Therefore, we performed a systematic assessment of microcentrifuge tubes obtained from different manufacturers in our proliferation assay.

A human T lymphocyte line from peripheral blood was established and kept in culture by repeated stimulation as previously published (2). Lymphocyte proliferation was measured after anti-CD3 stimulation by a thymidine incorporation assay as previously published (1). In short, 96-well, flat-bottom plates were precoated with goat-anti-mouse antibody and then coated with a mouse-antihuman CD3 mAb (Ortho-

Figure 2. Effect of heating time on amplification of a 182-bp PCR product from various mouse tissue samples. Control DNA (C) was prepared by a traditional DNA extraction method. HotSHOT DNA was heated as described in Figure 1, followed by PCR amplification.

Benchmarks

clone OKT® 3; Janssen, Beerse, Belgium). After being washed to remove unbound mAb, 100 000 T lymphocytes were added in 100- μ L cell culture medium (RPMI; Life Technologies, Rockville, MD, USA) with 10% fetal calf serum (Sigma, St. Louis, MO, USA). Then, 100 μ L of a test substance were added. After a three-day culture, each well was pulsed with 10 μ Ci 3 H-deoxythymidine (Amersham Pharmacia Biotech, Dübendorf, Switzerland), and cells were harvested 18 h later. The total amount of incorporated radiolabel, as a measure of T cell proliferation, was quantified by scintillation counting (Microman; Wallac, Turku, Finland).

To test the effect of various types of microcentrifuge tubes, sterile Hanks balanced salt solution (HBSS; Sigma) was stored in the respective tube for 30 min at room temperature before introduction into the assay. All tubes were previously sterilized by autoclaving.

HBSS from a glass tube 16 $\times$ 160 mm (22 mL; Merck, Dietikon, Switzerland) was used as a negative control. In addition, experiments were performed in which an inhibitor of lymphocyte proliferation, isolated from human intestinal epithelial cells and currently being investigated by our laboratory, was serially diluted in HBSS in a specific type of microcentrifuge tube and characterized for inhibition of T cell proliferation. Triplicate determinations were made for all experiments. Mean values $\pm$ SD of counts per minute are shown and were compared by student's *t* test. To compensate for the need to perform multiple *t* tests, stringent criteria for *P* was used and only *P* < 0.005 was considered significant.

As shown in Table 1, the proliferation rate is reduced to approximately one-third if HBSS stored in a specific brand of microcentrifuge tubes (Semadeni, Ostermundigen, Switzerland)

is used in the assay (*P* < 0.001). This effect was not observed with any of the other tubes tested, including other brands of microcentrifuge tubes and larger plastic tubes. Similarly, inhibition only in this specific brand of microcentrifuge tube was confirmed in a second set of experiments using a T cell line from another patient. Consistent with these results, the inhibitory effect of an inhibitor of lymphocyte proliferation could be reduced but not eliminated by serial dilutions in microcentrifuge tubes from Semadeni.

A T lymphocyte inhibitory substance currently under investigation in our laboratory shows over 99% inhibition of proliferation at a tenfold dilution. The inhibitory activity gradually decreased to 60% inhibition when serial dilutions were performed up to 500-fold dilution. However, additional dilutions did not further diminish the inhibitory activity. Washing Semadeni

Table 1. Effect of Brand of Tube on T Cell Proliferation

Manufacturer or Distributor	Composition	Proliferation or cpm, mean $\pm$ sd
Glass tube (negative control)	Glass	20 076 $\pm$ 162
Eppendorf 1.5 mL Hamburg, Germany	PP	32 560 $\pm$ 6495
Treff 1.5 mL Winiger; Wohlen, Switzerland	PP	23 924 $\pm$ 1227
Biocon 1.5 mL Winiger; Wohlen, Switzerland	PP	25 115 $\pm$ 2644
Winiger Safelock 1.5 mL Wohlen, Switzerland	PP	21 289 $\pm$ 665
Semadeni 1.5 mL Ostermundigen, Switzerland	PP	6464 $\pm$ 2680
Nunc cryotube 2 mL Nalge Nunc Int'l, Roskilde, Denmark	PP	21 027 $\pm$ 2667
Sarstedt 2 mL Nümbrecht, Germany	PP	18 804 $\pm$ 865
Winiger 2 mL (store brand)	PP	19 714 $\pm$ 1061
Falcon® 5 mL Becton-Dickinson; Franklin Lakes, NJ, USA	PS	20 306 $\pm$ 422
Falcon 15 mL	PS	18 149 $\pm$ 1611

HBSS was stored for 30 min at room temperature in tubes of the respective manufacturer and subsequently used in a T cell proliferation assay. Proliferation was determined by ³H-deoxythymidine incorporation. PP = polypropylene; PS = polystyrene; P < 0.001 compared with glass tubes.

microcentrifuge tubes three times with distilled water before autoclaving did not improve the proliferative response outcome for HBSS (33.1% compared to glass tubes, $P < 0.001$).

We conclude that this brand of microcentrifuge tube is not suited for certain cell biology experiments. This tube appears to interfere with cell growth and proliferation in vitro. At present, it is uncertain whether plastic softeners or residual components of the manufacturing process cause this problem. Thus, in addition to factors commonly associated with poor cell growth, such as expired shelf life of nutrients and media or altered compositions of different batches of serum, the brand of microcentrifuge tube used for storage of growth factors and other additives may need to be considered for some cell biology experiments.

REFERENCES

1. Christ, A.D., S.P. Colgan, S.P. Balk and R.S. Blumberg. 1997. Human intestinal epithelial

cell lines produce factor(s) that inhibit CD3-mediated T-lymphocyte proliferation. *Immunol. Lett.* 58:159-165.

2. Probert, C.S., A.D. Christ, L.J. Saubermann, J.R. Turner, A. Chott, D. Carr-Locke, S.P. Balk and R.S. Blumberg. 1997. Analysis of human common bile duct-associated T cells: evidence for oligoclonality, T cell clonal persistence, and epithelial cell recognition. *J. Immunol.* 158:1941-1948.

Address correspondence to Dr. Andreas D. Christ, Gastroenterology Division, University Hospital, CH-8091 Zurich, Switzerland. e-mail: andreas.christ@dim.usz.ch

Received 11 January 2000; accepted 16 March 2000.

Y. Wei, R.J. Mersny¹, R.S. Blumberg², M. Fried and A.D. Christ

*University Hospital
Zurich, Switzerland*

¹*Genentech*

South San Francisco, CA

²*Brigham and Women's Hospital
Boston, MA, USA*