the interpretation that the anti-BrdU staining seen corresponds to Bromo-Uridine that has been incorporated into nascent transcripts and not incorporation of BrUTP into DNA or nonspecific binding within the cells.

Figure 1, panel I, shows the Bromo-Uridine staining pattern on polytene chromosomes in a salivary gland cell that has been squashed to spread the chromosomes. Approximately 250 distinct bands or sites of Bromo-Uridine incorporation can be made out along with other staining that is less distinct. There is a wide range of staining intensities along the chromosomes, most likely indicating that transcription is occurring at different rates at the various sites and/or that there are multiple genes being transcribed at a particular site. In Figure 1, panel J, salivary glands were incubated in TB1 containing 1 mM 20-hydroxyecdysone (ecdysone) for 1 h before labeling for 20 min with BrUTP (in the presence of ecdysone). Ecdysone is known to induce six "early" developmental puffs. The two largest puffs occur at 74EF (a site that encodes an *ets*-related transcription factor) and 75B (a site that encodes a steroid receptor DNA-binding protein). These puffs are easily identifiable and show a tremendous amount of Bromo-Uridine incorporation.

The results presented here clearly demonstrate that nascent transcripts can be visualized on polytene chromosomes of *Drosophila* through Bromo-Uridine incorporation. The level of resolution of transcript localization is unsurpassed compared to other methods, and in many instances, the fluorescent band observed on the chromosomes can be attributed to the transcription of a single gene. Using antibodies to detect RNA/DNA hybrids offers similar resolution, but the BrUTP labeling method has the added advantage of being able to visualize nascent transcripts, thus making it possible to compare global transcription patterns before and after a specific gene-inducing event. However, it does have the disadvantage of having to add BrUTP to perform the experiment. The BrUTP labeling method may also be suitable for pulse-chase experiments. In addition, the reagents used to perform these experiments are readily available through commercial sources.

REFERENCES

1. Bonner, J.J. 1981. Induction of *Drosophila* heat-shock puffs in isolated polytene nuclei. *Dev. Biol.* 86:409-418.
2. Bonner, J.J. and M.L. Pardue. 1977. Polytene chromosome puffing and in situ hybridization measure different aspects of RNA metabolism. *Cell* 12:227-234.
3. Haukenes, G. and K.-H. Kalland. 1998. Labelling of eukaryotic transcripts with BrUTP using a non-liposomal transfection reagent. *Biochemica* 3:45-46.
4. Haukenes, G., A.M. Szilvay, K.A. Brokstad, A. Kanestrom and K.H. Kalland. 1997. Labeling of RNA transcripts of eukaryotic cells in culture with BrUTP using a liposome transfection reagent (DOTAP). *BioTechniques* 22:308-312.
5. Jackson, D.A., A.B. Hassan, R.J. Errington and P.R. Cook. 1993. Visualization of focal sites of transcription within human nuclei. *EMBO J.* 12:1059-1065.
6. Pardue, M.L., S.A. Gerbi, R.A. Eckardt and J.G. Gall. 1970. Cytological localization of DNA complementary to ribosomal RNA in polytene chromosomes of diptera. *Chromosoma* 29:268-290.
7. Pelling, C. 1959. Chromosomal synthesis of ribonucleic acid as shown by incorporation of uridine labeled with tritium. *Nature* 184:655-656.
8. Rudkin, G.T. and B.D. Stollar. 1977. High resolution detection of DNA-RNA hybrids in situ by indirect immunofluorescence. *Nature* 265:472-473.
9. Wansink, D.G., W. Schul, I. van der Kraan, B. van Steensel, R. van Driel and L. de Jong. 1993. Fluorescent labeling of nascent RNA reveals transcription by RNA polymerase II in domains scattered throughout the nucleus. *J. Cell Biol.* 122:283-293.
10. Weaver, R.F. 1999. *Molecular Biology*, p. 268-272. WCB McGraw-Hill, New York.
11. Westwood, J.T., J. Clos and C. Wu. 1991. Stress-induced oligomerization and chromosomal relocalization of heat-shock factor. *Nature* 353:822-827.

This work was supported by a grant from NSERC Canada to J.T.W. Address correspondence to Dr. J. Timothy Westwood, Department of Zoology, University of Toronto at Mississauga, 3359 Mississauga Rd., Mississauga, ON, Canada L5L 1C6. e-mail: twestwoo@credit.erin.utoronto.ca

Received 4 February 2000; accepted 2 August 2000.

W.Y. Chang, N.A. Winegarden, J.P. Paraiso, M.L. Stevens and J.T. Westwood
*University of Toronto
Mississauga, ON, Canada*

Microarray-Based Detection of Select Cardiovascular Disease Markers

BioTechniques 29:936-944 (November 2000)

Assessment of the genetic risk for cardiovascular disease is complex. Genes linked to hypertension and/or thrombophilia are among the key components. Both the angiotensinogen (8) and α -adducin (3) genes play a role in hypertension. An adenine to guanine mutation in the promoter region at nucleotide-6 of the angiotensinogen gene (AGT-6) alters the binding of the nuclear protein, which results in increased gene transcription and elevated plasma angiotensinogen levels (7). α -Adducin (ADD) regulates sodium-potassium transport in the renal tubule through modifications in the actin cytoskeleton; individuals with a point mutation (G460T) in the ADD gene are reported to be more sensitive to changes in sodium balance (4). Proteolytic inactivation of factor V by activated protein C is a critical regulatory step that limits clot formation during activation of the coagulation cascade. A point mutation (Leiden) in the factor V gene (G1,691A) predicts the generation of an abnormal factor V molecule in which arginine at amino acid position 506 is replaced by glutamine. This mutation (FVL) renders factor V resistant to proteolytic inactivation by activated protein C and shifts the coagulation cascade toward one of clot formation (i.e., the clinical condition of thrombophilia) (2).

DNA microarray technology provides an ideal platform for investigating the combination of polymorphisms and mutations described above. We recently reported a strategy based on the combined use of multiplex PCR/ligase detection reaction (PCR/PCR/LDR) with "zip-code" hybridization to DNA probes on glass slides for the simultaneous detection of such genetic variants (5,6). This approach provides for an accurate, inexpensive and high-throughput assay that does not exhibit false-positive or false-negative signals, thus making it highly suitable for gene-based testing in high-incidence, low-complexity diseases.

Benchmarks

Table 1. Oligonucleotide Primers

PCR Primers (5'@3'): Boldface sequences represent universal primers.

ADD Forward: **GGAGCACGCTATCCCGTTAGAC**GGCACTTGCTCCCCACTCAGA

ADD Reverse: **CGCTGCCAACTACCGCACATC**GCAGACACCGGACGAGAGAGAC

AGT-6 Forward: **GGAGCACGCTATCCCGTTAGAC**GGCTCACTCTGTTTCAGCAGTGAAAC

AGT-6 Reverse: **CGCTGCCAACTACCGCACATC**CTCCTAGCCACAGCTCAGTTAC

FVL Forward: **GGAGCACGCTATCCCGTTAGAC**AACACCATGATCAGAGCAGTTCAAC

FVL Reverse: **CGCTGCCAACTACCGCACATC**CTAGCCAGAAGAAATTCTCAGAATTTTC

Universal Primer A: **GGAGCACGCTATCCCGTTAGAC**

Universal Primer B: **CGCTGCCAACTACCGCACATC**

LDR Primers: Boldface sequences represents zip-code complements; underlined bases indicate the locations of the polymorphisms.

ADD discriminating primer T: Cy3-CGACGAAGCTTCCGAGGAAT

ADD discriminating primer G: Cy5-CGACGAAGCTTCCGAGGAAG

ADD common primer-cZip13: p-GGCAGAATGGAAGCAGTCCCAAC**CGCACGATAGGTGGTCTACCGCTG**

AGT-6 discriminating primer G: Cy3-CATCGTGACCCGGCC

AGT-6 discriminating primer A: Cy5-CATCGTGACCCGGCC

AGT-6 common primer-cZip1: p-GGGGAAGAAGCTGCCGTTGT**GCTGAGGTGCGATGCTGAGGTGCGA**

FVL discriminating primer G: Cy3-GCAGATCCCTGGACAGGC

FVL discriminating primer A: Cy5-GCAGATCCCTGGACAGGC

FVL common primer-cZip25: p-AGGAATACAGGTATTTTGTCTTGAAGTAAC**GGTCTACCTACCCGCACGATGGTC**

Oligonucleotides were obtained as custom synthesis products from IDT (Coralville, IA, USA) or synthesized in-house on a Model 394 DNA Synthesizer (Applied Biosystems).

Genomic DNA is amplified by a two-step PCR using locus-specific primers containing universal tags on the 5' ends. The initial PCR step is directed by the locus-specific primers. This is followed by a second PCR step directed by the universal primers (experimental details below). This strategy, referred to as multiplex PCR/PCR, minimizes differences in amplification efficiency of each gene in the multiplex format (1). The resulting PCR products are then subjected to multiplex LDR, which creates a series of products that discriminates polymorphic sequences. This is achieved using the enzyme *Tth* ligase, which seals the nick between two adjacent oligonucleotides annealed to a complementary target only if the nucleotides are perfectly base-paired at the junction. Ligase reactions are ideal

for multiplexing because several primer sets can ligate along a gene without interference. A unique zip-code complement is affixed to the 3' end of each common LDR primer. Zip-codes are oligonucleotides, 24 bases in length, that have been designed to have closely matched melting temperature (T_m) values (70°C–82°C) and minimal sequence homology with the natural genome. During hybridization, the zip-code complement guides generated LDR product to the corresponding zip-code attached covalently to the prepared surface (described below). Since the T_m of a correct zip-code-mediated hybridization is at least 24°C higher than that of any incorrect hybridization (calculated using Oligo 6.0; Molecular Biology Insights, Cascade, CO, USA), cross-hybridization is prevented. De-

tection of hybridized LDR product is accomplished by the visualization of a specific fluorophore, either Cy3 or Cy5, attached to the 5' end of the discriminating primer (Figure 1). Discrimination among the polymorphisms is based on the spatial coordinates and identity of the visible fluorophore. Thus, one zip-code is associated with one gene, but two fluorophores can be distinguished at one zip-code location on the solid support. It should be pointed out that the PCR/PCR/LDR used in the analysis of the ADD, AGT-6 and FVL genes appends the zip-code complement to the 3' end of the common primer and thus doubles throughput with respect to the detection strategy described in our original paper (6), wherein the position of the zip-code complement is reversed.

Benchmarks

array and orient the viewer. (Note: zip-code complements cannot hybridize to the fiducials.)

Hybridizations of the LDR products to the microarrays are carried out in 300 mM MES, pH 6.0, 0.1% SDS, 10 mM MgCl₂ for 1 h at 65°C with gentle rotational mixing. Following hybridization, the arrays are washed briefly using 300

mM bicine, pH 8.0, 0.1% SDS, 10 mM MgCl₂ for 10 min at 50°C (6). Fluorescent signal is detected using a ScanArray[®] 5000 (GSI Lumonics, Watertown, MA, USA) with two lines (one exciting at 543.5 nm, Cy3, and the other exciting at 632.5 nm, Cy5). The use of zip-codes enables hybridizations to be carried out in a minimum amount of time (1 h) with no need for stringent washing procedures, while still producing fluorescent signals that are strong (signal-to-noise ratio of > 25:1) and specific.

Multiplex PCR/PCR/LDR followed by zip-code hybridization on the microarray for the three gene markers described above was performed on 10 DNA samples derived from a variety of cultured cell lines. Resultant genotypes for all samples are listed in Table 2; results are in complete accord with data based on conventional gel technology. An example of a hybridized array is shown in Figure 3.

In summary, we report an assay whereby mutations and polymorphisms are resolved at the amplification step (multiplex PCR/PCR combined with LDR), with a separate detection step involving the use of zip-codes to localize the high-fidelity products. We conclude that this approach produces a fluorescent signal that accurately reports the mutation (does not suffer from the occurrence of false positives or false negatives) with a signal-to-noise ratio of greater than 25:1. While the study described above only reports the results from three genes, this strategy provides the methodology to analyze polymor-

phisms or mutations in multiple (50 or more) cardiovascular genes simultaneously in an unambiguous, straightforward and rapid manner.

REFERENCES

1. Belgrader, P., M.M. Marino, M. Lubin and F. Barany. 1996. A multiplex PCR-ligase detection reaction assay for human identity testing. *Genome Sci. Technol.* 1:77-87.
2. Bertina, R.M., B.P.C. Koeleman, T. Koster, F.R. Rosendaal, R.J. Dirven, H. de Ronde, P.A. van der Velden and P.H. Reitsma. 1994. Mutation in blood coagulation factor V associated with resistance to activated protein C. *Nature* 369:64-67.
3. Casari, G., C. Barlassina, D. Cusi, L. Zagato, R. Muirhead, M. Righetti, P. Nembri, K. Amar et al. 1995. Association of the α -adducin locus with essential hypertension. *Hypertension* 25:320-326.
4. Cusi, D., C. Barlassina, T. Azzani, G. Casari, L. Citterio, M. Devoto, N. Glorioso, C. Lanzani et al. 1997. Polymorphisms of α -adducin and salt sensitivity in patients with essential hypertension. *Lancet* 349:1353-1357.
5. Favis, R., J. Day, N.P. Gerry, C. Phelan, S.A. Narod and F. Barany. 2000. Multiplex PCR/PCR/LDR using a universal DNA array detects small insertions/deletions in BRCA1 and BRCA2. *Nat. Biotechnol.* 18:561-564.
6. Gerry, N.P., N.E. Witowski, J. Day, R.P. Hammer, G. Barany and F. Barany. 1999. Universal DNA microarray method for multiplex detection of low abundance point mutations. *J. Mol. Biol.* 292:251-262.
7. Inoue, I., T. Nakajima, C.S. Williams, J. Quackenbush, R. Puryear, M. Powers, T. Cheng, E.H. Ludwig et al. 1997. A nucleotide substitution in the promoter of human angiotensinogen is associated with essential hypertension and affects basal transcription in vitro. *J. Clin. Invest.* 99:1786-1797.
8. Jeunemaitre, X., F. Soubrier, Y.V. Kotelevtsev, R.P. Lifton, C.S. Williams, A. Charru, S.C. Hunt, P.N. Hopkins et al. 1992. Molecular basis of human hypertension: role of angiotensinogen. *Cell* 71:169-178.

We gratefully acknowledge helpful discussions with Dr. Francis Barany and Susie LaFreniere. This work was supported in part by a grant from PE Biosystems (to F.B. and G.B.), the University of Minnesota Graduate School New Research Initiatives Program (to G.B. and R.C.M.) and by the Hypertension Genetic Epidemiology Network (HyperGEN) NIH/NHLBI grant no. 1 U10 HL54472. Address correspondence to Dr. George Barany, Distinguished McKnight University Professor, Department of Chemistry, University of Minnesota, 207 Pleasant Street SE, Minneapolis, MN 55455-0431, USA. e-mail: barany@umn.edu

Figure 2. Array preparation. With the aid of a mechanical spotter, nine unique zip-code oligonucleotides are deposited and subsequently coupled to an activated copolymer matrix. A quartet of spots constitutes one address. Spot diameters are approximately 170 nm with centers at approximately 250 nm apart. Amf is used to create three fiducials. Zip1 detects LDR product for AGT-6, Zip13 for ADD and Zip25 for FVL.

Figure 3. Scanner analysis of DNA specimen 7. (Left panel) Cy3 (red) is visible for three addresses, Zips1, 13 and 25. (Right panel) Cy5 (blue) is visible only at Zip13. The green quartets represent the amf fiducials overlaid on the image to facilitate orientation.

Benchmarks

Received 30 May 2000; accepted 2 August 2000.

N.E. Witowski, C. Leindecker-Foster, N.P. Gerry¹, R.C. McGlennen and G. Barany

*University of Minnesota
Minneapolis, MN*

¹*Joan and Sanford I.*

*Weill Medical College
of Cornell University
New York, NY, USA*

Nonisotopic, PCR-Based Method for 5¢ Insulin Gene VNTR Allele Class Assignment

BioTechniques 29:944-948 (November 2000)

The identification of genes that contribute to the susceptibility to polygenic human diseases is a laborious task that requires the analysis of multiple polymorphic markers in search of variants putatively associated with the disease phenotype. Besides, it is often necessary to study large cohorts of affected and unaffected individuals to achieve acceptable levels of significance. Once preliminary evidence of association has been obtained, a confirmation of results in independent samples from different populations is also required. If this type of research is to be cost effective, genotyping methods must be straightforward, easily scalable to large numbers of samples and results must be obtained in a short time.

A clear example of a polygenic disease is insulin-dependent (type 1) diabetes mellitus, an autoimmune disorder in which T-cell-mediated destruction of the insulin-producing pancreatic beta cell occurs, and is currently the object of intense genetic investigation in the search for susceptibility genes. During the last few years, several groups have focused their attention towards the allel-

ic variation at the polymorphic region situated 5¢ upstream of the human insulin gene (*INS*) on chromosome 11p15.5, which arises from a variable number of tandem repetitions (VNTR) of a 14–15 bp oligonucleotide with the consensus sequence 5¢-ACAGGGGT(G/C)(T/C)-GGGG-3¢. Based on Southern blot analysis (2), these VNTR alleles fall into three distinctive size classes according to the number of repeats of the oligonucleotide (classes I, II and III average 570, 1200 and 2200 bp, respectively).

It has been shown that in Caucasians, class I alleles are positively associated with type I diabetes, class II alleles are rare and class III alleles are dominantly protective. More recently, a PCR-based method has been used for class I allele subtyping, and allele-specific effects on disease susceptibility have been described (3,8). However, all reported methods for PCR amplification of the *INS*-VNTR polymorphism are limited to the analysis of class I alleles and fail to amplify either class II or class III alleles (3,5,6) probably because of (i) the increased length of these alleles, (ii) the strong secondary structure derived from the high GC-rich content of the template and (iii) the high frequency of class I/class III heterozygous in which the amplification of the short class I alleles is strongly favored over the long class III alleles. In fact, long alleles may only be detected by traditional Southern blot hybridization with an internal radioactive-specific probe, and this is a laborious, time-consuming technique unsuitable for large population studies requiring large amounts of purified, high molecular weight DNA, which is not always available.

The analysis of diallelic markers in the 5¢ region of the insulin gene, especially the -23/*HphI* RFLP, has proven useful as an alternative to Southern blot typing of the *INS*-VNTR, at least in populations of European descent, in which the VNTR locus may be considered diallelic (class I/class III) and tight linkage disequilibrium between -23/*HphI* and VNTR alleles has been demonstrated (4). However, it might be necessary to investigate both the frequency of class II alleles and the degree of linkage disequilibrium between the proposed marker and the *INS*-VNTR locus when a non-Caucasian popula-

tion is investigated for the first time. We report a set of PCR conditions that allow efficient amplification of *INS*-VNTR alleles of all classes, offering a nonradioactive and rapid alternative to the conventional, tedious Southern blot technique for reliable VNTR typing in large samples from any population.

Genomic DNA was isolated from blood samples using NucleoSpin[®] Blood kit (Macherey-Nagel, Düren, Germany). DNA (150 ng) was subjected to PCR amplification with flanking primers (5) (5FP1: 5¢-CACCTTGGCC-CATCCATGGCGGCATC-3¢ and 5FP2: 5¢-CTCCAGGAGAGCAAAGCCCTCACCTG-3¢, which anneal 21 and 64 bp beyond the 5¢ and 3¢ ends of the VNTR, respectively). The downstream primer is labeled with fluorochrome Cy5 on its 5¢ end, so the pair of primers is routinely used for *INS*-VNTR class I subtyping on an ALFexpress[®] automatic DNA sequencing apparatus (Amersham Pharmacia Biotech, Uppsala, Sweden). PCR amplifications were performed in 100-µL reactions containing NH₄-based PCR buffer (16 mM ammonium sulfate, 67 mM Tris-HCl, pH 8.8, and 0.01% Tween[®] 20), 2 mM MgCl₂, 10% glycerol, 0.4 mM

Figure 1. Example of *INS*-VNTR typing by PCR in a family, showing the ability of the method to detect class I, II and III alleles. Samples are run on a denaturing 3.5% polyacrylamide gel, and bands are visualized by silver staining. Lane M is the DNA Molecular Weight Marker III (Roche Diagnostics S.L., Basilea, Switzerland).