Short Technical Reports

for full-length protein (6,10) results in a powerful tool to rapidly isolate mutants for biochemical or genetic studies.

REFERENCES

1. **Duttweiler, H.M.** 1996. A highly sensitive and non-lethal β -galactosidase plate assay for yeast. *Trends Genet.* 12:340-341.
2. **Fields, S. and O. Song.** 1989. A novel genetic system to detect protein-protein interactions. *Nature* 340:245-246.
3. **Gietz, R.D. and R.A. Woods.** 1994. High efficiency transformation with lithium acetate, p. 121-134. *In* J.R. Johnston (Ed.), *Molecular Genetics of Yeast: A Practical Approach*. IRL Press, Oxford.
4. **Liang, J., S.R. Jaffrey, W. Guo, S.H. Snyder and J. Clardy.** 1999. Structure of the PIN/LC8 dimer with a bound peptide. *Nat. Struct. Biol.* 6:735-740.
5. **Puthalakath, H., D.C.S. Huang, L.A. O'Reilly, S.M. King, and A. Strasser.** 1999. The pro-apoptotic activity of the Bcl-2 family member Bim is regulated by interaction with the dynein motor complex. *Mol. Cell* 3:287-296.
6. **Shih, H.M., P.S. Goldman, A.J. DeMaggio, S.M. Hollenberg, R.H. Goodman, and M.F. Hoekstra.** 1996. A positive genetic selection for disrupting protein-protein interactions: identification of CREB mutations that prevent association with the coactivator CBP. *Proc. Natl. Acad. Sci. USA* 93:13896-13901.
7. **Tochio, H., S. Ohki, Q. Zhang, M. Li, and M. Zhang.** 1998. Solution structure of a protein inhibitor of neuronal nitric oxide synthase. *Nat. Struct. Biol.* 5:965-969.
8. **Vidal, M.** 1997. The reverse two hybrid system, p. 109-143. *In* P. Bartel and S. Fields (Eds.), *The Two Hybrid System*. Oxford University Press, Oxford.
9. **Vidal, M., R.K. Brachmann, A. Fattaey, E. Harlow, and J.D. Boeke.** 1996. Reverse two-hybrid and one-hybrid systems to detect dissociation of protein-protein and DNA-protein interactions. *Proc. Natl. Acad. Sci. USA* 93:10315-10320.
10. **Vidal, M. and P. Legrain.** 1999. Yeast forward and reverse 'n'-hybrid systems. *Nucleic Acids Res.* 27:919-929.
11. **Wach, A., H. Pick, and P. Phillipsen.** 1994. Procedures for isolating yeast DNA for different purposes, p. 1-16. *In* J.R. Johnston (Ed.), *Molecular Genetics of Yeast: A Practical Approach*. IRL press, Oxford.

We thank K. Newton for comments on the manuscript and M. Vidal for generous gifts of reagents. A.S. is a Scholar and D.C.S.H. a Special Fellow of the Leukemia and Lymphoma Society. This work was supported by the National Health and Medical Research Council (Canberra; Reg. Key 973002), the U.S. National Cancer Institute (grant no. CA80188), the Dr. Josef Steiner Cancer Foundation, the Cancer Research

Institute, and the Sylvia and Charles Viertel Charitable Foundation. Address correspondence to Dr. David Huang, The Walter and Eliza Hall Institute of Medical Research, PO Royal Melbourne Hospital, Melbourne, Victoria 3050, Australia. e-mail: huang_d@wehi.edu.au

Received 7 September 2000; accepted 21 November 2000.

Hamsa Puthalakath, Andreas Strasser, and David C.S. Huang
*The Walter and Eliza Hall Institute
Melbourne, Australia*

Screening for Ribosomal-Based False Positives Following Prokaryotic mRNA Differential Display

BioTechniques 30:988-996 (May 2001)

ABSTRACT

Differential display (DD) and the closely related RNA arbitrarily primed PCR (RAP-PCR) have become the molecular tools of choice for identifying and isolating differentially expressed genes in both eukaryotic and prokaryotic systems. However, one of the current drawbacks of both techniques is the high number of false positives generated. In prokaryotic applications, the many false positives typically generated by DD are subsequently identified as rRNAs because of their greater abundance compared to mRNAs. To circumvent this problem, full-length 16S and 23S rDNA probes, derived from Pseudomonas putida G7 and Pseudomonas aeruginosa FRD1, respectively, were used as a pre-screening approach to discriminate between those bands, which appear to be differentially expressed mRNAs, but in fact are rRNAs, following prokaryotic mRNA DD.

Short Technical Reports

INTRODUCTION

Differential gene expression is a central component common to all biological systems that permits proper functioning under normal physiological conditions or adaptation during periods of prolonged stress or disease. The differential display (DD) (8) and RNA arbitrarily primed PCR (RAP-PCR) (10, 14) procedures have provided both sensitive and versatile approaches for identifying differentially expressed genes.

Our laboratory has continued to use prokaryotic mRNA DD as a tool to monitor bacterial gene expression patterns in response to a number of environmental cues, primarily chemical exposure to heavy metals and radionuclides. Although various technical advances (13) have been employed to refine DD (and to a lesser extent RAP-PCR), one of the continued drawbacks of each approach is the generation of false positives (i.e., PCR-based cDNA products that appear to be differentially expressed on the gel but are not verified as such following northern blot or RNA slot blot analysis). In our laboratory, most false positives generated from mRNA prokaryotic DD are rRNA-derived PCR products. This is attributed primarily to the lack of poly (A)⁺ tails in prokaryotes (2,12) that prevent selective amplification of the mRNA pool. This is acutely apparent when arbitrary primers are used during the RT step because rRNAs comprise the majority (approximately 95%) of the total cellular RNA pool. To solve this problem, several current approaches have been employed (4,11) as a means of discriminating between mRNA-derived PCR products and rRNA-derived PCR products, before or following DD. The first approach is to derive the primer set(s) from the model organism of interest specifically for the mRNA pool. However, known sequence data must be available for the organism, and computer software must be available for statistical analysis to generate the mRNA-derived primer(s) of interest. An alternative approach is to use the bacterial transcriptional inhibitor, rifampicin, originally developed by our laboratory as a preliminary screening approach for detecting mRNAs on the DD gel. However, the primary drawback of this approach is its applicability only to gram-negative and

not gram-positive bacteria, which because of their thick cell membranes, prevent rifampicin from effectively penetrating into bacterial cytoplasm (12).

In this report, we adopted a simple and reliable screening approach for identifying ribosomal-derived false positives following prokaryotic mRNA DD. This entails the use of full-length 16S and 23S rDNA PCR products as antisense probes to screen putative differentially expressed cDNAs following prokaryotic mRNA DD. Such an approach should serve as a surrogate for other screening approaches that may not be applicable to the experimental organism of interest.

MATERIALS METHODS

Culture Conditions and RNA Extraction Protocol

Pseudomonas putida G7 was grown overnight at 30°C in a yeast extract peptone glucose (YEPG) medium. Before sterilization, the medium was adjusted to pH 7.2. Following overnight incubation, a 2-mL aliquot culture was transferred to 98 mL fresh YEPG media and grown to an A₆₀₀ of approximately 0.3. At this point, 15 mL culture were added to four separate conical tubes. This was followed by the addition of 200 µM CdCl₂ to two of the cultures. Bacterial cells from the cadmium-treated and control cultures were harvested following a 2-h incubation and processed for total RNA extraction. Total cellular RNA was isolated using a modified hot phenol RNA extraction method (5). Total RNA was precipitated at -20°C and subsequently quantified and diluted to approximately 1 µg/µL for DD experiments.

Reverse Transcription and Polymerase Chain Reaction (RT-PCR) for DD

For first-strand cDNA synthesis, 2 µg of total RNA were reverse transcribed (11) using two arbitrary 14-mers: Pp-2: 5'-GGTACTAAGGCTGC-3' and Pp-3: 5'-CTGCTTGATGGAGC-3'. One-tenth of the RT reaction was used for second-strand synthesis (11), using a degenerate primer designed from the Shine Dalgarno (SD) sequence (5'-GGGGAACGAC-

GATG-3') (5) and the arbitrary primer used for first-strand synthesis. Following second-strand synthesis, the PCR products were resolved on a 4.5% denaturing polyacrylamide gel (Beckman Coulter, Fullerton, CA, USA) to visualize the RNA fingerprint. Those candidate bands, which were uniquely expressed or showed a definitive differential expression pattern, were excised from the gel, re-amplified using the same DD-PCR conditions, and subjected to electrophoresis on a 1% agarose gel to ensure that the correct molecular weight band had been amplified.

Isolation of Full-Length 16S and 23S rRNAs Using Touchdown PCR (TD-PCR)

Amplification of the 16S rRNA

Figure 1. DD gel following exposure to cadmium. Arrows depict putative differentially expressed cDNAs isolated from the gel. Duplicate reactions were performed using the primer combinations Pp-3 + SD: lane 1, control 1; lane 2, Cd2⁺-exposed 1; lane 3, control 2; and lane 4, Cd2⁺-exposed 2. Bands excised from the gel (indicated by the arrows) were designated as follows: band 1, Pp-3 Cd no. 1; band 2, Pp-3 Cd nos. 2 and 3; Pp-3 Cd no. 3; band 4, Pp-3 Cd no. 4; band 5, Pp-3 control no. 1; band 6, Pp-3 control no. 2.

Short Technical Reports

gene from *P. putida* G7 was performed using degenerate primers denoted as 27f (5'-AGAGTTTGATCMTGGCTCA-3') and 1492r (5'-TACGGYTACCTTGTTACGACTT-3') (7). We used the following TD-PCR protocol (6,15): initial denaturing step at 94°C for 5 min, followed by 94°C for 45 s, 60°C for 45 s, and 72°C for 1 min, for a series of 15 cycles, with a decrement of 1°C per cycle. This was followed by a series of 25 cycles at 94°C for 45 s, 45°C for 45 s, 72°C for 1 min, and a final extension step at 72°C for 7 min. Amplification of the 23S rRNA gene from *P. aeruginosa* FRD1 was performed using a specific set of primers derived from *P. aeruginosa* (GenBank® accession no. Y00432): 23f (5'-GGTCAAGTGAAGAAGCGCATACGG-3') and 23r (5'-CCGACTTCGGCCAG

GGTCCCATACC-3'). A modified TD-PCR protocol was followed: initial denaturing step at 94°C for 5 min, followed by 94°C for 1 min, 70°C for 1 min, and 72°C for 2 min, for a series of 15 cycles, with a decrement of 1°C per cycle. This was followed by a series of 25 cycles at 94°C for 1 min, 60°C for 1 min, 72°C for 2 min, and a final extension step at 72°C for 7 min. Each of the PCR products was resolved on a 1% agarose gel for size verification. The remaining reactions were purified using the QIAquick™ PCR purification kit (Qiagen, Valencia, CA, USA), and sequenced (University of Tennessee, Molecular Biology Resource Facility). Following the sequence verification, both PCR products were cloned into the TOPOTM TA Cloning® vector (Invitrogen, Carlsbad, CA, USA).

Figure 2. DNA dot blot profile of putative differentially expressed cDNAs following hybridization with full-length 16S and 23S rDNA antisense gene probes. (A) DNA dot blot profile following ³²P-labeled probing with the full-length 16S rDNA gene. (B) DNA dot blot profile following ³²P-labeled probing with the full-length 23S rDNA gene. Individual dots are labeled numerically: 1, 23S rDNA standard; 2, 16S rDNA standard; 3, Pp-3 Cd no. 1; 4, Pp-3 Cd no. 2; 5, Pp-3 Cd no. 3; 6, Pp-3 Cd no. 4; 7, Pp-3 control no. 1; 8, Pp-3 control no. 2.

Figure 3. Verification of differential expression via total RNA slot blot analysis. Total RNA (10 µg) from control and induced cells were blotted onto nylon membranes and probed with (A) putative differentially expressed cDNAs obtained by DD and (B) ³²P-labeled 16S rDNA to confirm the equal loading of total RNA. Pairs of RNA slots [control (upper) and induced (lower)] are numbered according to the cDNA probe used: 1, Pp-3 Cd no. 1; 2, Pp-3 Cd no. 2; 3, Pp-3 control no. 1; 4, Pp-3 control no. 2.

Short Technical Reports

Ribosomal Probing of Candidate cDNAs and Verification of Differential Expression

A total of 10 ng from each candidate PCR-re-amplified cDNA were resuspended in a 0.5 M NaOH solution, followed by denaturation in a boiling water bath for 5 min. Each sample was then placed in an ice-cold water bath for several minutes and then loaded onto a dot-blot manifold using an analogous protocol described by Sambrook et al. (16). Prehybridization and hybridization experiments were carried out at 75°C using a hybridization buffer, pH 7.2, consisting of 0.5 M NaH₂PO₄, 1 mM EDTA, and 7% SDS (3). For hybridization experiments, we generated [³²P]-dCTP-labeled antisense probes from 16S and 23S PCR products using the Prime-A-Probe™ kit (Ambion, Austin, TX, USA). The resulting labeled reactions were purified using a NucTrap®

column (Stratagene, La Jolla, CA, USA) and denatured in a boiling water bath for 5 min, followed by incubation in an ice-cold water bath for several minutes. Approximately $5-8 \times 10^7$ total cpm from each labeled sample were added to the appropriate bag and hybridized overnight. Following hybridization, blots were washed with 0.1% SDS, 0.1× SSPE, three times for 20 min at 75°C. Dot blots were exposed overnight to a phosphorimager intensifying screen and visualized using the Molecular Dynamics Storm 840 Imaging System (Molecular Dynamics, Sunnyvale, CA, USA).

To confirm differential expression, 10 µg total RNA was loaded and blotted onto nylon membranes using a slot-blot manifold (16). Blots were air dried, baked at 80°C for 2 h, and prehybridized with a formamide-based hybridization solution (Ambion) for 4–6 h at 45°C. The candidate differential cDNAs were labeled with [³²P]-dCTP using a

linear PCR protocol supplied with the Prime-A-Probe kit. Labeled double-stranded PCR products were purified, boiled for 5 min, added to the appropriate hybridization bag, and hybridized overnight at 45°C. This was followed by three washes at 45°C for 20 min using the same wash solution in diethylpyrocarbonate (DEPC)-treated water. Total RNA blots were exposed from one to two days, depending on the signal following the washing.

RESULTS AND DISCUSSION

Figure 1 depicts an mRNA fingerprint resolved on a DD gel following second-strand synthesis using the primer combination of Pp-3 and SD (lanes 1–4). No observable differences in gene expression were detected using the primer combination of Pp-2 and SD (data not shown). The bands indicating

a reproducible expression pattern between duplicate samples were chosen to further eliminate any potential artifacts. A total of six candidate cDNAs were identified as unique or showing putative differential regulation in either the control or the cadmium-exposed samples (denoted by the arrows). The candidate bands were excised from the DD gel, eluted in 50 μ L RNase-free water, and re-amplified using the same PCR protocol to generate adequate template for subsequent rRNA-based DNA dot blot experiments.

Two DNA dot blots were prepared by blotting 10 ng of each re-amplified PCR product onto a nylon membrane. The blots were subsequently hybridized using the full-length antisense 16S and 23S rDNA probe, respectively. Figure 2 represents the results following probing and imaging. Compared to the 23S rDNA standard, no differences in signal intensity were detected in any of the nu-

merical dot positions corresponding to each candidate cDNA of interest following hybridization with the 16S rDNA probe (Figure 2A). However, dot positions 5 and 6, corresponding to Pp-3 Cd no. 3, and Pp-3 Cd no. 4, respectively, revealed an intense hybridization signal following hybridization with the 23S rDNA probe (Figure 2B). As a further precaution, to test the validity of this screening approach, both PCR products were sequenced and analyzed using the GenBank Blast-N and Blast-X algorithms (1). As expected, both sequences revealed a 95%–97% sequence similarity to a number of 23S rRNA coding regions. Thus, both sequences could be dismissed from further analysis, and the four remaining cDNAs of interest were cloned and processed for differential expression verification experiments.

Figure 3A depicts four separate total RNA blots probed with each of the remaining candidate cDNAs. Levels of

transcriptional induction between the control- and cadmium-exposed samples were calculated using the ImageQuant™ software package version 5.0 (Molecular Dynamics). The two bands corresponding to Pp-3 Cd no. 1 and Pp-3 Cd no. 2 were up-regulated following a 2-h post-exposure to cadmium, threefold and fourfold, respectively, compared to control samples. The two bands corresponding to Pp-3 control no. 5 and Pp-3 control no. 6 were down-regulated threefold and twofold, respectively, compared to the cadmium-exposed samples. The confirmed differential cDNAs were examined using the Blast-N and Blast-X algorithms to identify putative related function(s) to other known sequences within the database. Interestingly, band Pp-3 Cd no. 1 showed the highest degree of similarity (76%) to the ArsH-resistant protein from several different organisms, suggesting that this protein is possibly involved in some form of metal resis-

Short Technical Reports

tance. Band Pp-3 Cd no. 2 showed the highest degree of similarity (72%) to putative cytoplasmic membrane-associated proteins from both *P. fluorescens* and *P. aeruginosa*. Band Pp-3 control no. 5 was identified to be the most similar (88%) to the phosphoribosylaminoimidazole carboxylase catalytic subunit involved in the purine biosynthetic pathway in *P. aeruginosa*. Band Pp-3 control no. 6 showed no significant matches to any sequence in the database, suggesting a putative novel sequence.

As demonstrated in this experiment, two of the six putative differentially expressed cDNAs displayed on the DD gel resulted in 23S rRNA-based false positives following DNA dot blot analysis. In our laboratory, this generally falls within the percentage (about 40%) of false positives generated following DD, if no prior screening method is employed. The primary advantages of this approach are (i) the application to gram-negative or gram-positive bacteria for which rifampicin may be ineffective, (ii) the application to bacteria for which sequence information may be limiting, (iii) the conservation of precious amounts of RNA template, and (iv) the employment of 16S and 23S rDNA PCR products as probes to screen cDNAs from other closely related organisms because typical evolutionary relationships show a relatively high sequence conservation (9,17).

REFERENCES

1. Altschul, S.F., T.L. Madden, A.A. Schaffer, J. Zhang, Z. Zheng, W. Miller, and D.J. Lipman. 1997. Gapped blast and PSI-blast: a new generation of protein database search programs. *Nucleic Acids Res.* 25:3389-3402.
2. Belasco, J.G. 1993. Experimental approaches to the study of mRNA decay, p. 475-493. In J.G. Belasco and G. Brawerman (Eds.), *Control of Messenger RNA Stability*. Academic Press, San Diego.
3. Church, G.M. and W. Gilbert. 1984. **Genomic sequencing**. *Proc. Natl. Acad. Sci. USA* 81:1991-1995.
4. Fislage, R., M. Berceanu, Y. Humboldt, M. Wendt, and H. Oberender. 1997. Primer design for a prokaryotic differential display RT-PCR. *Nucleic Acids Res.* 25:1830-1835.
5. Fleming, J.T., W.-H. Yao, and G.S. Sayler. 1998. Optimization of differential display to prokaryotic mRNA: application to pure culture and soil microcosms. *Appl. Environ. Microbiol.* 64:3698-3706.
6. Hecker, K.H. and K.H. Roux. 1996. High and low annealing temperatures increase both specificity and yield in touchdown and step-down PCR. *BioTechniques* 20:478-485.
7. Lane, D.J. 1991. 16S/23S rRNA sequencing, p. 115-148. In E. Stackebrandt and M. Goodfellow (Eds.), *Nucleic Acid Techniques in Bacterial Systematics*. John Wiley & Sons, New York.
8. Liang, P. and A.B. Pardue. 1992. Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction. *Science* 257:967-971.
9. Ludwig, W., R. Rossello-Mora, R. Aznar, S. Klugbauer, S. Spring, K. Reetz, C. Beimehr, E. Brockman et al. 1995. Comparative sequence analysis of 23S rRNA from proteobacteria. *Syst. Appl. Microbiol.* 18:164-188.
10. McClelland, M., R. Honeycutt, F. Mathieu-Daude, T. Vogt, and J. Welsh. 1997. Fingerprinting by arbitrarily primed PCR. *Methods Mol. Biol.* 85:13-24.
11. Nagel, A., J.T. Fleming, and G.S. Sayler. 1999. Reduction of false positives in prokaryotic mRNA differential display. *BioTechniques* 26:641-648.
12. Nierlich, D.P. and G.J. Murakawa. 1996. The decay of bacterial messenger RNA. *Prog. Nucleic Acid Res. Mol. Biol.* 52:153-215.
13. Pardue, A.B. and M. McClelland. 1999. Reducing false positives, p. 163-207. In *Expression Genetics: Differential Display*. Eaton Publishing, Natick, MA.
14. Rothschild, C.B., C.S. Brewer, and D.W. Bowden. 1997. DD/AP-PCR: combination of differential display and arbitrarily primed PCR of oligo (dT) cDNA. *Anal. Biochem.* 245:48-54.
15. Roux, K.H. 1994. Using mismatched primer-template pairs in touchdown PCR. *BioTechniques* 16:812-814.
16. Sambrook, J., E.F. Fritsch, and T. Maniatis. 1989. *Molecular Cloning: A Laboratory Manual*, 2nd ed. CSH Laboratory Press, Cold Spring Harbor, NY.
17. Woese, C.R. 1987. Bacterial evolution. *Microbiol. Rev.* 51:221-271.

This work was supported by grant no. 1081/8 from the Science and Engineering Research Development Program (SERDP). Address correspondence to Dr. James T. Fleming, The University of Tennessee, Knoxville, The Center for Environmental Biotechnology, 676 Dabney Hall, Knoxville, TN 37996, USA. e-mail: jtf@utk.edu

Received 11 July 2000; accepted 13 November 2000.

Aaron C. Nagel¹, James T. Fleming¹, Gary S. Sayler¹, and Kenneth L. Beattie^{1,2}

¹*The University of Tennessee*

Knoxville

Knoxville, TN

²*Oak Ridge National*

Laboratory

Oak Ridge, TN, USA