Short Technical Reports

- naire, and J. Mallet. 1997. Anchoring a defined sequence to the 5' ends of mRNAs. The bolt to clone rare full-length mRNAs. *Methods Mol. Biol.* 67:261-278.
6. Edery, I., L.L. Chu, N. Sonenberg, and J. Pelletier. 1995. An efficient strategy to isolate full-length cDNAs based on an mRNA cap retention procedure (CAPture). *Mol. Cell Biol.* 15:3363-3371.
7. England, T.E., A.G. Bruce, and O.C. Uhlenbeck. 1980. Specific labeling of 3' termini of RNA with T4 RNA ligase. *Methods Enzymol.* 65:65-74.
8. England, T.E. and O.C. Uhlenbeck. 1978. Enzymatic oligoribonucleotide synthesis with T4 RNA ligase. *Biochemistry* 17:2069-2076.
9. Fu, D.J., N.E. Broude, H. Koster, C.L. Smith, and C.R. Cantor. 1995. A DNA sequencing strategy that requires only five bases of known terminal sequence for priming. *Proc. Natl. Acad. Sci. USA* 92:10162-10166.
10. Goode, B.L. and S.C. Feinstein. 1994. Identification of a novel microtubule binding and assembly domain in the developmentally regulated inter-repeat region of tau. *J. Cell Biol.* 124:769-782.
11. Gubler, U. and B.J. Hoffman. 1983. A simple and very efficient method for generating cDNA libraries. *Gene* 25:263-269.
12. Hamilton, D.L. and K. Abremski. 1984. Site-specific recombination by the bacteriophage P1 lox-Cre system. Cre-mediated synopsis of two lox sites. *J. Mol. Biol.* 178:481-486.
13. Harada, K. and L.E. Orgel. 1993. In vitro selection of optimal DNA substrates for T4 RNA ligase. *Proc. Natl. Acad. Sci. USA* 90:1576-1579.
14. Hyone-Myong, E. 1996. *Enzymology Primer for Recombinant DNA Technology*, p. 477-489. Academic Press, San Diego.
15. King, R.W., K.D. Lustig, P.T. Stukenberg, T.J. McGarry, and M.W. Kirschner. 1997. Expression cloning in the test tube. *Science* 277:973-974.
16. Maruyama, K. and S. Sugano. 1994. Oligocapping: a simple method to replace the cap structure of eukaryotic mRNAs with oligoribonucleotides. *Gene* 138:171-174.
17. Okayama, H. and P. Berg. 1982. High-efficiency cloning of full-length cDNA. *Mol. Cell Biol.* 2:161-170.
18. Pesole, G., S. Liuni, G. Grillo, and C. Saccone. 1997. Structural and compositional features of untranslated regions of eukaryotic mRNAs. *Gene* 205:95-102.
19. Sambrook, J., E.F. Fritsch, and T. Maniatis. 1989. *Molecular Cloning: A Laboratory Manual*, 2nd ed. CSH Laboratory Press, Cold Spring Harbor, NY.
20. Schmidt, W.M. and M.W. Mueller. 1999. CapSelect: a highly sensitive method for 5' CAP dependent enrichment of full-length cDNA in PCR-mediated analysis of mRNAs. *Nucleic Acids Res.* 27:e31.
21. Sugahara, Y., P. Carninci, M. Itoh, K. Shibata, H. Konno, T. Endo, M. Muramatsu, and Y. Hayashizaki. 2001. Comparative evaluation of 5'-end sequence quality of clones in CAP trapper and other full-length cDNA libraries. *Gene* 263:93-102.
22. Suh, D., Y. Yuan, D. Henning, and R. Reddy. 1989. Secondary structure of 7SK and 7-2 small RNAs. Possible origin of some 7SK pseudogenes from cDNA formed through self-priming by 7SK RNA. *Eur. J. Biochem.* 186:221-226.
23. Summers, D.K. 1991. The kinetics of plasmid loss. *Trends Biotechnol.* 9:273-278.
24. Troutt, A.B., M.G. McHeyzer-Williams, B. Pulendran, and G.J. Nossal. 1992. Ligation-anchored PCR: a simple amplification technique with single-sided specificity. *Proc. Natl. Acad. Sci. USA* 89:9823-9825.

We thank A. Hara, A. Yasunishi, N. Hayatsu, K. Satoh, M. Nakamura, and Y. Hodoyama for technical assistance. This study has been supported by a Research Grant for the RIKEN Genome Exploration Research Project from the Science and Technology Agency of the Japanese Government. This work was also supported by a Grant-in-Aid for Scientific Research on Priority Areas and Human Genome Program, from the Ministry of Education, Science and Culture, and by a Grant-in-Aid for a Second Term Comprehensive 10-Year Strategy for Cancer Control from the Ministry of Health and Welfare to Y. H. Address correspondence to Dr. Piero Carninci, Genome Exploration Research Group, RIKEN Genomic Sciences Center (GSC) and Genome Science Laboratory, RIKEN Tsukuba Institute, 3-1-1 Koyadai, Tsukuba 305-0074, Japan. e-mail: rgscerg@gsc.riken.go.jp

Received 31 July 2000; accepted 12 December 2000.

**Y. Shibata^{1,2}, P. Carninci¹,
A. Watahiki^{1,3}, T. Shiraki¹,
H. Konno¹, M. Muramatsu¹,
and Y. Hayashizaki^{1,3}**
¹RIKEN Tsukuba Institute
Ibaraki
²Nippon Gene
Tonyamachi, Toyama
³University of Tsukuba
Ibaraki, Japan

RT-PCR Method to Quantify Vascular Endothelial Growth Factor Expression

BioTechniques 30:1254-1260 (June 2001)

ABSTRACT

The vascular endothelial growth factor (VEGF) is implicated in the progression of cancers. Its expression is well correlated with tumor growth and metastases. The availability of a rapid and sensitive method to detect the amounts of VEGF mRNA in biological samples of limited size, very small biopsies, or samples containing relatively few cells could provide an interesting prognostic tool for clinicians. We have developed an RT-PCR method that allows us to detect the VEGF mRNA from as little as 3 µg total mRNA. We have also shown that this protocol can be generalized to all cell lines tested. This method constitutes a very potent tool for the analysis of VEGF mRNA expression in different contexts.

INTRODUCTION

The vascular endothelial growth factor (VEGF) is one of the major angiogenic factors implicated in the neovascularization of human tumors. High levels of VEGF correlate with aggressivity and bad prognosis in human tumors (5). VEGF mRNAs are usually analyzed by classical northern blotting, which is time consuming and difficult to apply to low amounts of mRNA, or by qualitative RT-PCR (10). Quantitative RT-PCR using endogenous mRNA as control (3) has also been described. However, the level of control mRNAs classically used, such as GAPDH, actin, or HPRT, could vary under certain conditions, rendering this approach poorly reliable. We have developed a new quantitative RT-PCR method to precisely determine the levels of VEGF mRNA in cell lines and tissues of different origins. This method is based on a competitive RT-PCR using an artificial RNA as an internal standard (11). It constitutes a practical tool for researchers and clinicians.

Short Technical Reports

MATERIALS AND METHODS

Construction of the Standard

One microgram of the two following overlapping oligonucleotides (Eurogentec, Liege, Belgium) complementary to the mouse VEGF 164 A cDNA (GenBank® accession no. M95200) were annealed (Oligo 1: 5'-GTTTCAGAGCGGAGAAAGCATTGTATGTTCCTGCAAAAACACAGACTCGC-

GTT-3'; Oligo 2: 5'-CACATCTGCAAGTACGTTTCGTTTAACTCAAGCTGCCTCGCCTTGCAACGGAGTCTGTGTTTTTGCAGGA-3') and then submitted to a polymerization reaction at 72°C using *Taq* DNA polymerase (Qiagen, Valencia, CA, USA) to fill in the un-annealed region. The resulting DNA fragment of 97 bp was inserted in the pCRII-TOPO® vector (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's protocol.

The plasmid was linearized with *NotI*, and the RNA for use as internal standard was generated in vitro by T7 RNA polymerase (Promega, Madison, WI, USA). The first 23 bases of Oligo 1, representing the forward primer (Figure 1), correspond to bases 434–456 of the coding region of the mouse VEGF 164 cDNA; the 30 following bases correspond to bases 483–511 of the same cDNA. Thus, we have artificially introduced a 27-base deletion in our internal standard. Oligo 2 corresponds to bases 559–487 of the mouse cDNA. The first 23 bases of Oligo 2 represent the reverse primer (Figure 1).

RNA Quantification by RT-PCR

The cDNAs were synthesized using the Omniscript™ kit (Qiagen) according to the manufacturer's protocol. The reverse transcription (RT) reaction was carried out using 0.5 µg total RNA in the presence of variable amounts of internal standard RNA (ranging from 1 pg to 10 ng; Figure 2B) and 100 ng reverse primer 5'-CACATCTGCAAGTACGTTTCGTTT-3' in a final volume of 20 µL. The PCR was performed using the *Taq* PCR Master Mix Kit (Qiagen) with 10 µL of the RT reaction, the reverse primer, and the forward primer 5'-GTTTCAGAGCGGAGAAAGCATTG-3'. The PCR was performed in a final volume of 50 µL in the presence of 0.133 µCi α[³²P]dATP in a T3 Thermocycler (Biometra) using the following conditions: 95°C for 3 min and then 28 cycles of 95°C for 30 s, 56.5°C for 30 s, 72°C for 30 s, and finally 72°C for 7 min. Amplified fragments were resolved on 10% polyacrylamide gels (Tris-borate EDTA) and quantitated by phosphorimager analysis (Fujix Bas 1000 Model IPR 1000; Fuji Photo Film, Elmsford, NY, USA). The expected size of the target fragment is 123 bp, and the expected size of the standard is 97 bp.

RESULTS

Description of the Quantitative RT-PCR Method

Quantification of VEGF mRNA levels was performed by co-amplification

Figure 1. Detection of VEGF-A mRNA by quantitative RT-PCR. (A) Specific competitor DNA was generated by annealing two complementary oligonucleotides (nos. 1 and 2). The first 23 bases of Oligo 1 represent the forward primer (FP), and the first 23 bases of Oligo 2 represent the reverse primer (RP). Oligo 1 has a 27-base deletion in its 5' end as compared to the VEGF cDNA. The forward and reverse primers will be used in the reactions. (B) Autoradiogram of a representative competitive RT-PCR experiment. Known concentration of competitor RNA was added to sample RNA and subjected to RT-PCR. Upper fragments (123 bp) represent the signals obtained from constant amount of target RNA. Lower fragments (97 bp) represent the signals obtained from decreasing amounts of competitor RNA (lanes 1–5). Lane 6; target RNA alone. Lane 7; standard RNA alone.

Short Technical Reports

of the target VEGF mRNA and an artificial RNA standard yielding a deletion of 27 bp (see Materials and Methods). Amplification of the shorter competitor RNA, which yields a distinguishable fragment, allows for control of the RT step. To establish linearity of the assay, 0.5 μ g total RNA isolated from the tissue or from the cells to be tested is mixed with serial dilutions (1:10) of the competitor RNA (ranging from 1 pg to 10 ng) and reverse transcribed with a specific oligonucleotide for VEGF A (reverse primer) (Figure 1A). This oligonucleotide also anneals to the 3' part of the internal RNA standard. The PCR is performed with 10 μ L RT reaction, the forward and reverse primer in the presence of α [33 P]dATP (Figure 1B). Both oligonucleotides correspond to highly conserved sequences among different species (human, mouse, rat, hamster, dog, bovine, and pig) of mouse VEGF. The radioactivity incorporated in the amplified DNA fragments, either corresponding to the VEGF target or the internal standard, are measured by phosphorimager analysis (Figure 2). The variable standard concentrations were plotted against the log of the ratio between the counts incorporated in the internal standard and in the target. The straight line obtained crosses the abscissa (log 0) when the corresponding counts from the standard and the VEGF RNA are the same. This method allows us to detect VEGF mRNA amounts ranging from 10 to 30 pg. Thus, the proposed RT-PCR method permits us to quantitate VEGF mRNA expression in samples in which detection by northern blotting analysis was impossible (7), such as total RNA from serum-deprived CCL39 Chinese hamster lung fibroblasts. Our oligonucleotides can anneal to mRNAs encoding four spliced variants of VEGF A (these spliced variants include mouse VEGF 205, 188, 183, and 164 or their human homologues VEGF 206, 189, 183, and 165). Variations in VEGF 144/145 and VEGF 120/121 expression (mouse/human, respectively), two others described spliced variants cannot be monitored because their mRNAs do not contain the sequence corresponding to Oligo 1 and Oligo 2. Whereas VEGF188/189, 164/165, and 120/121

are the predominant forms expressed in most tissues and cells examined, VEGF 144/145 and 205/206 are rare variants (2). Although VEGF 188/189, 164/165, and 120/121 are co-regulated most of the time, upon certain conditions VEGF 188/189 and VEGF 120/121 are expressed at higher levels than VEGF 164/165 (1,4,8,12). Our method will not allow the detection of modifications

in VEGF 120/121 expression. To validate our RT-PCR method, we have compared the VEGF mRNA expression in non-transformed CCL39 cells stably expressing the Raf-1: ER chimera stimulated, or not, with estradiol. We have previously shown by northern blot analysis that, in this model system, estradiol treatment leads to a robust induction of VEGF mRNA expression,

Figure 2. Comparison between the quantitative RT-PCR method and a classical northern blot analysis. (A) Total RNA (20 μ g) isolated from Raf-1:ER cells quiescent (0) or estradiol-stimulated for 5 h (5H) were analyzed by northern blotting for the expression of VEGF A. Ethidium bromide staining of 18S ribosomal RNA is shown as loading control. (B) The amounts of VEGF mRNA in the samples analyzed above by northern blot were quantified by the competitive RT-PCR method. The radioactivity incorporated in the internal standard, and the target DNA fragments was quantified using a phosphorimager. Variable dilutions of the internal standard RNA were plotted against the log of the ratio between the radioactivity incorporated in the DNA fragments corresponding to the internal standard DNA fragment and those corresponding to the target VEGF DNA fragment. The heavy arrow shown on the abscissa corresponds to difference in RNA between the two conditions. (C) Estradiol-induced stimulation of VEGF A mRNA levels obtained in (B) is plotted. This result is representative of four independent experiments.

which is maximal after 5 h of stimulation (7). Thus, we have used RNA isolated from quiescent cells as a control of basal expression to determine the fold induction in VEGF mRNA after 5 h of estradiol stimulation by our RT-PCR technique. For that purpose, we compared the amounts of standard RNA necessary to obtain the crossing of the abscissa in control or estradiol-stimulated RNA samples. The ratio between the corresponding values determines the fold induction of VEGF mRNA levels between both conditions. Figure 2 compares the results obtained by northern blotting (Figure 2A) with those using RT-PCR method (Figure 2, B and C). Using the proposed quantitative RT-PCR method, it can be seen that VEGF mRNA levels are induced 10 times following estradiol stimulation. We can now easily quantify basal VEGF mRNA levels, whereas it was impossible by northern blot. These re-

sults demonstrate that our RT-PCR approach is more precise and sensitive than our previously used analysis.

Expression of VEGF mRNA in Different Cell Lines

To validate our technique, we have compared the expression of VEGF A cell lines from different species and with varying tumoral potential including serum-deprived Raf-1:ER hamster cells (as a basal control), Raf-1:ER cells stimulated for 5 h with estradiol, human colon carcinoma HT-29 cells, rat colon carcinoma PRO cells (6), and Ras-transformed CCL39 cells (9). Figure 3 shows that the highly transformed PRO cells, HT-29 cells, and Ras-transformed cells express 15- to 30-fold more VEGF mRNA than the quiescent Raf-1:ER control cells. This result correlates with their capacity to induce highly vascularized tumors.

DISCUSSION

The search for even better molecular biologic factors for cancer determination is speeding up because an efficient treatment relies on a more precise prognosis. VEGF is one of the major growth factors implicated in tumoral angiogenesis. Humanized anti-VEGF antibodies are currently being tested for their capacity to repress tumoral angiogenesis. Assessment of VEGF levels in different tumors could provide a valuable prognostic parameter for optimization of therapeutic protocols. However, to adapt their therapeutic protocols, clinicians need a sensitive and efficient mean of evaluating VEGF levels in the different tumors that they treat. This evaluation is not possible using a classical ELISA kit because of the limitation in cultivating the resected tumoral cells. ELISA can be performed on patient serum, but significant increases

Short Technical Reports

Figure 3. Analysis of VEGF A mRNA in different cell lines by quantitative RT-PCR. Total RNA isolated from quiescent Raf-1:ER cells (column 1), Raf-1:ER cells stimulated for 5 h with estradiol (column 2), exponentially growing PRO cells (column 3), HT-29 cells (column 4), and Ras-transformed cells (column 5) were analyzed by RT-PCR for VEGF mRNA amounts. These data are representative of three independent experiments.

are detectable when the tumor has reached a critical size. Our results demonstrate that we are able to precisely quantify the amounts of VEGF mRNA in various samples using only 1 µg total RNA. These very low amounts of RNA are comparable to those extracted from biopsy specimens. Beyond its importance in fundamental research, our method constitutes a practical tool for clinical evaluation to estimate the potential of tumor cells to induce highly aggressive vascularized tumors. Determination of precise VEGF amounts will enable a more tailored selection of individualized therapeutic options.

REFERENCES

1. **Bacic, M., N.A. Edwards, and M.J. Merrill.** 1995. Differential expression of vascular endothelial growth factor (vascular permeability factor) forms in rat tissues. *Growth Factors* 12:11-15.
2. **Burchardt, T., M. Burchardt, M.W. Chen, R. Buttyan, A. de la Taille, A. Shabsigh, and R. Shabsigh.** 1999. Expression of VEGF splice variants 144/145 and 205/206 in adult mouse tissues. *IUBMB Life* 48:405-408.
3. **Iizuka, M., M. Yamauchi, K. Ando, N. Hori, Y. Furusawa, H. Itsukaichi, K. Fukutsu, and H. Moriya.** 1994. Quantitative RT-PCR assay detecting the transcriptional

induction of vascular endothelial growth factor under hypoxia. *Biochem. Biophys. Res. Commun.* 205:1474-1480.

4. **Krussel, J.S., B. Behr, A.A. Milki, J. Hirschenhain, Y. Wen, P. Bielfeld, and M.L. Polan.** 2001. Vascular endothelial growth factor (VEGF) mRNA splice variants are differentially expressed in human blastocysts. *Mol. Hum. Reprod.* 7:57-63.
5. **Lee, Y.H., T. Tokunaga, Y. Oshika, R. Suto, K. Yanagisawa, M. Tomisawa, H. Fukuda, H. Nakano et al.** 1999. Cell-retained isoforms of vascular endothelial growth factor (VEGF) are correlated with poor prognosis in osteosarcoma. *Eur. J. Cancer* 35:1089-1093.
6. **Martin, M., K. Ahlen, M. Dimanche-Boitrel, D. Mendrick, D. Turner, K. Rubin, and F. Martin.** 1996. Colon-cancer cell variants producing regressive tumors in syngeneic rats, unlike variants yielding progressive tumors, attach to interstitial collagens through integrin alpha2beta1. *Int. J. Cancer* 65:796-804.
7. **Milanini, J., F. Vinals, J. Pouyssegur, and G. Pagès.** 1998. p42/p44 MAP kinase module plays a key role in the transcriptional regulation of vascular endothelial growth factor gene in fibroblasts. *J. Biol. Chem.* 273:18165-18172.
8. **Rout, U.K., K. Oomen, and M.P. Diamond.** 2000. Altered expressions of VEGF mRNA splice variants during progression of uterine-peritoneal adhesions in the rat. *Am. J. Reprod. Immunol.* 43:299-304.
9. **Seuwen, K., A. Lagarde, and J. Pouyssegur.** 1988. Dereglulation of hamster fibroblast proliferation by mutated ras oncogenes is not mediated by constitutive activation of phospho-

inositide-specific phospholipase C. *EMBO J.* 7:161-168.

10. **Tober, K.L., R.E. Cannon, J.W. Spalding, T.M. Oberyzy, M.L. Parrett, A.I. Rackoff, A.S. Oberyzy, R.W. Tennant, and F.M. Robertson.** 1998. Comparative expression of novel vascular endothelial growth factor/vascular permeability factor transcripts in skin, papilloma and carcinomas of v-Ha-ras Tg.AC transgenic mice and FVB/N mice. *Biochem. Biophys. Res Commun.* 247:644-653.
11. **Wang, A.M., M.V. Doyle, and D.F. Mark.** 1989. Quantitation of mRNA by polymerase chain reaction. *Proc. Natl. Acad. Sci. USA* 86:9717-9721.
12. **Zhang, H.T., P.A. Scott, L. Morbidelli, S. Peak, J. Moore, H. Turley, A.L. Harris, M. Ziche, and R. Bicknell.** 2000. The 121 amino acid isoform of vascular endothelial growth factor is more strongly tumorigenic than other splice variants in vivo. *Br. J. Cancer* 83:63-68.

We thank Dr. Edurne Berra and Ellen Van Obberghen for reading the manuscript. This work was supported by grants from the Centre National de la Recherche Scientifique (CNRS), l'Université de Nice Sophia-Antipolis, le Ministère de la Recherche et de la Technologie, l'Association pour la Recherche contre le Cancer (ARC), and le Groupement des Entreprises Françaises et Monégasque dans la Lutte contre le Cancer et la Fondation de France. Address correspondence to Dr. Gilles Pagès, CNRS UMR 6543, Centre Antoine Lacassagne, 33 Avenue de Valombrose, 06189 Nice, France. e-mail: gpages@unice.fr

Received 16 October 2000; accepted 8 January 2001.

Frédéric Bonino, Julie Mila-nini, Jacques Pouyssegur, and Gilles Pagès
CNRS UMR 6543
Centre Antoine Lacassagne
Nice, France