Benchmarks

used in combination with any other M13-tailed forward primer.

We then tested whether the method could be extended to analysis on an automatic capillary sequencer. Three M13 primers end-labeled with WellRED fluorescent dyes D2, D3, and D4 (Beckman Coulter, Fullerton, CA, USA) were ordered from Research Genetics (Huntsville, AL, USA). These primers were used for the amplification of labeled alleles using the same protocol described previously for ^{32}P -labeled primers. After amplification, the PCR products were pooled (i.e., up to nine different products, three size ranges, and three dyes). For desalting, wells of a MultiScreen®-FB filter plate (Millipore, Bedford, MA, USA) were pre-wetted with binding buffer [7 M guanidine-HCl and 200 mM 2-(N-morpholino) ethanesulfonic acid, pH 5.6], the pooled PCR products were then mixed in a 1:1 ratio with binding buffer and loaded onto individual wells by centrifugation at $1000\times g$ for 5 min. The wells were then washed twice with 80% ethanol, and the products were eluted with 50 μL TE buffer (0.1 mM EDTA and 10 mM Tris-HCl, pH 8.0). Fifteen microliters of eluate were mixed with 25 μL deionized formamide and 0.5 μL size standards (Beckman Coulter) labeled with WellRED fluorescent dye D, then loaded and analyzed on a CEQ™ 2000XL capillary DNA analysis system (Beckman Coulter). Figure 2 shows an analysis of a run containing six multiplexed PCR products. Genomic DNA from strain A amplified by three markers along with M13 primer end-labeled with WellRED fluorescent dye D2 is compared with genomic DNA from strain B amplified with the same markers along with M13 primer end-labeled with fluorescent WellRED dye D3. As observed with ^{32}P -labeled markers, PCR products migrate as clean-laddered fluorescence peaks without interfering artifactual bands and, thus, allow clear identification of the origin of the alleles. We have obtained similar results when multiplexing was initiated at the level of PCR, by amplifying genomic DNA with several primer combinations simultaneously. It is reasonable to anticipate that this method could be used with dyes designed for other sequencing platforms. For instance, M13 primers labeled with

FAM and HEX fluorescent dyes have already been used for other applications involving the analysis of fragments on an ABI PRISM™ 377 automated DNA sequencer (Applied Biosystems, Foster City, CA, USA) (1).

In summary, the use of M13-tailed primers has the potential to facilitate microsatellite analysis, regardless of whether allele sizing is performed on manual gels or on automatic sequencers. The benefits of using M13-tailed primers include the possibility of increasing the genotyping output with greater convenience, reducing costs, and increasing specificity.

REFERENCES

1. **Boutin, P., E.H. Hani, F. Vasseur, C. Roche, B. Bailleul, J. Hager, and P. Froguel.** 1997. Automated fluorescence-based screening for mutation by SSCP: use of universal M13 dye primers for labeling and detection. *BioTechniques* 23:358-362.
2. **Oetting, W.S., H.K. Lee, D.J. Flanders, G.L. Wiesner, T.A. Sellers, and R.A. King.** 1995. Linkage analysis with multiplexed short tandem repeat polymorphisms using infrared fluorescence and M13 tailed primers. *Genomics* 30:450-458.
3. **Steffens, D.L., S.L. Sutter, and S.C. Roemer.** 1993. An alternate universal forward primer for improved automated DNA sequencing of M13. *BioTechniques* 15:580-582.

This work was supported in part by grant nos. MOP-14086 and MOP-36449 from the Medical Research Council of Canada/Canadian Institutes of Health Research (MRCC/CIHR) and by a grant from the Fondation des Maladies du Cœur du Québec to C.F.D. Address correspondence to Dr. Christian Deschepper, Laboratory Director, Experimental Cardiovascular Biology, Institut de Recherches Cliniques de Montréal (IRCM), 110 Pine Ave West, Montréal, QC, Canada, H2W 1R7. e-mail: deschec@ircm.qc.ca

Received 5 January 2001; accepted 11 April 2001.

**Isabelle Boutin-Ganache,
Marco Raposo, Martine
Raymond, and Christian F.
Deschepper**
*Institut de Recherches Cliniques
de Montréal (IRCM)
Montréal, QC, Canada*

Human Duplex Sex Determination PCR

BioTechniques 31:— (July 2001)

Applications for the development of molecular sex diagnostics are numerous. One of the first such applications was pre-implantation sex diagnosis to provide choice to known female carriers of sex-linked disorders (3,9). This was a logical extension of the earlier cytogenetic methods of prenatal sex diagnosis that enabled decision making for pregnancy termination before specific gene tests became available (14). Molecular determination of sex often accompanies and supports "sex assignments" after cytogenetic studies of patients presenting with ambiguous genitalia because, "although determination for the sex-of-rearing is generally routine, in certain disorders of sexual differentiation considerable deliberation to determine the sex is necessary at birth" (21). Anatomically and cytogenetically remarkable patients are often enlightened as to their sex by these studies (11,15,16). In athletics, gender verification of female athletes is required in Olympic games (20,21). In many forensic situations, establishing the sex of a tissue can rule in or out suspects or victims (18,19). In addition, every laboratory occasionally experiences labeling discrepancies. A sex diagnostic can help to resolve these and to provide a firm result when the choice of correct assignments is between two members of the opposite sex.

In laboratory medicine, accurate knowledge of a patient's sex can become critical for test interpretation. For example, in the course of simultaneous cytogenetic and molecular testing for Fragile X Syndrome carriers and affected patients, cytogenetic characterization usually lags behind the molecular results. This is a reflection of the speed of technology for cytogenetics versus molecular genetics. Because only one allele is seen in males and homozygous females using fluorescent semi-automated sizing (5,8), sex assignments made by FMR-1 PCR must be verified by patient record or cytogenetics before molecular test results can be issued. Without verification, many full muta-

Benchmarks

tion carrier females risk misdiagnosis. If the sex denoted on patient records is inaccurate, incorrect reports will be issued. A simple, rapid PCR for sex determination would expedite the issuance of patient results in such a circumstance.

We developed two pairs of primers, one pair in the coding region of the sex-determining gene, SRY (22), on the human Y-chromosome, and one pair in the coding region of an ATP-Binding Cassette (ABC) gene, ABCD1, that is the etiologic cause of X-linked adrenoleukodystrophy (X-ALD) (13). These primers, each used at a final concentration of 0.2 μ M, are as follows: SRY344F: 5'-GCATTCATCGTGTGGTCTCG-3' and SRY574R: 5'-GCAAT-TCTTCGGCAGCATCT-3' and ALD1F: 5'-CACCAGCCCCAGTCCCTAC-3' and ALD2R: 5'-GGTAGCAGGGAG-GGCGAT-3'. In preliminary experiments, we found that some of the genomic DNAs provided to us did not amplify well. Under conditions of poor

initial amplification, DNA was subjected to a Gentra Generation® Capture Column (Gentra Systems, Minneapolis, MN, USA) according to the manufacturer's instructions, except that DNA in water or buffer was applied in lieu of freshly mixed whole blood. Subsequent to this rapid column purification, each DNA amplified well. PCRs were multiplexed through the use of one unit of AmpliTaq Gold® thermostable polymerase (Applied Biosystems, Foster City, CA, USA) in 50- μ L reaction mixtures, using 50 ng genomic DNA per reaction. Reactions were optimized through use of the PCR Optimizer™ Kit (Invitrogen, Carlsbad, CA, USA), which was used according to the manufacturer's instructions, except that dimethyl sulfoxide (DMSO) was diluted 10-fold before beginning second-level optimization. As a result of kit optimization, we settled on the following cycling buffer: Invitrogen Buffer A [15 mM (NH₄)₂SO₄, 1.5 mM MgCl₂, 60 mM Tris-HCl, pH 8.5] with 1% (v/v) DMSO

and 1 mM each dNTP. Samples were denatured by heating to 95°C for 10 min, followed by 10 cycles consisting of 30 s at 95°C, 45 s at 65°C reduced by one-half degree each cycle to 60°C, and a 1.5-min extension at 72°C. This was followed by 30 cycles of 95°C for 30 s, 58°C for 45 s, and 1.5 min at 72°C, with a final 7-min hold at 72°C. Cycling was carried out on a model 9700 thermocycler (Applied Biosystems) set to emulate model 9600 thermal profiles.

Following amplification, 12- μ L aliquots of PCR products were loaded in a 2% (w/v) agarose gel (Life Technologies, Rockville, MD, USA) containing Tris-Acetate EDTA buffer and separated by electrophoresis for 1 h at 100 W. The gel was then stained with ethidium bromide and electronically imaged using a UV transilluminator mounted with a digital camera.

PCR using these primers resulted in amplification of a 475-bp fragment entirely within the exon 1 coding region of ABCD1 (12) on the X chromosome and a 231-bp fragment entirely within the coding region of the SRY gene. Twenty-four known female and 24 known male DNAs were subjected to the multiplex; each produced the expected gel bands. Two representative female and two representative male PCR products are shown in Figure 1A, lanes 2–5. We then subjected two unknown samples to analysis using this technique. These two samples were suspected of being mislabeled in our laboratory, as is explained below and illustrated in Figure 1B.

Although a number of sex determination PCRs have been described in the literature, many of these are flawed. Among the problems encountered in previous sex determination PCR systems are the following: (i) the use of primers within the TTCCA repeat regions of DYZ1, (ii) the use of primers in the zinc finger (Zfy) locus, (iii) the use of obligate nested primers, (iv) the lack of multiplexing, (v) the presence of intronic primers, and (vi) the use of the amelogenin locus.

Use of the portions of the 3.4-kb Y-specific TTCCA repeat regions of DYZ1 has become a notorious source of sex misdiagnosis (3,9). Because this repetitive sequence is similar to chromosome 1-derived satellite III, it can be

Figure 1. Sex determination gel and sample mix-up pedigree. (A) Lane 1, 100-bp ladder; lane 2, known female DNA 1; lane 3, known female DNA 2; lane 4, known male DNA 1; lane 5, known male DNA 2; lane 6, person labeled “196” on the tube; lane 7, person labeled “197” on the tube. (B) Pedigree of a nuclear family affected with NBCC and subject to a sample mix-up. Numeric labels correspond with the numbers on the tubes; note that these are NOT the correct sex assignments. For details, see text.

Benchmarks

deleted in males or even carried by females if translocated to autosomes or chromosome X (3,9). A confounding factor accompanying the use of the kuppel Zfy locus (approximately 200 kb proximal to the pseudoautosomal boundary) and of the amelogenin locus (23) for sex determination studies is that these may not be present in all presenting cryptic males (11,16). In particular, Zfy has not been detected in patients who are XX true hermaphrodites nor is it present in many XX male pseudohermaphrodites (1,15,25). In addition, the amelogenin locus has recently been implicated in forensic misdiagnosis through the presence of a polymorphism underlying a common PCR primer (19). Although SRY was discovered to be the male-determining locus in humans approximately 10 years ago and has been the subject of many sex determination PCR methods (2,4,7,10,17,24), none of these are as robust as the assay described here. Most of the SRY PCRs published to date have required the use of "nested" primer pairs (3,7,9,10,17). The use of such nesting has become notorious for its susceptibility to contamination (26). The need for "open-tube" nesting for even very small samples may be obviated through the use of degradable dUMP primers (6). Unfortunately, a few methods for sex determination PCR that use SRY have not been multiplexed, so there is no internal control to make certain that the PCR was proceeding (2,24). Such reactions are inadequately controlled for unambiguous sex diagnosis.

We used the duplexed sex determination diagnostic described here to resolve a sample interchange that occurred during the study of one family affected with Nevroid Basal Cell Carcinoma (NBCC; no. BCC3382). In this nuclear family with characteristics of NBCC, the mother's sample was apparently mislabeled "196". Through the use of the multiplexed X and Y chromosome-specific PCR, we concluded that the tube labeled "196" was the father's DNA (Figure 1, A and B). Mother and son are affected with NBCC in this family. It is now clear that the maternal G1513C mutation in one copy of the PTCH gene segregates with this disease.

We believe the sex determination test described here has high utility in molec-

ular diagnostics laboratories. We have demonstrated only one of many possible applications in this short communication. However, we believe that, after further validation in a variety of research and clinical settings, a non-nested SRY duplex will be found preferable to other molecular sex determination diagnostics currently available.

REFERENCES

1. Abbas, N., G. Novelli, N.C. Stella, O. Triolo, F. Corrado, M. Fellous, M. Chery, S. Gilgenkrantz, and B. Dallapiccola. 1990. A 45,X male with molecular evidence of a translocation of Y euchromatin onto chromosome 1. *Hum Genet.* 86:94-98.
2. Choi, S.K., J.W. Kim, S.Y. Park, Y.M. Kim, J.M. Kim, H.M. Ryu, J.S. Yang, and S.R. Yoon. 1999. Retroactive DNA analysis for sex determination and dystrophin gene by polymerase chain reaction with archived cytogenetic slides. *Exp. Mol. Med.* 31:36-41.
3. Cui, K.H., G.M. Warnes, R. Jeffrey, and C.D. Matthews. 1994. Sex determination of preimplantation embryos by human testis-determining-gene amplification. *Lancet* 343:79-82.
4. Finch, J.L., R.M. Hope, and A. van Daal. 1996. Human sex determination using multiplex polymerase chain reaction (PCR). *Sci. Justice* 36:93-95.
5. Fu, Y., D.P. Kuhl, A. Pizutti, M. Pieretti, J.S. Sutcliffe, S. Richards, A.J. Verkerk, J.J. Holden, R.G. Fenwick, Jr., S.T. Warren et al. 1991. Variation of the CGG repeat at the Fragile X site results in genetic instability: resolution of the Sherman paradox. *Cell* 67:1047-1058.
6. Grace, M.B., G.S. Buzard, M.R. Hughes, and R.E. Gore-Langton. 1998. Degradable dUMP outer primers in merged tandem (M/T)-nested PCR: low- and single-copy DNA target amplification. *Anal. Biochem.* 263:85-92.
7. Hahn, S., X.Y. Zhong, M.R. Burk, C. Troeger, and W. Holzgreve. 2000. Multiplex and real-time quantitative PCR on fetal DNA in maternal plasma. A comparison with fetal cells isolated from maternal blood. *Ann. NY Acad. Sci.* 906:148-152.
8. Hamdan, H., J.A. Tynan, R.A. Fenwick, and J.A. Leon. 1997. Automated detection of trinucleotide repeats in Fragile X syndrome. *Mol. Diagnosis* 2:259-269.
9. Handyside, A.H., E.H. Kontogianni, K. Hardy, and R.M. Winston. 1990. Pregnancies from biopsied human preimplantation embryos sexed by Y-specific DNA amplification. *Nature* 344:768-770.
10. Hussey, N.D., H. Donggui, D.A. Froiland, D.J. Hussey, E.A. Haan, C.D. Matthews, and J.E. Craig. 1999. Analysis of five Duchenne muscular dystrophy exons and gender determination using conventional duplex polymerase chain reaction on single cells. *Mol. Hum. Reprod.* 11:1089-1094.
11. Kusz, K., M. Kotecki, A. Wojda, M. Szarras-Czapnik, A. Latos-Bielenska, A. Warenik-Szymankiewicz, A. Ruszczynska-Wolska, and J. Jaruzelska. 1999. Incomplete masculinisation of XX subjects carrying the SRY gene on an inactive X chromosome. *J. Med. Genet.* 36:452-456.
12. Lachtermacher, M.B., H.N. Seunaez, A.B. Moser, H.W. Moser, and K.D. Smith. 2000. Determination of 30 X-linked adrenoleukodystrophy mutations, including 15 not previously described. *Hum. Mutat.* 15:348-353.
13. Mosser, J., A.M. Douar, C.O. Sarde, P. Kioschis, R. Feil, H. Moser, A.M. Poustka, J.L. Mandel, and P. Aubourg. 1993. Putative X-linked adrenoleukodystrophy gene shares unexpected homology with ABC transporters. *Nature* 361:726-730.
14. Mulley, J.C., A.M. Turner, A.K. Gedeon, V.A. Berdoukas, T.H. Huang, D.H. Ledbetter, H. Grierson, and D.T. Purtilo. 1992. X-linked lymphoproliferative disease: prenatal detection of an unaffected histocompatible male. *Clin Genet.* 42:76-79.
15. Nakagome, Y., S. Seki, K. Fukutani, S. Nagafuchi, Y. Nakahori, and T. Tamura. 1991. PCR detection of distal Yp sequences in an XX true hermaphrodite. *Am. J. Med. Genet.* 41:112-114.
16. Osipova, G.R., M.E. Karmanov, S.I. Kozlova, and O.V. Evgrafov. 1998. PCR detection of Y-specific sequences in patients with Ullrich-Turner syndrome: clinical implications and limitations. *Am. J. Med. Genet.* 76:283-287.
17. Palmirotta, R., F. Verginelli, A. Cama, R. Mariani-Costantini, L. Frati, and P. Battista. 1998. Origin and gender determination of dried blood on a statue of the Virgin Mary. *J. Forensic Sci.* 43:431-434.
18. Reynolds, R. and J. Varlaro. 1996. Gender determination of forensic samples using PCR amplification of ZFX/ZFY gene sequences. *J. Forensic Sci.* 41:279-286.
19. Roffey, P.E., C.I. Eckhoff, and J.L. Kuhl. 2000. A rare mutation in the amelogenin gene and its potential investigative ramifications. *J. Forensic Sci.* 45:1016-1019.
20. Serrat, A. and A. Garcia de Herreros. 1996. Gender verification in sports by PCR amplification of SRY and DYZ1 Y chromosome specific sequences: presence of DYZ1 repeat in female athletes. *Br. J. Sports Med.* 30:310-312.
21. Simpson, J.L., A. Ljungqvist, A. de la Chapelle, M.A. Ferguson-Smith, M. Genel, A.S. Carlson, A.A. Ehrhardt, and E. Ferris. 1993. Gender verification in competitive sports. *Sports Med.* 16:305-315.
22. Sinclair, A.H., P. Berta, M.S. Palmer, J.R. Hawkins, B.L. Griffiths, M.J. Smith, J.W. Foster, A.M. Frischauf, R. Lovell-Badge, and P.N. Goodfellow. 1990. A gene from the human sex-determining region encodes a protein with homology to a conserved DNA-binding motif. *Nature* 346:240-244.
23. Sullivan, K.M., A. Mannucci, C.P. Kimpston, and P. Gill. 1993. A rapid and quantitative DNA sex test: fluorescence-based PCR analysis of X-Y homologous gene amelogenin. *BioTechniques* 15:636-641.
24. Utsumi, K. and A. Iritani. 1993. Embryo sexing by male specific antibody and by PCR

using male specific (SRY) primer. *Mol. Reprod. Dev.* 36:1238-1241.

25. **Waibel, F., G. Scherer, M. Fraccaro, T.W. Hustinx, J. Weissenbach, J. Wieland, A. Mayerova, E. Back, and U. Wolf.** 1987. Absence of Y-specific DNA sequences in human 46,XX true hermaphrodites and in 45,X mixed gonadal dysgenesis. *Hum. Genet.* 76:332-336.
26. **Young, K.K., J.J. Archer, O. Yokosuka, M. Omata, and R.M. Resnick.** 1995. Detection of hepatitis C virus RNA by a combined reverse transcription PCR assay: comparison with nested amplification and antibody testing. *J. Clin. Microbiol.* 33:654-657.

The authors wish to thank Dr. Alisa Goldstein for providing NBCC family samples and Dr. Stephen O'Brien for supplying sex-identifiable controls. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. Government. This project has been funded in whole or in part with Federal Funds from the National Cancer Institute, National Institutes of Health. Address correspondence to Dr. Bert Gold, Laboratory of Genomic Diversity, The National Cancer Institute at Frederick, Building 560, Room 11-85, Frederick, MD 21702, USA. e-mail: goldb@ncifcrf.gov

Received 29 January 2001; accepted 26 April 2001.

Bert Gold, Julie Bergeron, Marcia Lachtermacher-Triunfol, and Michael Dean
*The National Cancer Institute at Frederick
Frederick, MD, USA*

Optimized Growth Conditions for Direct Amplification of cDNA Clone Inserts from Culture

BioTechniques 31:___-___ (July 2001)

Microarray technology plays a significant role in today's quest for gene discovery and expression. DNA mi-

Table 1. Reaction Master Mixture Prepared for Each 96-Well Microplate

Milli-Q® water	3.2 mL	
10× PCR buffer	500 µL	
MgCl ₂ (25 mM)	500 µL	
M13 forward primer (10 µM)	100 µL	
M13 reverse primer (10 µM)	100 µL	
dNTP mixture (2 mM/dNTP) ^a	500 µL	(Life Technologies, Rockville, MD, USA)
Platinum Taq (5 U/µL)	20 µL	(Life Technologies)
Total:	4.92 mL	
In the mixture, the M13 amplification primers are M13 forward primer, 5'-GTTTTCCTCCAGT-CACGACGTTG-3' and M13 reverse primer, 5'-TGAGCGGATAACAATTCACACAG-3'.		
^a 10 µL each dNTP as 100 mM stock in 460 µL Milli-Q water.		

croarrays are typically constructed by arraying a large number of PCR products on derivatized glass slides. Gene expression is then assayed in a co-hybridization assay that allows two distinct tissue types, developmental stages, or cellular phenotypes to be compared (2). High-quality, high-yield PCR products are a crucial element in obtaining accurate, consistent, and reliable results from microarray assays. PCR products can be prepared using either purified plasmid templates or directly from bacterial culture; many groups preparing microarrays have opted for the second approach because it avoids the cost and labor associated with plasmid template preparation (1). To date, we have amplified more than 30 000 cDNA clones from bacterial culture for printing microarrays. We observed that the manner in which the cultures are handled before PCR amplification has a significant impact on both the quality and quantity of the products. For re-arrayed libraries used for microarray construction, it is often difficult to obtain uniform growth from frozen glycerol stocks and consequently difficult to achieve uniform PCR amplification. To address this problem, we investigated a variety of growth conditions and optimized these for direct PCR amplification from bacterial culture.

For each growth method, clone inserts were amplified in 50 µL reactions in 96-well microplates (Applied Biosystems, Foster City, CA, USA) using an identical protocol. Table 1 shows the master mixture that was prepared for each 96-well microplate.

For each clone, 48 µL master mixture is added to 2 µL supernatant prepared from bacterial culture. Reactions are amplified in a thermocycler (PTC-225 Tetrad™; MJ Research, Waltham, MA, USA) using the following cycling protocol: initial denaturation at 95°C for 2 min, 30 cycles each of denaturation at 95°C for 30 s, annealing at 52°C for 30 s, extension at 72°C for 2 min, and hold at 4°C.

Bacterial supernatant was prepared from culture, starting with -80°C glycerol stocks, using each of the methods described below. PCR amplification products were checked by agarose gel electrophoresis in a 1.5% gel, stained with ethidium bromide, and visualized by UV transillumination.

In Method 1 (Figure 1A), 5 µL cells from the -80°C glycerol stock plate were added directly to 95 µL double-distilled water in a 96-well, U-bottom microplate (Falcon®). The microplate was then placed in a 95°C laboratory oven for 10 min to lyse the cells and release the plasmids. Following lysis, the microplate was centrifuged at 1200× g for 3 min. The supernatant (2 µL) was then added to the 96-well microplate containing the PCR master mixture and amplified as previously described.

In Method 2A (Figure 1B), 5 µL cells from the -80°C stock plate were inoculated into a 96-well, deep-well block (Qiagen, Valencia, CA, USA) with each well containing 1.25 mL LB/ampicillin (50 µg/µL). The cells were grown for 16 h in a 37°C shaking incubator at 200 rpm. Following overnight growth, 5 µL culture from each well