Short Technical Reports

used for the screening of deletion mutants or point-mutated proteins for interactions. Finally, our system will enable the large-scale production of modified proteins by co-expression of a substrate protein with an enzyme that will modify it (e.g., a protein kinase, a protease, a methylase, etc.). This system could also be used as a rapid and selective screening method for mapping such modifications without the interference of other enzymes that tend to complicate experiments in eukaryotic cells. For example, proteins normally phosphorylated by multiple kinases could be studied using one kinase at a time.

REFERENCES

1. **Bray, D.** 1998. Signaling complexes: biophysical constraints on intracellular communication. *Ann. Rev. Biophys. Biomol. Struct.* 27:59-75.
2. **Canagarajah, B.J., A. Khokhlatchev, M.H. Cobb, and E.J. Goldsmith.** 1997. Activation mechanism of the MAP kinase ERK2 by dual phosphorylation. *Cell* 90:859-869.
3. **Chen, F.E., S. Kempik, D.-B. Huang, C. Phelps, and G. Ghosh.** 1999. Construction, expression, purification and functional analysis of recombinant NF κ B p50/p65 heterodimer. *Protein Eng.* 12:423-428.
4. **Chen, X.S., T. Stehle, and S.C. Harrison.** 1998. Interaction of polyomavirus internal protein VP2 with the major capsid protein VP1 and implications for participation of VP2 in viral entry. *EMBO J.* 17:3233-3240.
5. **Jonckheere, H., K. De Vreese, Z. Debyser, J. Vandekerckhove, J. Balzarini, J. Desmyter, E. De Clercq, and J. Anne.** 1996. A two-plasmid co-expression system in *Escherichia coli* for the production of virion-like reverse transcriptase of the human immunodeficiency virus type 1. *J. Virol. Methods* 61:113-125.
6. **Robbins, D.J., E. Zhen, H. Owak, C.A. Vanderbilt, D. Ebert, T.D. Geppert, and M.H. Cobb.** 1993. Regulation and properties of extracellular signal-regulated protein kinases 1 and 2 in vitro. *J. Biol. Chem.* 268:5097-5106.
7. **Sambrook, J., E.F. Fritsch, and T. Maniatis.** 1989. *Molecular Cloning. A Laboratory Manual*, 2nd ed. CSH Laboratory Press, Cold Spring Harbor, NY.
8. **Saxena, M. and T. Mustelin.** 2000. Extracellular signals and scores of phosphatases: all roads lead to MAP kinase. *Sem. Immunol.* 12:387-396.
9. **Saxena, M., S. Williams, J. Brockdorff, J. Gilman, and T. Mustelin.** 1999. Inhibition of T cell signaling by MAP kinase-targeted hematopoietic tyrosine phosphatase (HePTP). *J. Biol. Chem.* 274:11693-11700.
10. **Saxena, M., S. Williams, K. Taskén and T. Mustelin.** 1999. Crosstalk between cAMP-dependent kinase and MAP kinase through hematopoietic protein tyrosine phosphatase (HePTP). *Nat. Cell Biol.* 1:305-311.
11. **Stebbins, C.E., W.G. Kaelin, Jr., and N.P. Pavletich.** 1999. Structure of the VHL-elonginC-elonginB complex: implications for VHL tumor suppressor function. *Science* 284:455-461.

We are grateful to Dr. E. Goldsmith for the Erk2 plasmid and Drs. K. Ely and A. Lombardo for valuable discussions and advice. This work was supported by National Institutes of Health grant nos. AI48032, AI35603, AI41481, and AI40552 to T.M. Address correspondence to Dr. Tomas Mustelin, Laboratory of Signal Transduction, La Jolla Cancer Research Center, The Burnham Institute, 10901 North Torrey Pines Road, La Jolla, CA 92037, USA. e-mail: tmustelin@burnham-inst.org

Received 12 October 2000; accepted 7 February 2001.

Natalya Kholod and Tomas Mustelin

*La Jolla Cancer Research Center
La Jolla, CA, USA*

Elimination of Transcriptional Interference between Tandem Genes in Plant Cells

BioTechniques 31:328-334 (August 2001)

ABSTRACT

Plant cells are commonly transformed with two or more tandemly arranged genes, but how orientation affects their expression is not well understood. We investigated the amount of transcriptional interference occurring between two adjacent genes by cloning luciferase and green fluorescent protein (GFP) genes (promoter - coding sequence - terminator) in all possible orientations and expressing the genes in tobacco protoplasts. When two genes are oriented head-to-tail ($\rightarrow \rightarrow$), the expression of the downstream gene was reduced 80% by the upstream gene. When two genes are oriented tail-to-tail ($\rightarrow \leftarrow$), the expression of the upstream gene was reduced 53% by the expression of the downstream gene. There was no interference when the orientation was head-to-head ($\leftarrow \rightarrow$). Using a chemically inducible gene expression system, we showed that the downstream gene expression was reduced 71% by the induction of an upstream gene. Inserting a mammalian transcription blocker sequence eliminated the interference between the genes in tail-to-tail orientation. The interference in the head-to-tail orientation was eliminated by inserting a 2322-bp λ phage DNA fragment. The terminators in gene constructs did not prevent the transcriptional interference, and the interference was eliminated by designing the orientation of genes and by placing a transcription blocker or a λ phage sequence between genes.

INTRODUCTION

Transcriptional interference refers to the phenomenon of reduced expression of a gene caused by the transcription from an upstream or downstream promoter. It can occur when different

Short Technical Reports

genes (i.e., promoter sequence + protein coding sequence + terminator/polyadenylation sequence) are arranged in tandem or even when the genes are separated by several kilobases (5). Even though the transcriptional interference is well documented in mammalian, yeast, and prokaryotic systems (5,6,13), the mechanism is poorly understood. Possible mechanisms for the reduction of transgene expression by an adjacent promoter include antisense RNA production, which can co-suppress the endogenous copy of the transgene (7,10), RNA polymerase "collision" (11), localized depletion of diffusible transcription factors, and DNA topological constraints (16).

Studies in transgenic plants showed that the expression of an upstream gene was reduced by placing a promoter at the 3' of the gene and in opposing orientation (8,11). A poly(A) signal sequence placed on either side of the transgene blocked the downstream transcriptional read-through and restored the upstream gene expression (8). Configuration of transfer DNA (T-DNA) and whether genes are placed at the left or right border of T-DNA of the binary plant transformation vectors (position effect) also have effects on transgene expression in plants (2,3,12). Tetracycline-dependent activation of an upstream promoter was shown to reduce the downstream gene expression in tomato (16).

Plant biologists routinely express transgenes in plant cells. Very often, more than one gene is cloned next to each other on plasmids before they are introduced into plant cells. The orientation of genes is usually determined by the availability of convenient restriction enzyme sites on the plasmid. How the orientation of a gene affects the transcription (expression) of an adjacent gene is not clearly documented. In this study, we examined the influence of a gene on the expression of an adjacent gene by cloning two genes in all possible orientations. We show that the transcriptional interference between two adjacent genes is dependent on how the genes are oriented and that the interference can be eliminated by inserting a transcription blocker (TB) sequence or a sequence from λ phage.

MATERIALS AND METHODS

Plasmid Constructs

Genes for luciferase (LUC) and green fluorescent protein (GFP) suitable for expression in plant cells were constructed in pBluescript® II SK⁻ using standard procedures (15). As illustrated in Figure 1A, the LUC gene construct contained the cassava vein mosaic virus (CsVMV) promoter (17), LUC coding sequence, and nopaline synthase (NOS) terminator sequence. The GFP gene construct contained a duplicated cauliflower mosaic virus 35S promoter (E35S), tobacco etch virus (TEV) enhancer, GFP coding sequence, and 35S terminator sequence (14). As an internal control for protoplast assays, a *Renilla* luciferase (RLUC) gene was constructed with a 35S promoter and NOS terminator. To obtain plasmids containing both genes in all possible orientations, the GFP gene was cloned before or after the LUC gene in four possible orientations (Figure 1B). In addition, a 154-bp mammalian TB sequence (5), a 702-bp *Bst*EII fragment, a 1519-bp *Stu*I fragment, or a 2322-bp *Hind*III fragment from λ phage DNA (GenBank® accession no. J02459) was cloned between the LUC and GFP genes (Figure 1C). An ecodyson receptor (EcR)-based, chemically inducible gene expression plasmid was constructed by cloning the chimeric GVE gene (CsVMV promoter - GAL4 DNA binding domain:VP16 activation domain:EcR ligand binding domain - NOS terminator), inducible LUC gene (5xGAL4 response element - minimal 35S promoter - LUC coding sequence - 35S terminator), and constitutive RLUC gene (35S promoter - RLUC coding sequence - 35S terminator) in the orientation shown in Figure 1D. All plasmid DNAs were prepared using QIAGEN® Plasmid Maxi kit (Qiagen, Valencia, CA, USA).

Protoplast Transfections

Protoplasts were prepared from actively growing tobacco BY-2 suspension cells and cultured essentially as described in Kikkawa et al. (9). One milliliter of protoplasts (5×10^5 to 1×10^6 cells) were mixed with plasmid DNA and electroporated at 0.56 V for

80 μ s using a electroporation system with petripulser electrode (BTX, San Diego, CA, USA). Equimolar plasmid DNA was added to transfections, and the amount of DNA added varied from 23 to 30 μ g depending on the plasmid size. To standardize the luciferase enzyme assays, 2 μ g RLUC plasmid DNA were used in each transfection as an internal control. Following the electroporation, protoplasts were diluted with 1 mL 2x protoplast culture medium, aliquoted as two 1-mL cultures, and incubated at 27°C for 16 h. Each plasmid construct was tested in 5–7 independent transfection experiments.

Luciferase Enzyme Assay

Protoplast cultures were lysed by adding 250 μ L 5x passive lysis buffer (Promega, Madison, WI, USA) and shaking at room temperature for 10 min. A 20- μ L lysate from each culture (two assays per transfection) was assayed using the Dual-Luciferase® Reporter Assay Kit (Promega) according to the manufacturer's suggestions. The assay was performed with an MLX luminometer plate reader (Dynex, Chantilly, VA, USA). The luciferase enzyme activities were expressed as ratios of luciferase to RLUC activities in relative light units (RLUs).

RESULTS AND DISCUSSION

To determine the transcriptional interference on the expression of one gene from its adjacent gene, two genes (promoter - coding sequence - terminator) were cloned in four possible orientations (Figure 1B). The promoters we used for the expression of GFP and LUC in plant cells were cauliflower mosaic virus 35S and CsVMV promoters. Both 35S and CsVMV promoters are commonly used to express transgenes in plant cells. Both the duplicated 35S promoter (E35S) and the CsVMV promoter showed similar promoter strength in tobacco BY-2 protoplasts in our previous experiments (data not shown).

No endogenous LUC enzyme activity was detected in tobacco BY-2 protoplasts. LUC enzyme activity was about 1 RLU with a LUC/RLUC ratio of

Short Technical Reports

0.0003 in protoplasts transfected with salmon sperm DNA or GFP gene (GFP →). In contrast, protoplasts transfected with the LUC gene (LUC →) showed LUC activities of about 50 000 RLUs with an average LUC/RLUC ratio of 7.07 (SD 0.9; Figure 2A). When the GFP gene was cloned upstream of the LUC gene in a head-to-tail orientation (GFP → → LUC), the LUC activity was reduced by approximately 80%. However, no inhibitory interference was detected on LUC gene expression when the GFP gene was downstream of the LUC gene in the head-to-tail (LUC → → GFP) orientation (Figure 2A). The LUC activity was reduced by 53% when GFP and LUC genes were in tail-to-tail (LUC → ← GFP) orientation. There was no decrease in LUC gene expression when the orientation was head-to-head (GFP ← ← LUC).

To test if the activation of an upstream promoter can cause transcriptional interference on downstream gene expression, we used an EcR-based, chemically inducible gene expression system (unpublished results). This system is similar to a glucocorticoid receptor based, dexamethasone-inducible gene expression system developed for plants by Aoyama and Chua (1). It contains a constitutively expressed chimeric GVE receptor gene (CsVMV promoter - sequences coding for GAL4 DNA binding domain, VP16 activation domain, and EcR ligand binding domain - NOS terminator) and an inducible promoter (five copies of GAL4 response element + minimal 35S promoter) to drive LUC gene expression (Figure 1D). When ligand methoxyfenozide (4) is applied to protoplasts, it binds to and activates the chimeric receptor. The activated receptor then induces LUC gene expression. This system was used in our experiment to activate the upstream LUC gene, and the influence of LUC activation on the downstream constitutive RLUC gene expression was then measured. After the addition of a ligand to protoplasts, the LUC gene expression increased from 3981 (SD 3602) to 26 967 (SD 2681) RLUs, while the RLUC expression decreased from 26 323 (SD 7487) to 7891 (SD 1644) RLUs (Figure 2B).

The above results show that the LUC gene expression was affected by the lo-

cation and the orientation of adjacent GFP gene, and induction of the upstream promoter reduced the downstream gene expression. Neither the 35S nor the NOS terminator was sufficient in preventing the transcriptional interference between tandem genes. In an earlier study (8), the interference was shown to be blocked by placing a termi-

nator (polyadenylation) sequence on either side of the transgene. In our construct GFP → → LUC, LUC activity was reduced 80%, even though there were terminators before the LUC gene and after LUC coding sequence. It is possible that the strength of the promoter is important in determining the degree of transcriptional interference. Our results

Figure 1. Schematic representation of gene constructs used in this study. (A) LUC, GFP, and RLUC genes for plant expression are shown. Coding sequences for LUC, GFP, and RLUC are flanked on the 5' end by the CsVMV, 35S, or E35S promoter and on the 3' end by the 35S terminator (35St) or NOS terminator (NOS). GFP gene in addition contained TEV enhancer sequence. All genes were cloned in pBluescript II SK. (B) Plasmid constructs containing GFP and LUC genes head-to-tail (→ →), tail-to-tail (← →), and head-to-head (→ ←) orientations. (C) TB sequence or a λ DNA fragment is placed between the GFP and LUC genes. (D) Plasmid used for chemical induction of LUC gene. GVE is a chimeric protein consisting of GAL4 DNA binding, VP16 activation, and EcR ligand binding domains. LUC coding sequence is placed behind the response element (where GVE can bind) consisting of five copies of GAL4 binding site (5xGAL) and minimal 35S promoter (M35S). All the above constructs were inserted into polylinker region of the pBluescript II SK in such a way that the sequence coding for the α-peptide of β-galactosidase is 5' to the constructs.

suggest that the transcriptional interference between the GFP and LUC genes was caused by RNA polymerase read-through and/or by antisense RNA production. The interference may not be due to the local depletion of diffusible

transcription factors or DNA topological constraints because the LUC gene expression was expected to be reduced by the expression of GFP in all orientations if depletion of factors or topological constraints were operating.

Figure 2. Transcriptional interference between tandemly cloned genes. (A) Reduced expression of LUC gene because of the adjacent GFP gene. Protoplasts were transfected with the constructs shown on the left along with the RLUC plasmid as an internal control. LUC and RLUC enzyme activities were measured in RLUs as described in Materials and Methods. LUC expression levels are shown as ratios of LUC/RLUC activities. The error bars represent standard deviations based on 5–7 independent transfections. (B) Activation of an upstream promoter using a chemically inducible system reduced the expression of the constitutive downstream gene. Protoplasts transfected with the plasmid shown in Figure 1D were divided into two parts, and one part received no chemical (uninduced) and the other part received 10 μ M methoxyfenozide (induced). LUC and RLUC activities are shown in RLUs. The error bars represent standard deviations based on three independent transfections. (C) Transcriptional interference is eliminated by cloning TB or λ DNA sequence between the genes. The TB sequence, a 702-, 1519-, or 2322-bp λ DNA fragments were cloned between GFP and LUC gene constructs shown on the left. Plasmid constructs were introduced into protoplasts along with the RLUC plasmid (as an internal control) and LUC and RLUC activities were assayed. LUC activities are shown as ratio of LUC/RLUC activities. The error bars represent standard deviations based on 5–7 independent transfections.

Short Technical Reports

Our results show that transcriptional interference can be avoided by cloning two genes in the head-to-head ($\leftarrow \rightarrow$) orientation. However, it is not possible to have head-to-head orientation when three or more genes are cloned in tandem. Also, the integration of a transgene into plants is random, and adjacent genes on the chromosome can cause interference. It was reported in mammalian cells that poly(A) signal and a downstream transcriptional pause site are required for transcriptional termination by RNA polymerase II. Placing a poly(A) signal, a transcription pause site or both between tandem genes blocked the transcriptional interference (5). On this basis, we examined if the transcriptional interference on a downstream gene expression could be eliminated by placing an additional sequence after the upstream gene terminator. We cloned a mammalian TB sequence between the GFP and LUC genes in head-to-tail [GFP $\rightarrow$ (TB) $\rightarrow$ LUC] and tail-to-tail [LUC $\rightarrow$ (TB) $\leftarrow$ GFP] orientations. The TB sequence is a 154-bp DNA fragment containing a synthetic poly(A) site and a transcription pause site from human α 2 globin gene (5). As shown in Figure 2A, the transcription interference was completely blocked by TB sequence in the construct LUC $\rightarrow$ (TB) $\leftarrow$ GFP and partially blocked in the construct GFP $\rightarrow$ (TB) $\rightarrow$ LUC. As expected, cloning of the TB sequence before the LUC gene in the construct (TB)LUC $\rightarrow$ $\rightarrow$ GFP and between the LUC and GFP genes in the construct LUC $\leftarrow$ (TB) $\rightarrow$ GFP did not increase the LUC activity as there was no interference in these orientations (Figure 2A).

We also examined if any other DNA fragments can be used to eliminate the transcriptional interference in the construct GFP $\rightarrow$ $\rightarrow$ LUC. Three different DNA fragments from λ phage (702 bp *Bst*EII, 1519 bp *Stu*I, or 2322 bp *Hind*III fragment) were tested by cloning them between the GFP and LUC genes. As shown in Figure 2C, the 702-bp and 1519-bp fragments partially negated the interference. The LUC/RLUC ratio for the construct GFP $\rightarrow$ $\rightarrow$ LUC increased from 1.41 (SD 0.3) to 4.42 (SD 1.0) when a 702-bp λ DNA fragment was inserted (GFP $\rightarrow$ (702) $\rightarrow$ LUC) and to 4.88 (SD 0.2) when a

1519-bp fragment was inserted (GFP $\rightarrow$ (1519) $\rightarrow$ LUC). Insertion of a 2322-bp λ DNA fragment between GFP and LUC genes (GFP $\rightarrow$ (2322) $\rightarrow$ LUC) not only eliminated the interference but also increased the LUC expression when compared to the construct LUC $\rightarrow$ (Figure 2C). The 702-bp and 1519-bp λ DNA fragments have 46% and 48% AT content, respectively. The 2322-bp fragment, in contrast, has 63% AT content. AT-rich sequences can act as transcription termination signals, and the elimination of interference by the 2322-bp fragment may be due to its higher AT content and/or longer length. Why the constructs GFP $\rightarrow$ (2322) $\rightarrow$ LUC showed increased LUC activity needs further investigation.

Our experiments show that the transcriptional interference between adjacent genes can be significant and that the interference can be eliminated by cloning two genes in head-to-head orientation and placing TB or λ sequences between the genes in other orientations.

REFERENCES

1. Aoyama, T. and N.-H. Chua. 1997. A glucocorticoid-mediated transcriptional induction system in transgenic plants. *Plant J.* 11:605-612.
2. Bhattacharyya, M.K., B.A. Stermer, and R.A. Dixon. 1994. Reduced variation in transgene expression from a binary vector with selectable markers at the right and left T-DNA borders. *Plant J.* 6:957-968.
3. Breynne, P., G. Gheysen, A. Jacobs, M.V. Montagu, and A. Depicker. 1992. Effects of T-DNA configuration on transgene expression. *Mol. Gen. Genet.* 235:389-396.
4. Dhadialla, T.S., G.R. Carlson, and D.P. Le. 1998. New insecticides with ecdysteroid and juvenile hormone activity. *Annu. Rev. Entomol.* 43:545-569.
5. Eggermont, J. and N.J. Proudfoot. 1993. Poly (A) signals and transcriptional pause sites combine to prevent interference between RNA polymerase II promoters. *EMBO J.* 12:2539-2548.
6. Greger, I.H., A. Aranda, and N.J. Proudfoot. 2000. Balancing transcriptional interference and initiation on the *GAL7* promoter of *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* 97:8415-8420.
7. Grierson, D., R.G. Fray, A.J. Hamilton, C.J.S. Smith, and C.F. Watson. 1991. Does co-suppression of sense genes in transgenic plants involve antisense RNA? *Trends Biotechnol.* 9:122-123.
8. Ingelbrecht, I., P. Breynne, K. Vancompernelle, A. Jacobs, M.V. Montagu, and A. Depicker. 1991. Transcriptional interference in transgenic plants. *Gene* 109:239-242.
9. Kikkawa, H., T. Nagata, C. Matsui, and I. Takebe. 1982. Infection of protoplasts from tobacco suspension culture by tobacco mosaic virus. *J. Gen. Virol.* 63:457-467.
10. Mol, J., R. van Blokland, and J. Looter. 1991. More about co-suppression. *Trends Biotechnol.* 9:182-183.
11. Paszty, C.J.R. and P.F. Lurquin. 1990. Inhibition of transgene expression in plant protoplasts by the presence in *cis* of an opposing 3'-promoter. *Plant Sci.* 72:69-79.
12. Peach, C. and J. Velten. 1991. Transgene expression variability (position effect) of CAT and GUS reporter genes driven by linked divergent T-DNA promoters. *Plant Mol. Biol.* 17:49-60.
13. Proudfoot, N.J. 1986. Transcriptional interference and termination between duplicated α -globin gene constructs suggests a novel mechanism for gene regulation. *Nature* 322:562-565.
14. Reichel, C., J. Mathur, P. Eckes, K. Langenkemper, C. Koncz, J. Schell, B. Reiss, and C. Maas. 1996. Enhanced green fluorescence by the expression of an *Aequorea victoria* green fluorescent protein mutant in mono- and dicotyledonous plant cell. *Proc. Natl. Acad. Sci. USA* 93:5888-5893.
15. Sambrook, J., E.F. Fritsch, and T. Maniatis. 1989. *Molecular Cloning: A Laboratory Manual*, 2nd ed. CSH Laboratory Press, Cold Spring Harbor, NY.
16. Thompson, A.J. and S.C. Myatt. 1997. Tetracycline-dependent activation of an upstream promoter reveals transcriptional interference between tandem genes within T-DNA in tomato. *Plant Mol. Biol.* 34:687-692.
17. Verdager, B., K. Alexandre, R.N. Beachy, and C. Fauquet. 1996. Isolation and expression in transgenic tobacco and rice plants, of the cassava vein mosaic virus (CVMV) promoter. *Plant Mol. Biol.* 31:1129-1139.

We thank Eileen Ryan, Diane Talaber, and Dusti Fisher for making some of the plasmid constructs and Eric Lifson, Michael Gore, and Dean Cress for reading the manuscript. Address correspondence to Dr. Malla Padidam, Biotech Research, Rohm and Haas Company, 727 Norristown Road, Spring House, PA 19477-0904, USA. e-mail: rahpad@rohmhaas.com

Received 26 September 2000; accepted 27 February 2001.

**Malla Padidam and
Yajuan Cao**
*Rohm and Haas Company
Spring House, PA, USA*

For reprints of this or
any other article, contact
Reprints@BioTechniques.com