Benchmarks

- Hildebrandt, and E. Fleck. 1997. Use of manganese in RT-PCR eliminates PCR artifacts resulting from DNase I digestion. *BioTechniques* 22:1128-1132.
3. Bolotin, A., S. Mauger, K. Malarme, S. Ehrlich, and A. Sorokin. 1999. Low-redundancy sequencing of the entire *Lactococcus lactis* IL1403 genome. *Antonie Van Leeuwenhoek* 76:27-76.
4. Buchman, G., D. Schuster, and A. Rashtchian. 1993. Selective RNA amplification: a novel method using dUMP-containing primers and uracil DNA glycosylase. *PCR Methods Appl.* 3:28-31.
5. Imboden, P., T. Burkart, and K. Schopfer. 1993. Simultaneous detection of DNA and RNA by differential polymerase chain reaction (DIFF-PCR). *PCR Methods Appl.* 3:23-27.
6. Ivarsson, K. and B. Weijdegard. 1998. Evaluation of the effects of DNase treatment on signal specificity in RT-PCR and in situ RT-PCR. *BioTechniques* 25:630-636.
7. Koo, K. and L. Jaykus. 2000. Selective amplification of bacterial RNA: use of a DNA primer containing mismatched bases near its 3' terminus to reduce false-positive signals. *Lett. Appl. Microbiol.* 31:187-192.
8. Kuipers, O., M. Beerthuyzen, R. Siezen, and W. De Vos. 1993. Characterization of the nisin gene cluster *nisABTCIPR* of *Lactococcus lactis*. Requirement of expression of the *nisA* and *nisI* genes for development of immunity. *Eur. J. Biochem.* 216:281-291.
9. Sambrook, J., E.F. Fritsch, and T. Maniatis. 1989. *Molecular Cloning: A Laboratory Manual*, 2nd ed. CSH Laboratory Press, Cold Spring Harbor, NY.
10. Shuldiner, A.R., K. Tanner, C.A. Moore, and J. Roth. 1991. RNA template-specific PCR: an improved method that dramatically reduces false positives in RT-PCR. *BioTechniques* 11:760-763.

Address correspondence to Dr. Wilbert Sybesma, NIZO Food Research, P.O. Box 20, 6710 BA, Ede, The Netherlands. e-mail: sybesma@nizo.nl

Received 2 February 2001; accepted 16 May 2001.

Wilbert Sybesma, Jeroen Hugenholtz, Igor Mierau, and Michiel Kleerebezem
Wageningen Centre For Food Sciences
Wageningen, The Netherlands

For reprints of this or
any other article, contact
Reprints@BioTechniques.com

Non-Disruptive PNA-FISH Protocol for Formalin-Fixed and Paraffin-Embedded Tissue Sections

BioTechniques 31:472-476 (September 2001)

Formalin fixation and subsequent paraffin embedding are commonly employed practices for the convenient and long-term storage of clinical specimens such as cancer tissues. For fluorescence in situ hybridization (FISH) analyses of these tissues, it is often necessary to isolate the nuclei from tissues to improve the accessibility of probes for hybridization. However, nucleus extraction inevitably disrupts the architectural structure of tissue and may therefore result in the loss of valuable information on the tissues. We have recently reported on human and mouse chromosome-specific peptide nucleic acid (PNA) probes for FISH. In our continued effort, we have developed a non-disruptive PNA-FISH protocol for use on sections of formalin-fixed, paraffin-embedded cancer tissues. Using human chromosome 9- and 11-specific PNA probes, we demonstrated that target signals in sections of colon cancer tissues are comparable to signals in the extracted nucleus. This PNA-FISH technique provides a robust cytogenetic method to study the pathology of tumor tissues embedded in paraffin.

Numerical or structural abnormalities in chromosomes are found in many kinds of tumor cells (6,7,22). FISH has been extremely valuable in the study of human chromosomal abnormalities (2,11,15,18,23,25,26). Formalin fixation and subsequent paraffin embedding are common procedures to store cancer tissues. However, FISH on formalin-fixed and paraffin-embedded tissues often results in poor target signal intensity and high nonspecific background signals. These problems have been attributed to complex cross-linkages of intra- and extracellular materials of tissues caused by formaldehyde fixation. The cross-linkages give rise to a reduction in the accessibility of nuclei to specific probes. Moreover, the size

of chromosome-specific DNA probes is relatively large (e.g., 300–600 bp), which further reduces the efficiency of hybridization in FISH. To overcome these hurdles, formaldehyde-fixed and paraffin-embedded tissues are often mechanically and enzymatically dissected to isolate the nucleus. Extraction of the nucleus from paraffin-embedded tissue was first described by Hedley et al. (12) and subsequently modified and improved in many ways (1,5,13,17,19, 24). However, several aspects of nucleus extraction make the method less attractive. First, nucleus isolation inevitably destroys the architectural integrity of tissues, resulting in a loss of structural information that could be important. Second, nucleus isolation leads to a significant loss of the starting materials. Third, this procedure destroys cytoplasm, and, accordingly, it is impossible to do immunocytochemistry after FISH. Simultaneous FISH and immunocytochemistry may provide valuable information on a given gene and its protein product. Fourth, the procedure is tedious and time consuming.

Previously, we reported on human and murine chromosome-specific PNA probe (3,4). PNAs are nucleic acid mimics that contain a pseudo-peptide backbone to which the nucleotides are attached via a methylene carbonyl linker (9,10,20). Because PNAs form a very stable duplex with complementary DNA sequences, PNA/DNA duplexes have a higher melting temperature than DNA/DNA duplexes and a greater sensitivity to mismatches (4,10,20). Recently, PNA probes have been applied to detect chromosomal abnormalities in human disease (4,8,14). In our continued effort to study chromosomal abnormalities of cancer specimens, we have developed a PNA-FISH protocol for sections of formalin-fixed, paraffin-embedded tissues without altering the architectural integrity. In this report, we present a non-disruptive PNA-FISH protocol using PNA probe specific for a segment of human chromosome 9 and 11.

Colon cancer tissues were used for this study. Formalin-fixed and paraffin-embedded tissue blocks were cut into 4- μ m sections. The slides were heated at 60°C for 1 h and deparaffinized in xylene twice for 10 min at room temperature. The slides were rehydrated in

a series of ethanol solutions (100%, 95%, and 75% for 5 min each) and then washed in PBS. The tissue was treated with 0.03% proteinase K at 37°C for 10 min. The tissue was subsequently washed with PBS, dehydrated in ethanol solution (70%, 80%, 90%, and 100%), and air-dried.

The PNA probe, labeled at the N-terminus with rhodamine and specific for repetitive sequences in the centromeric regions of chromosome 9 and 11, was obtained from PerSeptive Biosystems (Framingham, MA, USA) (4). The PNA probe sequences specific for chromosomes 9 and 11 are 5'-AAT-CAACCCGAGTGCAA-3' and 5'-AG-GGTTTCAGAGCTGCTC-3', respectively. The PNA probe was dissolved in trifluoroacetic acid (0.1%) and heated for 30 min in a 50°C water bath. After heating, the PNA probe was lyophilized and stored at -30°C at a concentration of 0.1 nmol.

Before hybridization, 0.1 nmol lyophilized PNA probe was dissolved in 200 µL prewarmed hybridization mixture [70% deionized formamide and 1% (w/v) blocking reagent (Roche Molecular Biochemicals, Indianapolis, IN, USA) in 10 mM Tris-HCl, pH 7.2]. The blocking reagent was used to reduce background. The dissolved PNA (20 µL for slide) was incubated at 37°C

in the dark for 15 min before application to the prepared slides. A coverslip was sealed onto the slide with rubber cement. Denaturation was then carried out at 80°C for 3 min on the PCR slide plate (PTC-100™; MJ Devices, Waltham, MA, USA), followed by hybridization in a dark, moist chamber at 37°C overnight. Because the PNA probe is directly conjugated with rhodamine, light should be avoided to prevent fluorescence bleaching.

The post-hybridization washes were carried out at room temperature: three in 70% formamide and 10 mM Tris-HCl, pH 7.2, for 10 min each, followed by three in 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, and 0.05% Tween® 20 for 5 min each. The slides were dehydrated in a graded series of ethanol solutions and then air-dried in the dark. Chromosomes were counterstained with 0.1 g/mL 4,6-diamidino-2-phenylindole-dehydrochloride (DAPI). Digital images were visualized with a Nikon® Eclipse E800™ fluorescence microscope equipped with a Sensys® CCD camera (Photometrics, Tucson, AZ, USA) (4).

Figure 1. PNA-FISH in colon cancer (A and B) and metaphase spread of peripheral lymphocyte (C and D). Rhodamine-labeled PNA probes specific for human chromosomes 9 and 11 were applied onto deparaffinized tissue sections and metaphase spread. rhodamine (red) and DAPI (blue) images are captured and merged using the IP Lab® program (Scanalytics, Fairfax, VA, USA).

Figure 1 shows the results from two PNA-FISH performed on intact tissue sections of formalin-fixed and paraffin-embedded colon cancer samples. The background signal is low with distinct chromosome 9- and 11-specific signals apparent in the interphase nuclei. Some nuclei do not show two distinct signals (Figure 1, A and B). We found that this was because part of the nucleus is lost by the sectioning of the tissue or because some signals are not on the focal plane of microscope. These nuclei may also result from an artifact caused by histological sectioning, sequence variations in homologous chromosomes (21), heteromorphism (16), hybridization efficiency, and many other reasons. Chromosome 9- and 11-specific probes were also applied to metaphase spread from peripheral lymphocytes to validate the specificity of each PNA probe (Figure 1, C and D). The structural integrity of the tissue is maintained, which is essential for further histological examination. For comparison, we performed the same PNA-FISH on nuclei extracted from tissues of formalin-fixed and paraffin-embedded colon cancer and the hematoxylin-eosin staining on the same slide after PNA-FISH to observe tissue architecture preservation (Figure 2, A and B). The PNA signals from the tissue sections were comparable to the signals from the extracted nuclei, and tissue architecture was also preserved well in counterstaining.

The objective of this study was to establish a non-disruptive PNA-FISH methodology for interphase cytogenetic analysis, especially for the study of numerical chromosome aberrations in paraffin-embedded cancer tissue. Unlike the previously described nucleus extraction methods, this technology uses the direct application of PNA probes onto deparaffinized tissue sections. The most attractive feature of this protocol is that investigators are able to perform FISH analysis while keeping the architectural integrity of the sample intact. This is especially important because cancer tissue specimens are usually contaminated with the surrounding normal tissues, which can make the cytogenetic analysis of isolated nuclei troublesome. We believe that this non-disruptive PNA-FISH protocol is useful for the study of nu-

Figure 2. PNA-FISH in extracted nuclei from the colon cancer tissue (A) and hematoxylin-eosin staining on the same slide (B) after PNA-FISH in colon cancer tissue to show the preservation of tissue architecture.

Benchmarks

merical chromosome aberrations directly in intact sections of formalin-fixed and paraffin-embedded cancer tissues.

REFERENCES

1. Arnoldus, E.P., E.J. Dreef, I.A. Noordermeer, M.M. Verheggen, R.F. Thierry, A.C. Peters, C.J. Cornelisse, M. Van der Ploeg et al. 1991. Feasibility of in situ hybridization with chromosome specific DNA probes on paraffin wax embedded tissue. *J. Clin. Pathol.* 44:900-904.
2. Blasco, M.A., H.-W. Lee, M.P. Hande, E. Samper, P.M. Lansdorp, R.A. DePinho, and C.W. Greider. 1997. Telomere shortening and tumor formation by mouse cells lacking telomerase RNA. *Cell* 91:25-34.
3. Chen, C., Y.K. Hong, S.D. Ontiveros, M. Egholm, and W.M. Strauss. 1999. Single base discrimination of CENP-B repeats on mouse and human chromosomes with PNA-FISH. *Mamm. Genome* 10:13-18.
4. Chen, C., B. Wu, T. Wei, M. Egholm, and W.M. Strauss. 2000. Unique chromosome identification and sequence-specific structural analysis with short PNA oligomers. *Mamm. Genome* 11:384-391.
5. Diebold, J., K. Mosinger, G. Peiro, U. Pannekamp, C. Kaltz, G.B. Baretton, W. Meier, and U. Lohrs. 2000. 20q13 and cyclin D₁ in ovarian carcinomas. Analysis by fluorescence in situ hybridization. *J. Pathol.* 190:564-571.
6. Dierlamm, J., I. Wlodarska, L. Michaus, M. Stefanova, K. Hinz, H. Van Den Berghe, A. Hagemeijer, and D.K. Hossfeld. 2000. Genetic abnormalities in marginal zone B-cell lymphoma. *Hematol. Oncol.* 18:1-13.
7. Dos Santos, N.R. and A.G. Van Kessel. 1999. Chromosomal abnormalities: detection and implications for cancer development. *Anticancer Res.* 19:4697-4714.
8. Drobniewski, F.A., P.G. More, and G.S. Harris. 2000. Differentiation of mycobacterium tuberculosis complex and nontuberculous mycobacterial liquid cultures by using peptide nucleic acid-fluorescence in situ hybridization probes. *J. Clin. Microbiol.* 38:444-447.
9. Egholm, M., O. Buchardt, L. Christensen, C. Behrens, S.M. Freier, D.A. Driver, R.H. Berg, S.K. Kim et al. 1993. PNA hybridizes to complementary oligonucleotides obeying the Watson-Crick hydrogen-bonding rules. *Nature* 365:566-568.
10. Egholm, M., O. Buchardt, P.E. Nielson, and R.H. Berg. 1992. Peptide nucleic acids (PNA). Oligonucleotide analogues with an achiral peptide backbone. *J. Am. Chem. Soc.* 114:1895-1897.
11. Hande, M.P., E. Samper, P. Lansdorp, and M.A. Blasco. 1999. Telomere length dynamics and chromosomal instability in cells derived from telomerase null mice. *J. Cell Biol.* 144:589-601.
12. Hedley, D.W., M.L. Friedlander, I.W. Taylor, C.A. Rugg, and E.A. Musgrove. 1983. Method for analysis of cellular DNA content of paraffin-embedded pathological material using flow cytometry. *J. Histochem. Cytochem.* 31:1333-1335.
13. Isaken, C.V., B. Ytterhus, and S. Skarsvag. 2000. Detection of trisomy 18 on formalin-fixed and paraffin-embedded material by fluorescence in situ hybridization. *Pediatr. Dev. Pathol.* 3:249-255.
14. Krishan, L.T., A.C. Elizabeth, C. James, and P.M. Lansdorp. 2001. Multicolor fluorescence in situ hybridization with peptide nucleic acid probes for enumeration of specific chromosomes in human cells. *Genes Chromosomes Cancer* 30:57-63.
15. Lansdorp, P.M., N.P. Verwoerd, F.M. van de Rijke, V. Dragowska, M.-T. Little, R.W. Dirks, A.K. Raap, and H.J. Tanke. 1996. Heterogeneity in telomere length of human chromosomes. *Hum. Mol. Genet.* 5:685-691.
16. Lo, A.W., G.C. Liao, M. Rocchi, and K.H. Choo. 1999. Extreme reduction of chromosome-specific α -satellite array is unusually common in human chromosome 21. *Genome Res.* 9:895-908.
17. Mark, H.F., M.E. Rausch, W.M. Taylor, A. Huth, S. Mark, K. Santoro, K. Zolnierz, K. Ferreira et al. 1998. Sequential flow cytometry and fluorescence in situ hybridization for the study of formalin-fixed, paraffin-embedded breast cancer cells. *Cancer Genet. Cytogenet.* 100:129-133.
18. Martens, U.M., J.M. Zijlmans, S.S.S. Poon, W. Dragowska, J. Yui, E.A. Chavez, R.K. Ward, and P.M. Lansdorp. 1998. Short telomeres on human chromosome 17p. *Nat. Genet.* 18:76-80.
19. Muresu, R., A. Cossu, A.M. Scarpa, E.V. Volpi, P.C. Rocca, A. Pintus, M.G. Tibiletti, F. Tanda et al. 1998. Numerical abnormalities of chromosomes 1 and 10 in endothelial adenocarcinoma: fluorescence in situ hybridization analysis of 23 archival paraffin-embedded samples. *Cancer Genet. Cytogenet.* 107:37-42.
20. Nielson, P.E., M. Egholm, R.H. Berg, and O. Buchardt. 1991. Sequence-selective recognition of DNA by strand displacement with a thymine-substituted polyamide. *Science* 254:1497-1500.
21. O'Keefe, C.L. and A.G. Matera. 2000. α satellite DNA variant-specific oligoprobes differing by a single base can distinguish chromosome 15 homologs. *Genome Res.* 10:1342-1350.
22. Parada, L.A., G. Bardi, M. Hallen, I. Hagerstrand, K.G. Tranberg, F. Mitelman, and B. Johansson. 1997. Cytogenetic abnormalities and clonal evolution in an adult hepatoblastoma. *Am. J. Surg. Pathol.* 21:1381-1386.
23. Taneja, K.L. 1998. Localization of trinucleotide repeat sequences in myotonic dystrophy cells using a single fluorochrome-labeled PNA probe. *BioTechniques* 24:472-476.
24. Tibiletti, M.G., B. Bernasconi, A. Dionigi, and C. Riva. 1999. The applications of FISH in tumor pathology. *Adv. Clin. Pathol.* 3:111-118.
25. Wan, T.S.K., U.M. Martens, S.S.S. Poon, S.-W. Tsao, L.C. Chan, and P.M. Lansdorp. 1999. Absence or low number of telomere repeats at junctions of dicentric chromosomes. *Genes Chromosomes Cancer* 24:83-86.
26. Zijlmans, J.M., U.M. Martens, S.S.S. Poon, A.K. Raap, H.J. Tanke, R.K. Ward, and P.M. Lansdorp. 1997. Telomeres in the mouse have large inter-chromosomal variations in the number of T₂A₃G₃ repeats. *Proc. Natl. Acad. Sci. USA* 94:7423-7428.

The authors acknowledge the support of U.S. Army Prostate Cancer Research Award no. DAMD17-99-1-9032 and National Institutes of Health grant no. NCI R21CA-81732 awarded to W.M.S. Address correspondence to Dr. William M. Strauss, Harvard Institute of Human Genetics, 4 Blackfan Circle, Boston, MA 02115, USA. e-mail: wstrauss@hih.med.harvard.edu

Received 29 November 2000; accepted 5 June 2001.

D.-H. Kim, Y.-K. Hong, M. Egholm¹, and W.M. Strauss
*Harvard Medical School
 Beth Israel Deaconess
 Medical Center
 Boston
¹Applied Biosystems
 Framingham, MA, USA*