Benchmarks

I believe that, with the whole genome sequence information for *Halobacterium* species NRC-1 now available (14), the novel protocol described here will be a useful alternative to the existing protocol and its modifications.

REFERENCES

1. Blaseio, U. and F. Pfeiffer. 1990. Transformation of *Halobacterium halobium* development of vectors and investigation of gas vesicle synthesis. *Proc. Natl. Acad. Sci. USA* 87:6772-6776.
2. Cline, W.S., W.L. Lam, R.L. Charlebois, L.C. Schalkwyk, and W.F. Doolittle. 1989. Transformation methods for halophilic archaeobacteria. *Can. J. Microbiol.* 35:148-152.
3. Cohen, S.N., A.C.Y. Chang, and L. Hsu. 1972. Nonchromosomal antibiotic resistance in bacteria: genetic transformation of *Escherichia coli* by R-factor DNA. *Proc. Natl. Acad. Sci. USA* 69:2110-2114.
4. DasSarma, S., U.L. RajBhandary, and H.G. Khorana. 1983. High-frequency spontaneous mutation in the bacterio-opsin gene in *Halobacterium halobium* mediated by transposable elements. *Proc. Natl. Acad. Sci. USA* 80:2201-2205.
5. Dower, W.J. 1990. Electroporation of bacteria: a general approach to genetic transformation. *Genet. Eng.* 12:275-295.
6. Gropp, F. and M.C. Betlach. 1994. The *bat* gene of *Halobacterium halobium* encodes a *trans*-acting oxygen inducibility factor. *Proc. Natl. Acad. Sci. USA* 91:5475-5479.
7. Gropp, F., R. Gropp, and M.C. Betlach. 1995. Effects of upstream deletions on light- and oxygen-regulated bacterio-opsin gene expression in *Halobacterium halobium*. *Mol. Microbiol.* 16:357-364.
8. Inoue, H., H. Nojima, and H. Okayama. 1990. High efficiency transformation of *Escherichia coli* with plasmids. *Gene* 96:23-28.
9. Jennert, K.C., C. Tardif, D.I. Young, and M. Young. 2000. Gene transfer to *Clostridium cellulolyticum* ATCC 35319. *Microbiology* 12:3071-3080.
10. Jore, J.P., N. van Lwijk, R.G. Luiten, M.J. van der Werf, and P.H. Pouwels. 2001. Efficient transformation system for *Propionibacterium freudenreichii* based on a novel vector. *Appl. Environ. Microbiol.* 67:499-503.
11. Lam, W.L. and W.F. Doolittle. 1992. Mevinolin-resistant mutations identify a promoter and the gene for an eukaryote-like 3-Hydroxy-3-methylglutaryl-coenzyme A reductase in the archaeobacterium *Haloferax volcanii*. *J. Biol. Chem.* 267:5829-5834.
12. Lorenz, M.G. and W. Wackernagel. 1994. Bacterial gene transfer by natural genetic transformation in the environment. *Microbiol. Rev.* 58:563-602.
13. Mandel, M. and A. Higa. 1970. Calcium-dependent bacteriophage DNA infection. *J. Mol. Biol.* 53:159-163.
14. Ng, W.V., S.P. Kennedy, G.G. Mahairas, B. Berquist, M. Pan, H.D. Shukla, S.R. Lasky, N.S. Baliga et al. 2000. Genome sequence of *Halobacterium* species NRC-1. *Proc. Natl. Acad. Sci. USA* 97:12176-12181.
15. Otomo, J. and T. Muramatsu. 1995. Over-expression of a new photo-active halorhodopsin in *Halobacterium salinarum*. *Biochim. Biophys. Acta* 1240:248-256.
16. Pompejus, M., K. Friedrich, M. Teufel, and H.-J. Fritz. 1993. High-yield production of bacteriorhodopsin via expression of a synthetic gene in *Escherichia coli*. *Eur. J. Biochem.* 211:27-35.
17. Suga, M., M. Isobe, and T. Hatakeyama. 2000. Cryopreservation of competent intact yeast cells for efficient electroporation. *Yeast* 16:889-896.
18. Wagner, G., D. Oesterhelt, G. Kripphal, and J.K. Lanyi. 1981. Bioenergetic role of halorhodopsin in *Halobacterium halobium* cells. *FEBS Lett.* 131:341-345.

Address correspondence to Dr. Arne Zibat, EPIDAUROS Biotechnologie AG, Am Neuland 1, D-82347, Bernried, Germany. e-mail: arne.zibat@epidauros.com

Received 16 May 2001; accepted 6 August 2001.

Arne Zibat
Georg-August-Universität
Göttingen
Göttingen, Germany

PCR Primers and Conditions for Non-Marine Ostracods

BioTechniques 31:1012-1020 (November 2001)

Non-marine ostracods, small bivalved crustaceans, are increasingly used as model organisms for evolutionary studies (3). This is the case for three main reasons. First, ostracods have an excellent fossil record (6) that allows fine-tuned calibration of molecular clocks. Second, the group shows a high frequency of asexual reproduction (4). Third, the ostracod family Darwinulidae is one of the two putative ancient asexual groups in the animal kingdom (7). All of this makes non-marine ostracods particularly interesting to study the "queen of evolutionary problems", the benefit of sexual reproduction (2,9).

This avenue of evolutionary research requires DNA sequence data, on which phylogenetic reconstructions and estimates of genetic diversities can be based. Although PCR-based techniques are now widely used, few known primers will allow cross-species amplification in ostracods because such primers were mostly derived from distantly related invertebrates, such as *Drosophila*. Specific primers for ostracods are thus far published for two species only (11, 12). Furthermore, these belong to two lineages that have been separate for several hundred million years.

Owing to the small size of non-marine ostracods, the amplification of multi-copy regions facilitates both PCR amplification and subsequent cycle sequencing. Such regions can be found in nuclear and mitochondrial DNA. Both genomes are further known to evolve at different rates (8), and some evolutionary questions such as the origin of asexual clones (11) require nuclear and mitochondrial sequence data. Table 1 therefore includes primers that either amplify part of the nuclear ITS1 region or of the mitochondrial COI or 16S genes. Universal primers are used successfully (5,14, Table 1b), and the obtained DNA sequence data form the base for the design of specific primers. Such specific primers have been found

Benchmarks

Table 1a. PCR Primers for Multi-Copy Regions

Primer Name	Locus	Species	Sequence (5'→3')	PCR Conditions	Size of PCR Product (bp)	Accession No. of Submitted Sequence
ITSPB FORW	ITS1	<i>Penthesilenula brasiliensis</i>	AACGCGCTCTCTGG GGTTTTCTCC	50°C	293	AJ319737
ITSPB REV			TGGCGCCCTGCAA TTCGCCGTGTT			
ITS1C FORW	ITS1	<i>Cytherissa sp.</i>	GAAGAGTTTCG TTGGTATCCAATCT	50°C	300	AJ319739
ITS1C REV			AGATCCTCCCCTA AGATAATCAAGGA			
LCOPB FORW	COI	<i>Penthesilenula brasiliensis</i>	CTAAGAGTCTTAATT CGATCCGAAT	45°C	330	AJ319738
HCOPB REV			GGTATTCGGTC TATGGGTATTCC	2 mM MgCl ₂		
LCOCYP FOR	COI	<i>Cyprideis sp.</i>	ATCCGATYTGAGY TCWCACACTCTGG	45°C	605	AJ319742
HCOCYP REV			TGCTGGTATAGA ATGGGGTCTCC	2 mM MgCl ₂		
16SDros FORW	16S	<i>Drosophila melanogaster</i>	CGCCTGTTTA ACAAACAT	45°C	375	AJ319744
16SDros REV		GenBank X53506	CCGGTTTGAA CTCAGATCATGT	2 mM MgCl ₂		

Table 1b. Primers for Cycle Sequencing of Multi-Copy Regions

Primer Name	Locus	Species	Sequence (5'→3')	Annealing Temperature for Cycle Sequencing	Accession No. of Submitted Sequence
HCOCYPR REV	COI	<i>Cyprideis sp.</i>	GGTCAAAAAA GGTGGTGTT	50°C	AJ319742
HCODONG REV	COI	<i>Cytherissa donquixotei</i>	AAAGAAGGAC AGTTGTAATT	50°C	AJ319740
HCOELON REV	COI	<i>Cytherissa elongata</i>	GAGAAGAACAG TCGTGATCAG	50°C	AJ319741
16SCYP FORW	16S	<i>Cyprideis sp.</i>	CAAAGGTAGCA TAATAATTTGGCT	50°C	AJ319743

to significantly improve the success rate of PCR and subsequent automatic sequencing. ITS1 primers (Table 1a) have been designed for *Penthesilenula brasiliensis* of the family Darwinulidae and for *Cytherissa sp.* of the Cytheri-

deidae. According to a BLAST (1) search, both primer pairs amplify the 5'-end of ITS1 and part of 5.8S rRNA. The *Cytherissa* primer set works equally well in other species from the same subfamily (Cytherideinae) and might

even be suitable for the family as a whole. The ITS primer pairs can be used for PCR, resulting in short amplification products that are suitable for SSCP analysis. Alternatively, they can be applied to the automatic sequencing

Benchmarks

Table 2a. Primers for Nuclear Genes

Primer	Locus	GenBank Accession No.	Organism	Sequence (5'→3')	PCR Conditions	Size of PCR Product (bp)	Accession No. of Submitted Sequence
Adh FORW	Alcohol Dehydrogenase	AF045113	<i>Drosophila saltans</i>	CCCTATGATGT	45°C	279	AJ319728
Adh REV		AF067280	<i>Drosophila affinis</i>	GACCGTGCCC	2 mM MgCl ₂		(<i>Darwinula stevensoni</i>)
Adh REV		AF059887	<i>Chymomyza</i> (Diptera)	CATTGAAACCCAG			
Adh EX Rev		AF059888	<i>Drosophila funebris</i>	TCACCTGAACCA	Taq DNA polymerase (Promega)		
Cad FOR	Calmodulin	X05948-X05951	<i>Drosophila melanogaster</i>	TACTCTGTAAATGGCCAGTT	42°C	448	AJ319729
Cad REV				TGGCAAAATAA	2 mM MgCl ₂		(<i>Darwinula stevensoni</i>)
Agk FORW	Arginine Kinase	U09809	<i>Limulus polyphemus</i> (Chelicerata)	AACCGTATAT	HotStar™ (Qiagen, Lab-trade, Miami, FL, USA)		
Agk REV		AI253904	<i>Penaeus monodon</i> (Malacostraca)	ATAAAGATGCA	45°C	338	AJ319730
CYC1 FORW	Cytochrome Oxidase I	AF091461-AF091462	<i>Tigriopus californicus</i> (Copepoda)	CATGTTCTCC	2 mM MgCl ₂		(<i>Darwinula stevensoni</i>)
CYC1 REV		M11382	<i>Manduca sexta</i> (Hexapoda)	GCTGGCTTCAAGAACTTGA	Taq DNA polymerase	288	AJ319731
Pgd FORW	Phospho-gluconate Dehydrogenase	M80598	<i>Drosophila melanogaster</i>	GGCAAGAAGAT	45°C		(<i>Darwinula stevensoni</i>)
Pgd REV		S67873	<i>Ceratitis capitata</i> (Diptera)	CTTTGTGCAGAA	2 mM MgCl ₂		
Tub HA FORW	Tubulin	HA41810	<i>Homarus americanus</i>	TACGACGAAGT	Taq DNA polymerase	253	AJ319732
Tub HA REV				GGCTGACTTTA	Progene (Techne)		(<i>Darwinula stevensoni</i>)
Tub HA FORW				TAATTGGTCAA	45°C	291	AJ319733
Tub HA REV				GGATCGGCG	2 mM MgCl ₂		(<i>Darwinula stevensoni</i>)
				TCCTTGAACCT	HotStar		
				TGGCCACCGT			
				TTCACTCTACAG	45°C		
				ACCGGCCAGT	2 mM MgCl ₂		
				GTTGGACTCTG	HotStar		
				CCTCTGTAAAC	Progene		

Benchmarks

Table 2b. Primers for nuclear genes of *Darwinula stvensoni*

Primer Name	Locus	Species	Sequence (5'→3')	PCR Conditions	Size of PCR Product (bp)	Accession No. of Submitted Sequence
TubDS FORW	Tubulin	<i>Darwinula stvensoni</i>	GGAATTCGATT GATTTGGAGCC	45°C, 2 mM MgCl ₂ HotStar Progene	270	AJ319734
CadDS FORW	Calmodulin	<i>Darwinula stvensoni</i>	TTCAGCAAAGCG TCCCTCAAGTTC	42°C	441	AJ319735
CadDS REV			ACATGTTCTCCA GAGAGGGACAC	HotStar		
AdhDS FORW	Alcohol Dehydro- genase	<i>Darwinula stvensoni</i>	TAATCCCTAG AAGATTGGCT	42°C HotStar	501	AJ319736

of amplification products with universal ITS1 and ITS4 primers (14) and then provide a nested set of primers.

In a similar way as for ITS, primers have been developed for the mitochondrial COI region of *P. brasiliensis* and *Cyprideis* sp.; the latter species belongs to the same subfamily as *Cytherissa*.

Nested primers for automatic sequencing of the reverse direction have further been designed for two endemic species from Lake Baikal (*Cytherissa elongata* and *Cytherissa donquixotei*). These primers also generate high-quality sequences from other species of the same genus. Based on *Drosophila* sequences from GenBank® (Table 1b), a PCR primer pair for the mitochondrial 16S region has been designed and successfully applied to non-marine ostracods. Again, a nested primer for automatic sequencing, this time for the forward direction, has substantially improved the quality of sequences from the genera *Cytherissa* and *Cyprideis*.

All three multi-copy regions are especially suitable to study genetic variability within or between species, although the observed variability might vary considerably between different lineages of non-marine ostracods. For example, populations of *Darwinula stvensoni* from different continents differed only about 2.2% in COI (12), whereas a maximum of 20.9% was reported between *Eucypris virens* populations within Europe (11).

Single-copy genes from non-marine ostracods are much harder to amplify, probably because of the small amount of template. Nevertheless, their sequences can provide important data, especially for the calibration of molecular clocks or for estimating substitution rates between homologue alleles. Table 2a gives an overview of primers from six different nuclear genes, which have been designed from GenBank sequences and

tested on non-marine ostracods. Based on the information available from GenBank, these primers are designed for exon regions that might flank introns, as in the case of Alcohol Dehydrogenase and Calmodulin. Primer design was independently from Reference 10 because these published primers have not been suitable for non-marine ostracods. The primers from Table 2a have been based on sequences from distantly related arthropods, which is why they are equally suitable for other crustaceans such as *Daphnia*, *Artemia*, etc. Because these nuclear genes evolve relatively slowly and code for proteins, they could also offer important, suitable new genetic tools, for example, to reconstruct phylogenies on higher taxonomic levels.

Table 2b summarizes PCR primers that are designed for three nuclear genes of the species *D. stvensoni*. Although these primers only amplify part of each gene, they significantly increase the overall success rate of PCR. They might furthermore not only be suitable for *D. stvensoni* but also for other non-marine ostracod species. Amplifications with primer sets from Table 2, a and b, only yield single PCR products; thus, no evidence for pseudogenes or gene duplications has been found.

DNA extractions from single individuals are conducted with a slightly modified protocol of the Chelex® method (13). If not specifically mentioned otherwise, PCRs are carried out in 25-mL volumes with the model 480 DNA thermal cycler (Applied Biosystems, Foster City, CA, USA). Standard

Benchmarks

conditions include 0.5 U *Taq* DNA polymerase (Amersham Pharmacia Biotech, Piscataway, NJ, USA) with the provided buffer, 1.5 mM MgCl₂, 10 mM dNTPs (Promega, Madison, WI, USA), 10 pmol each primer and mineral oil (Sigma, St. Louis, MO, USA). A hot start (denaturation step of 5 min without *Taq* DNA polymerase before the cycles) is used at all times to increase specificity of the PCR. Modifications of this protocol and the appropriate annealing temperatures are given in Tables 1 and 2. PCR products of multi-copy regions are sequenced directly. PCR products of two successful amplifications are pooled and cleaned with the GFX[®] PCR/DNA and Gel Band Purification Kit (Amersham Pharmacia Biotech). The Thermo Sequenase fluorescently labeled primer cycle sequencing kit (Amersham Pharmacia Biotech) is used to carry out the direct automatic sequencing, following the supplier's protocol. A premixed 6% acrylamide matrix is used (ReproGel-High[™] Resolution; Amersham Phar-

maciac Biotech) for the sequencing gel, which is polymerized under UV light (Reprogel[™]; Amersham Pharmacia Biotech) for 10 min. Sequencing is conducted on an ALFexpress[™] (Amersham Pharmacia Biotech) automatic sequencer with the following conditions: 1500 V, 60 mA, 30 W, 55°C, sample intervals 2 s, running time 750 min. Sequences are automatically processed using ALFwin[™] Sequence Analyzer 2.11 (Amersham Pharmacia Biotech) and are also checked by eye.

Amplification products of single or low copy number genes are cloned before sequencing into the pGEM-T[™] Easy system (Promega), following the manufacturer's protocol. Positive clones are screened by PCR for insert size, using M13 Forward and Reverse primers, standard PCR conditions, and the following program: 3 min at 95°C and 30 cycles of 50 s at 95°C, 50 s at 55°C, and 2 min at 72°C in a Genius (Techne, Princeton, NJ, USA) thermal cycler. Minipreps of positive clones are conducted with the QIAprep[™] Spin Miniprep Kit (Qiagen, Valencia, CA, USA) and automatically sequenced in both directions on ABI PRISM[®] 377 or ABI PRISM[®] 3700 DNA analyzers with M13 primers and the BigDye[™] Terminator Kit (Applied Biosystems), following the manufacturer's protocol. Sequences are automatically processed with Chromas 1.45 and are also checked manually.

The identity of all sequences is verified by BLAST search (1) in GenBank. One sequence of each amplification product has been submitted to GenBank accession nos. AJ319728-AJ319744 (see also Tables 1 and 2).

REFERENCES

1. Altschul, S.F., T.L. Madden, A.A. Schaffer, J. Zhang, Z. Zhang, W. Miller, and D.J. Lipman. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* 25:3389-3402.
2. Bell, G. 1982. *The Masterpiece of Evolution, the Evolution and Genetics of Sexuality*. Croom Helm, London.
3. Butlin, R.K. and P. Menozzi. 2000. Open questions in evolutionary ecology: do ostracods have the answers? *Hydrobiologia* 419:1-14.
4. Chaplin, J.A., J.E. Havel, and P.D.N. Hebert. 1994. Sex and ostracods. *Trends Ecol. Evol.* 9:435-439.
5. Folmer, O., M. Black, W. Hoeh, R. Lutz, and R. Vrijenhoek. 1994. DNA primers for amplification of mitochondrial Cytochrome C Oxidase subunit I from diverse metazoan invertebrates. *Mol. Marine Biol. Biotechnol.* 3:294-299.
6. Horne, D.J. and K. Martens. 2000. Ostracoda and the four pillars of evolutionary wisdom. *Hydrobiologia* 419:vii-xi.
7. Judson, P.O. and B.B. Normark. 1996. Ancient asexual scandals. *Trends Ecol. Evol.* 11:41-46.
8. Li, W.-H. 1998. *Molecular Evolution*. Sinauer Associates, Sunderland, Massachusetts.
9. Martens, K. (Ed.) 1998. *Sex and Parthenogenesis. Evolutionary Ecology of Reproductive Modes in Non-Marine Ostracods*. Bakkhuys Publishers, Leiden, The Netherlands.
10. Palumbi, S.R. 1996. Nucleic acids II. The polymerase chain reaction, p. 205-248. In D.M. Hillis, C. Moritz, and B.K. Mable (Eds.), *Molecular Systematics*. Sinauer Associates, Sunderland, Massachusetts.
11. Schön, I., A. Gandolfi, E. Di Masso, V. Rossi, H.I. Griffiths, K. Martens, and R.K. Butlin. 2000. Long-term persistence of asexuality through mixed reproduction in *Eucypris virens*. *Heredity* 84:161-169.
12. Schön, I., R.K. Butlin, H.I. Griffiths, and K. Martens. 1998. Slow molecular evolution in an ancient asexual ostracod. *Proc. R. Soc. Lond. B Biol. Sci.* 265:235-242.
13. Walsh, P.S., D.A. Metzger, and R. Higuchi. 1991. Chelex[®] 100 as medium for simple extraction of DNA for PCR-based typing from forensic material. *BioTechniques* 10:506-513.
14. White, T.J., T. Bruns, S. Lee, and J. Taylor. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics, p.315-322. In M.A. Innis, D.G. Gelfand, J.J. Sninsky, and T.J. White (Eds.), *PCR Protocols: A Guide to Methods and Applications*. Academic Press, London.

I thank my colleagues Drs. K. Martens and T. Backeljau for comments on an earlier version of the manuscript. The European Commission is acknowledged for my Marie-Curie Post-Doc fellowship BIO4-CT 98-5086 in the programme Biotechnology. Address correspondence to Dr. Isabelle Schön, Royal Belgian Institute of Natural Sciences, Freshwater Biology, Vautierstraat 29, 1000 Brussels, Belgium. e-mail: isa.schoen@naturalsciences.be

Received 21 May 2001; accepted 7 August 2001.

Isabelle Schön
*Royal Belgian Institute of
Natural Sciences
Brussels, Belgium*

**For reprints of this or
any other article, contact
Reprints@BioTechniques.com**