Short Technical Reports

- scription-PCR on oligo(dT)-immobilized polypropylene microplates after capturing total mRNA from crude cell lysates. *Clin. Chem.* 44:2256-2263.
5. **Hyndman, D., A. Cooper, S. Pruzinsky, D. Coad, and M. Mitsuhashi.** 1996. Software to determine optimal oligonucleotide sequences based on hybridization simulation data. *BioTechniques* 20:1090-1097.
6. **Kawasaki, E.S. and A.M. Wang.** 1989. Detection of gene expression, p. 89-97. *In* E.A. Erlich (Ed.), *PCR Technology*. Stockton Press, Stockton, New York.
7. **Myers, T.W. and D.H. Gelfand.** 1991. Reverse transcription and DNA amplification by a *Thermus thermophilus* DNA polymerase. *Biochemistry* 30:7661-7666.
8. **Okamoto, T., M. Mitsuhashi, and Y. Kikkawa.** 1994. Fluorometric nuclear run-on assay with oligonucleotide probe immobilized on plastic plates. *Anal. Biochem.* 221:202-204.
9. **Rieber, M. and R.M. Strasberg.** 2000. Apoptosis-inducing levels of UV radiation and proteasome inhibitors produce opposite effects on p21(WAF1) in human melanoma cells. *Int. J. Cancer* 86:462-467.
10. **Roos, D., A.A. Voetman, L.J. Meerhof, and B. Lubin, inventors; Stichting Vrienden Van De Stichting Dr. Karl Landsteiner; assignee.** 1986. Enucleated granulocytes, their preparation and use. U.S. Patent 4 623 620. 1986 Nov 18.
11. **Urdea, M.S., T. Horn, T.J. Fultz, M. Anderson, J.A. Running, S. Hamren, D. Ahle, and C.A. Chang.** 1991. Branched DNA amplification multimers for the sensitive, direct detection of human hepatitis viruses. *Nucleic Acids Symp. Ser.* 24:197-200.
12. **Walker, G.T., M.C. Little, J.G. Nadeau, and D.D. Shank.** 1992. Isothermal in vitro amplification of DNA by a restriction enzyme/DNA polymerase system. *Proc. Natl. Acad. Sci. USA* 89:392-396.
13. **Wang, J.A., S. Fan, R.Q. Yuan, Y.X. Ma, Q. Meng, I.D. Goldberg, and E.M. Rosen.** 1999. Ultraviolet radiation down-regulated expression of the cell-cycle inhibitor p21Cip1/Waf1 in human cancer cell independently of p53. *Int. J. Radiat. Biol.* 75:301-316.

We would like to thank Dr. M. Kobrin, Mr. T. Rothwell, Ms. M. Ogura, Mr. T. Brooks, and Ms. S. Powell (RNAture) for their technical help and product support related to the RiboCap filter plate and the GenePlate. We would also like to thank Mr. S. Nozaki and Mr. B. Powell (Nissei Sangyo America), Ms. Marieta and Dr. C.E. Ribak (Pathology, University of California, Irvine), and Dr. C. Wen-an (School of Engineering, University of California, Irvine) for their help with the electron microscopy analysis. We would like to thank Dr. J.L. Redpath (Radiation Oncology, University of California, Irvine) for his critical review. This study was supported by grants from the Hitachi Chemical Research Center and Hi-

tachi Ltd., Medical Systems Division (Tokyo, Japan). Address correspondence to Dr. Masato Mitsuhashi, Hitachi Chemical Research Center, 1003 Health Sciences Road West, Irvine, CA 92612, USA. e-mail: mmitsuha@uci.edu

Received 10 September 2001; accepted 4 January 2002.

K. Matsuda^{1,2}, S. Tomozawa^{1,2}, S. Fukusho^{1,3}, T. Yoshino^{1,4}, T. Murakami⁵, and M. Mitsuhashi^{1,5}

¹University of California
Irvine, CA, USA

²Tokyo University

³Tokyo Institute of Technology

⁴Tokyo University of Agriculture
and Technology

Tokyo, Japan

⁵Hitachi Chemical Research
Center
Irvine, CA, USA

Cyclic Nucleotide-Dependent Screening of a λ Expression Library for Nucleic Acid Binding Activities

BioTechniques 32:1020-1028 (May 2002)

ABSTRACT

Here we describe a modified ligand screening strategy for the expression cloning of mammalian proteins that require the activation of protein kinase cascades to activate ligand binding activity. The manipulation of prokaryotic signaling pathways by the application of appropriate inhibitors or agonists to the nitrocellulose filter during functional screening of standard bacteriophage cDNA expression libraries can permit detection of activities that would not

otherwise be found in their active state. We have applied this strategy to a λ expression library to clone a novel renal cDNA that exhibits cAMP-dependent RNA binding activity when subsequently tested by ectopic expression in mammalian cells.

INTRODUCTION

Following the initial introduction of a method to isolate cDNA clones by antibody screening of bacteriophage expression libraries by Young and Davis (22), double-stranded dNTPs (20), proteins (6,18), and RNA ligands (15,16) have been used to screen expression libraries for ligand binding activities of expressed recombinant fusion proteins. Functional screening methods offer the advantage that a protein can be cloned without the knowledge of its partial amino acid sequence or the need to prepare an antibody. However, the ligand binding activity of many proteins is reversibly regulated by cell signaling pathways. In these cases, standard bacteriophage library screening protocols may fail to detect those fusion proteins that require appropriate activation by protein kinase/phosphatase(s).

The modified ligand screening strategy described here offers new possibilities for cloning mammalian proteins that require the activation of protein kinase cascades to activate ligand binding activity. There is now a large body of evidence that prokaryotic cells possess a number of "eukaryotic-type" signaling pathways (1). The manipulation of these pathways during functional screening of standard λ cDNA expression libraries can be accomplished by the application of appropriate inhibitors or agonists to the nitrocellulose filter before overlaying it on the plaques, which permits the detection of activities that would not otherwise be found in their active state.

We have applied this strategy to clone a novel cDNA that exhibits cAMP-dependent RNA binding activity when tested by expression in mammalian cells. The untranslated regions of mRNAs contain highly conserved sequence motifs that regulate the rate of mRNA turnover and thus influence the steady-state level of their protein product (5,19). We have recently identified

Short Technical Reports

a 120-nucleotide uridine-rich element (URE) in the 3'-untranslated region of sodium glucose co-transporter (SGLT1) mRNA, which is required for message stabilization in response to cyclic AMP elevation (7,9,13). Protein binding to the URE is stimulated after protein kinase A activation (7). To identify cDNAs encoding cellular proteins that bind the SGLT1 URE, we utilized an RNA ligand-screening strategy (15) modified to incorporate both IPTG and the phosphodiesterase inhibitor 3-isobutyl-1-methylxanthine (IBMX) into the nitrocellulose filter used for overlay and plaque lifts. The use of IBMX has the additional advantage that the elevated cAMP in *E. coli* increases the activity of the lac promoter in the pBluescript® phagemid (Stratagene, La Jolla, CA, USA) by binding to the catabolite activator protein (4), which permits a more robust expression of the recombinant fusion proteins.

MATERIALS AND METHODS

Materials

The [α - 32 P]uridine triphosphate (3000 Ci/mmol) was obtained from ICN Biomedicals (Costa Mesa, CA, USA). NTPs were from Roche Applied Science (Indianapolis, IN, USA). RNA polymerases T3 and T7 were purchased from Promega (Madison, WI, USA). RNase T1 (*Aspergillus oryzae*) was from Calbiochem-Novabiochem (San Diego, CA, USA). The IBMX, yeast tRNA, and other molecular biology-grade chemicals were obtained from Sigma (St. Louis, MO, USA). The anti-c-myc tag IgG₁ monoclonal antibody was from Calbiochem-Novabiochem (Ab1) and recognizes residues 410–419 of the human c-myc sequence.

Library Screening

A custom cDNA library in λ ZAP II™ (Stratagene) was prepared commercially using poly(A)⁺ RNA from differentiated LLC-PK₁ pig kidney epithelial cells. *E. coli* strain XL-1 Blue (Stratagene) was infected with bacteriophage and then plated at 5×10^4 pfu/dish in NZY top agar (Difco, Detroit, MI, USA) on 20×15 -cm plates

containing NZY bottom agar and 10 μ g/mL tetracycline using the standard Stratagene protocol. To enhance the detection of cyclic nucleotide-dependent binding activities, IBMX was added during the IPTG induction of protein expression. After 3.5–4 h, when pin-point plaques first appeared, nitrocellulose membranes (15 cm, BA 85; Schleicher & Schuell BioScience, Keene, NH, USA) that had first been soaked for 30 min in filter-sterilized 10 mM IPTG and 1 mM IBMX and air dried for 15 min on Whatman® 3-mm filter

paper (Fisher Scientific, Pittsburgh, PA, USA) were overlaid on the plates and incubated for 4 h. Filters were removed, washed briefly in 10 mM Tris-HCl, pH 7.6, 50 mM NaCl to remove loosely bound bacterial debris, air dried for 15 min on Whatman 3-mm filter paper, and then incubated on a rotary shaker for 1 h at 37°C in a baked glass tray containing 1 mg/mL yeast tRNA, 10 mM Tris-HCl, pH 7.6, 40 mM NaCl, 10 mM KCl, 5% glycerol, and 1 mM IBMX. The filters were transferred to an RNA binding buffer containing 1

Figure 1. Expression of URBP36 fusion protein in *E. coli*. Bacterial cultures transformed with plasmid pBS-URBP36 or with the empty pBluescript plasmid (vector) were induced for 4 h with 10 mM IPTG for comparison with uninduced controls. Because of its growth inhibitory effects in liquid culture, 1 mM IBMX was added to the indicated samples only during the last two hours of IPTG induction. Total cell proteins were extracted from cell pellets at 4°C by suspension at 1:4 (w/v) in lysis buffer containing 10 mM Tris-HCl, pH 8.0, 10 mM KCl, 50 mM NaCl, 1 mM EDTA, 1 μ M PMSF, 1 mM EDTA, 1 μ g/mL aprotinin, 0.4 mM dithiothreitol and 10% sucrose. Lysozyme was added to 1 mg/mL final concentration and incubated for 10 min on ice. Nonidet™ P-40 was then added to a final concentration of 0.1%, and the suspension was incubated for 10 min on ice. Suspensions were centrifuged for 1 h at 16 000 $\times$ g, and supernatants were removed and stored at -80°C. (A) Aliquots (10 μ g protein) of each extract were resolved by 10% SDS-PAGE and stained using Coomassie blue. (B) RNA binding activities of bacterial extracts (20 μ g protein) were assayed by the UV cross-linking assay using the indicated 32 P-labeled sense RNA probe.

Short Technical Reports

mg/mL yeast tRNA, 10 mM Tris-HCl, pH 7.6, 40 mM NaCl, 10 mM KCl, 1× Denhardt's solution, and 4×10^6 cpm/mL of a ^{32}P -labeled 120-nucleotide sense RNA SGLT1 URE probe that had been prepared by transcription in vitro as previously described (7). Incubation was carried out for 1 h at room temperature on a shaker to permit RNA/protein binding. Filters were washed four times at room temperature with 10 mM Tris-HCl, pH 7.6, 50 mM NaCl, 1 mM EDTA, and 0.5× Denhardt's solution, air dried on Whatman 3-mm filter paper, covered in Saran® Wrap, and exposed to X-Ray film (Eastman Kodak, Rochester, NY, USA) in the presence of intensifying screens for two days at -80°C . Positive plaques were cored, tittered, and subjected to secondary and tertiary screenings. The recombinant pBluescript plasmids were excised from positive bacteriophage plaques using the R408 helper phage (Strata-

gene) and *E. coli* BB4 host strain, as described in the Stratagene instructions, and then colony-purified.

One cDNA clone, URBP36, which exhibited positive plaques only in the presence of IBMX, was chosen for further analysis. The DNA was isolated using a Wizard® kit (Promega) and sequenced on both strands by automated DNA sequencing (Lone Star Labs, Houston, TX, USA).

Plasmid Construction

URBP36 cDNA was subcloned into the mammalian expression vector pMyc-over (8,14), a pCAT3 derivative (Promega) in which the CAT coding sequence has been replaced by a sequence encoding the myc epitope tag. The pBluescript plasmid pBS-URBP36 was digested with *BsmI* and *SpeI*, followed by fill-in using Klenow DNA polymerase. After ligation to double-strand-

ed *SalI* linkers, the URBP36 fragment was digested with *SalI*, purified, and then ligated to the vector pMyc-over, which had been prepared by digestion with *SalI* and treatment with calf intestinal phosphatase. The resulting plasmid, pMyc-over URBP36, was verified by DNA sequencing to be in frame with the N-terminal myc epitope tag.

Assay of RNA/Protein Complex Formation

Radiolabeled RNA probe-binding to protein was determined using the UV cross-linking and gel mobility shift assays described previously (7,8). The indicated sense RNA probes were synthesized by transcription in vitro in the presence of [α - ^{32}P]uridine triphosphate (3000 Ci/mmol). Nucleotide numbers refer to the porcine SGLT1 sequence (GenBank® accession no. M34044) (12). The plasmids 122 nt/*Bam* and 122

nt m2/*Bam*(7) that contains a TT → GG substitution at nucleotides 2622 and 2623 were linearized with *Bbs*I and transcribed with T3 DNA polymerase to yield wild-type and mutant 120 nucleotide RNA transcripts, respectively, that contain SGLT1 nucleotides 2576–2695. Plasmid T3 adenine + uridine-rich element (ARE) *fos* (21) was obtained from Dr. Ann-Bin Shyu, University of Texas Medical School, Houston. A 251-nucleotide *c-fos* ARE transcript was synthesized using T3 DNA polymerase from *Hind*III-linearized plasmid T3 ARE *fos* and contained 207 nucleotides of the human *c-fos* ARE and 44 nucleotides from plasmid pT7/T3α-19.

Gel mobility supershift analysis was carried out as previously described (8). Briefly, nuclear lysates from LLC-PK₁ cells transiently transfected with either pMyc-over URBP36 DNA or vector controls were incubated for 15 min with the ³²P-labeled 120-nucleotide SGLT1

RNA probe, unprotected RNA was digested using ribonuclease T1, and then either an anti-myc tag IgG₁ monoclonal antibody or a nonspecific control monoclonal antibody was diluted 1:10 into the RNA/protein binding mixtures. Incubation was continued for an additional 30 min in the presence of 1 U RNasin®, and complexes were resolved on non-denaturing 6% polyacrylamide gels.

RESULTS AND DISCUSSION

The URBP36 sequence (GenBank accession no. AF203905) contains an open reading frame of 322 amino acids and both a start and stop codon. The sequence encodes a novel putative protein with a predicted molecular mass of 35.9 kDa. The selection of this positive clone did not result as an artifact of nucleic acid hybridization with the RNA probe since there was no significant ho-

mology between the cloned cDNA sequence and the RNA probe. Northern analysis of the poly (A)⁺ RNA isolated from LLC-PK₁ cells using a riboprobe based on the antisense URBP36 sequence detected a single 2.5-kb mRNA species (data not shown).

Results shown in Figure 1A demonstrate that pBS-URBP36 cultures in *E. coli* induced with both IPTG and IBMX expressed high levels of a 37-kDa protein detectable by Coomassie® blue staining. This band was greatly diminished in the absence of either IPTG or IBMX and was not expressed in cultures transformed with the empty pBluescript vector. These results indicate that recombinant expression induced by IPTG was greatly increased when IBMX was also added during the last two hours of expression. The 37-kDa apparent molecular mass of this band is close to the 35.9-kDa size predicted from the cloned URBP36 sequence.

Short Technical Reports

The extracts from pBS-URBP36 cultures that were treated with both IPTG and IBMX formed a ^{32}P -labeled RNA/protein complex with the wild-type 120-nucleotide SGLT1 probe when assayed by UV cross-linking (Figure 1B). This band (indicated by the arrow) was not formed under these conditions in extracts from cells bearing the empty vector. Since this complex consists of protein cross-linked to a protected RNA fragment remaining after ribonuclease T1 digestion, this technique does not permit the estimation of the molecular mass of the RNA binding protein.

The specificity of the binding was tested using a mutant RNA probe 120 nt m2 (Figure 1B). This probe differed from the wild-type 120-nucleotide probe in that it contained a UU $\rightarrow$ GG substitution at positions corresponding to nucleotides 2622 and 2623 in the porcine SGLT1 cDNA sequence. This mutation prevented cAMP-dependent stabilization of a reporter message and greatly reduced protein binding to the 120-nucleotide sequence (7). The results shown in Figure 1B indicate that complex formation was greatly reduced when IPTG-induced, IBMX-treated bacterial cell extracts were assayed using the 120 nt m2 labeled RNA probe.

Furthermore, this complex was not formed with a 251-nucleotide *c-fos* ARE probe (Figure 1B). This adenine, uridine (AU)-rich sequence binds a number of cellular RNA-binding proteins directly implicated in regulating mRNA stability (3,17).

Taken together, these results indicate that the URBP36 RNA binding activity exhibits binding specificity for the 120-nucleotide SGLT1 URE sequence used as ligand in the library screening. Binding is abolished by a substitution mutation within this sequence and is not observed using another AU-rich sequence. Furthermore, URBP36 binding activity is stimulated after the IBMX treatment of the bacterial cultures, paralleling the increase in levels of a 36-kDa protein detectable by Coomassie blue staining.

URBP36 cDNA was subcloned into plasmid pMyc-over (14), and the recombinant protein, bearing an in-frame N-terminal myc epitope tag, was transiently expressed in LLC-PK $_1$ pig kidney

epithelial cells (Figure 2). The expression of the recombinant protein was demonstrated by Western analysis of transfected LLC-PK $_1$ cell extracts using an anti-myc epitope tag antibody (Figure 2A). The antibody recognized a 45-kDa band in mycURBP36-transfected cultures treated with IBMX. This band was greatly diminished in untreated cultures and in cultures transfected with the empty pMyc-over vector. Its apparent molecular weight agrees well with that predicted from the myc-tagged URBP36 cDNA sequence. These data indicate that the recombinant protein is expressed in transfected LLC-PK $_1$ cells.

To detect RNA binding activity of the expressed recombinant protein, we performed gel mobility supershift analysis using nuclear extracts from transfected cells (Figure 2B). In this analysis, extracts were incubated with the ^{32}P -labeled 120-nucleotide SGLT1

RNA probe, unprotected RNA was digested with ribonuclease T1, and the mixture was then incubated with either an anti-myc epitope tag monoclonal antibody or a control, nonspecific monoclonal antibody. Complexes were immediately resolved on a non-denaturing polyacrylamide gel (10). This analysis permits the detection of multi-protein noncovalent complexes with the RNA probe. Results in Figure 2B demonstrate that the anti-myc epitope tag monoclonal antibody produced a pronounced supershifted complex in nuclear extracts from IBMX-treated, mycURBP36-transfected cultures. No supershift was observed in extracts from cells transfected with the empty vector. The supershifted band was undetectable in the absence of IBMX treatment or after the incubation of extracts with a control, nonspecific monoclonal antibody. Because immunodetectable myc-tagged

Figure 2. Expression of myc-tagged URBP36 cDNA in LLC-PK $_1$ cells. LLC-PK $_1$ cells were transiently transfected with either 20 μg pMyc-over URBP36 DNA or the empty pMyc-over vector DNA/10-cm dish using the calcium phosphate procedure and cell culture conditions previously described (7). Where indicated, 1 mM IBMX (final concentration) was added to the cells immediately after the removal of DNA, and cell extracts were prepared after 48 h. (A) Western analysis of total cell extracts (15 μg protein) using a 1:5000 dilution of the anti-myc tag antibody, HRP-conjugated second antibody and detection using the ECL $^{\circ}$ Western Blotting kit (Amersham Biosciences, Piscataway, NJ, USA) as previously described (8). (B) Gel mobility supershift analysis of nuclear lysates (15 μg) using an anti-myc tag antibody and a labeled 120-nucleotide SGLT1 URE riboprobe (10). The arrow indicates the supershifted band. The "+" indicates anti-myc tag antibody was added and the "-" indicates control, nonspecific monoclonal antibody was added. The figure is cropped to show only the supershifted region.

Short Technical Reports

protein was expressed in the absence of IBMX treatment (Figure 2A), these findings suggest that binding activity is activated after IBMX treatment and protein kinase A activation. The binding activity was predominantly nuclear, with diminished activity in cytosolic extracts (data not shown).

Homology comparisons in sequence databases indicated that the URBP36 sequence is novel. It does not exhibit the typical RNA recognition motifs that are associated with most RNA binding proteins (2). It shared 87% protein sequence identity with a 421-amino-acid protein of unknown function (GenBank accession no. AK058054) cloned from human testis cDNA. A database search using BLASTP version 2.2.1 revealed 25%–35% amino acid identity to a number of DNA binding proteins, including a human putative transposase similar to the pogo element (accession no. U49973), a human centromere protein CENP-B (accession no. AAB21673.1), the yeast autonomously replicating sequence binding protein 1 (accession no. U39079) (11), a human jerky gene protein homolog (accession no. AF004715), a putative centromere autoantigen B/ARS-binding protein (accession no. AC006161), and hypothetical fruit fly transposon pogoR11 (accession no. S20478). The porcine URBP36 sequence also exhibited 85% identity to the base pair 529–543 region of dbEST file AI566011 cloned from human kidney cDNA and 83% identity to base pair 402–432 of dbEST file AI758641 cloned from human testis cDNA.

In conclusion, we have utilized a cyclic AMP-dependent RNA ligand expression screening strategy to clone a novel 36-kDa protein URBP36 from LLC-PK₁ pig kidney epithelial cells. The URBP36 exhibits sequence-specific, cyclic AMP-dependent binding to a regulatory element in the 3'-untranslated region of sodium glucose co-transporter SGLT1 mRNA. Further analysis is needed to determine the physiological significance of this interaction.

This strategy for the manipulation of signaling pathways in functional screening of expression libraries can be readily applied to existing protocols for screening with DNA or protein ligands and extended to other cell-permeable agonists.

REFERENCES

1. **Bakal, C.J. and J.E. Davies.** 2000. No longer an exclusive club: eukaryotic signaling domains in bacteria. *Trends Cell Biol.* 10:32-38.
2. **Burd, C.G. and G. Dreyfuss.** 1994. Conserved structures and diversity of functions of RNA-binding proteins. *Science* 265:615-621.
3. **Chen, C.Y. and A.B. Shyu.** 1995. AU-rich elements: characterization and importance in mRNA degradation. *Trends Biochem. Sci.* 20:465-470.
4. **de Crombrughe, B., S. Busby, and H. Buc.** 1984. Cyclic AMP receptor protein: role in transcription activation. *Science* 224:831-838.
5. **Duret, L. and P. Bucher.** 1997. Searching for regulatory elements in human noncoding sequences. *Curr. Opin. Struct. Biol.* 7:399-406.
6. **Helin, K., J.A. Lees, M. Vidal, N. Dyson, E. Harlow, and A. Fattaey.** 1992. A cDNA encoding a pRB-binding protein with properties of the transcription factor E2F. *Cell* 70:337-350.
7. **Lee, W.Y., P. Loflin, C.J. Clancey, H. Peng, and J.E. Lever.** 2000. Cyclic nucleotide regulation of Na⁺/glucose cotransporter (SGLT1) mRNA stability: interaction of a nucleocytoplasmic protein with a regulatory domain in the 3' untranslated region critical for stability. *J. Biol. Chem.* 275:33998-34008.
8. **Loflin, P., C.-Y. Chen, and A.B. Shyu.** 1999. Unraveling a cytoplasmic role for hnRNP D in the in vivo mRNA destabilization directed by the AU-rich element. *Genes Dev.* 13:1884-1897.
9. **Loflin, P. and J.E. Lever.** 2001. A cis-dominant cyclic nucleotide-dependent regulatory domain in the 3'-untranslated region of Na⁺/glucose cotransporter (SGLT1) mRNA. *FEBS Lett.* 492:233-237.
10. **Loflin, P. and J.E. Lever.** 2001. HuR binds a cyclic nucleotide-dependent stabilizing domain in the 3' untranslated region of Na⁺/glucose cotransporter (SGLT1) mRNA. *FEBS Lett.* 509:267-271.
11. **Murakami, Y., J.A. Huberman, and J. Hurwitz.** 1996. Identification, purification, and molecular cloning of autonomously replicating sequence-binding protein 1 from fission yeast *Schizosaccharomyces pombe*. *Proc. Natl. Acad. Sci. USA* 93:502-507.
12. **Ohta, T., K.J. Isselbacher, and D.B. Rhoads.** 1990. Regulation of glucose transporters in LLC-PK₁ cells: effects of D-glucose and monosaccharides. *Mol. Cell. Biol.* 10:6491-6499.
13. **Peng, H. and J.E. Lever.** 1995. Regulation of Na⁺-coupled glucose transport in LLC-PK₁ cells. Message stabilization induced by cyclic AMP elevation is accompanied by binding of an M_r = 48,000 protein to a uridine-rich domain in the 3'-untranslated region. *J. Biol. Chem.* 270:23996-24003.
14. **Peng, S.S., C.Y. Chen, N. Xu, and A.B. Shyu.** 1998. RNA stabilization by the AU-rich element binding protein, HuR, an ELAV protein. *EMBO J.* 17:3461-3470.
15. **Qian, Z. and J. Wilusz.** 1993. Cloning of a cDNA encoding an RNA binding protein by screening expression libraries using a northwestern strategy. *Anal. Biochem.* 212:547-554.
16. **Sagesser, R., E. Martinez, M. Tsagris, and M. Tabler.** 1997. Detection and isolation of RNA-binding proteins by RNA-ligand screening of a cDNA expression library. *Nucleic Acids Res.* 25:3816-3822.
17. **Shyu, A.B. and M.F. Wilkinson.** 2000. The double lives of shuttling mRNA binding proteins. *Cell* 102:135-138.
18. **Sikela, J.M. and W.E. Hahn.** 1987. Screening an expression library with a ligand probe: isolation and sequence of a cDNA corresponding to a brain calmodulin-binding protein. *Proc. Natl. Acad. Sci. USA* 84:3038-3042.
19. **Spicher, A., O.M. Guicherit, L. Duret, A. Aslanian, E.M. Sanjines, N.C. Denko, A.J. Giaccia, and H.M. Blau.** 1998. Highly conserved RNA sequences that are sensors of environmental stress. *Mol. Cell. Biol.* 18:7371-7382.
20. **Vinson, C.R., K.L. LaMarco, P.F. Johnson, W.H. Landschulz, and S.L. McKnight.** 1988. In situ detection of sequence-specific DNA binding activity specified by a recombinant bacteriophage. *Genes Dev.* 2:801-806.
21. **You, Y., C.Y. Chen, and A.B. Shyu.** 1992. U-rich sequence-binding proteins (URBPs) interacting with a 20-nucleotide U-rich sequence in the 3' untranslated region of *c-fos* mRNA may be involved in the first step of *c-fos* mRNA degradation. *Mol. Cell. Biol.* 12:2931-2940.
22. **Young, R.A. and R.W. Davis.** 1983. Efficient isolation of genes by using antibody probes. *Proc. Natl. Acad. Sci. USA* 80:1194-1198.

This work was supported by the National Institutes of Health grant no. DK27400 to J.E.L. We thank Dr. Ann-Bin Shyu for the plasmid pMyc-over and Constance J. Clancey and Zheng Li for library screening and sequence analysis. Address correspondence to Dr. Julia Lever, Department of Biochemistry & Molecular Biology, University of Texas-Houston Medical School, P.O. Box 20708, Houston, TX 77225, USA. e-mail: julia.e.lever@uth.tmc.edu

Received 29 October 2001; accepted 11 January 2002.

Paul Loflin and Julia E. Lever
*University of Texas-Houston
Medical School
Houston, TX, USA*

For reprints of this or
any other article, contact
Reprints@BioTechniques.com