Short Technical Reports

different mutagenesis reactions. We conclude that the high fidelity, low background, and the relatively short hands-on time make TPS a useful alternative mutagenesis procedure.

REFERENCES

1. Barlet, I., F. Maurier, M. Duchesne, E. Guittard, B. Tocque, and F. Schweighoffer. 1997. A role for Sam68 in cell cycle progression antagonized by a spliced variant within the KH domain. *J. Biol. Chem.* 272:3129-3132.
2. Bauer, C.E., S.D. Hesse, D.A. Waechter-Brulla, S.P. Lynn, R.J. Gumpert, and J.F. Gardner. 1985. A genetic enrichment for mutations constructed by oligodeoxynucleotide-directed mutagenesis. *Gene* 37:73-81.
3. Bernard, P., K.E. Kézdy, L.V. Melder, J. Steyaert, L. Vyns, M.L. Pato, P.N. Higgins, and M. Couturier. 1993. The F plasmid CcdB protein induces efficient ATP-dependent DNA cleavage by gyrase. *J. Mol. Biol.* 234:534-541.
4. Boren, J., I. Lee, M.J. Callow, E.M. Rubin, and T.L. Innerarity. 1996. A simple and efficient method for making site-directed mutants, deletions, and fusions of large DNA such as P1 and BAC clones. *Genome Res.* 6:1123-1130.
5. Deng, W.P. and J.A. Nickoloff. 1992. Site-directed mutagenesis of virtually any plasmid by eliminating a unique site. *Anal. Biochem.* 200:81-88.
6. Haas, J., E.C. Park, and B. Seed. 1996. Codon usage limitation in the expression of HIV-1 envelope glycoprotein. *Curr. Biol.* 6:315-324.
7. Hutchison, C.A., S. Phillips, M.H. Edgell, S. Gillam, and M. Smith. 1978. Mutagenesis at a specific position in a DNA sequence. *J. Biol. Chem.* 253:6551-6560.
8. Inouye, S. and M. Inouye. 1987. Oligonucleotide-directed site-specific mutagenesis using double-stranded plasmid DNA, p. 181-206. In S.A. Narang (Ed.), *Synthesis and Application of DNA and RNA*. Academic Press, New York.
9. Jaffé, A., T. Ogura, and S. Hiraga. 1985. Effects of the ccd function of the F plasmid on bacterial growth. *J. Biol. Chem.* 163:841-849.
10. Kunkel, T.A. 1985. Rapid and efficient site-specific mutagenesis without phenotypic selection. *Proc. Natl. Acad. Sci. USA* 82:488-492.
11. Lewis, M.K. and D.V. Thompson. 1990. Efficient site directed in vitro mutagenesis using ampicillin selection. *Nucleic Acids Res.* 18:3439-3443.
12. Reddy, T.R., W. Xu, J.K. Mau, C.D. Goodwin, M. Suhasini, H. Tang, K. Frimpong, D.W. Rose, and F. Wong-Staal. 1999. Inhibition of HIV replication by dominant negative mutants of Sam68, a functional homolog of HIV-1 Rev. *Nat. Med.* 5:635-642.
13. Reddy, T.R. 2000. A single point mutation in the nuclear localization domain of Sam68 blocks the Rev/RRE-mediated transactivation. *Oncogene* 19:3110-3114.
14. Taylor, J.W., J. Ott, and F. Eckstein. 1985. The rapid generation of oligonucleotide-directed mutations at high frequency using phosphorothioate-modified DNA. *Nucleic Acids Res.* 13:8765-8785.
15. Taylor, S.J. and D. Shalloway. 1994. An RNA-binding protein associated with Src through its SH2 and SH3 domains in mitosis. *Nature* 368:867-871.
16. Xu, W., B.A. Smith-Franklin, P.L. Li, C. Wood, J. He, Q. Du, G.J. Bhat, C. Kankasa, et al. 2001. Potent neutralization of primary human immunodeficiency virus clade C isolates with a synergistic combination of human monoclonal antibodies raised against clade B. *J. Hum. Virol.* 4:55-61.

This work was supported by National Institutes of Health grant no. 19972 to B.A.M., GM56089 to F.W.-S., AI46240 to T.R.R., and P01AI48240 to R.M.R. The authors thank Susan Sharp for her assistance in the preparation of this manuscript. Address correspondence to Dr. Ruth M. Ruprecht, Dana-Farber Cancer Institute, 44 Binney Street, Boston, MA 02115, USA. e-mail: ruth_ruprecht@dfci.harvard.edu

Received 20 November 2001; accepted 17 January 2002.

W. Xu^{1,2,3}, Y. Zhang¹, L.-Y. Yeh¹, C.R. Ruprecht¹, F. Wong-Staal², B.A. McFadden³, T.R. Reddy⁴, and R.M. Ruprecht¹
¹Dana-Farber Cancer Institute
Boston, MA
²University of California
at San Diego
La Jolla, CA
³Washington State University
Pullman, WA
⁴Wayne State University
Detroit, MI, USA

For reprints of this or
any other article, contact
Reprints@BioTechniques.com

Minimal FLAG Sequence Useful in the Functional Epitope Tagging of H-Ras

BioTechniques 32:1270-1280 (June 2002)

ABSTRACT

Epitope tagging can interfere with normal protein function, indicating the need for an unobtrusive epitope tag. The FLAG[®] epitope (DYKDDDDK) was examined for a minimal epitope useful in the tagging of H-Ras. The heptapeptide tag, F7 (MDYKDD), was found to retain reactivity with M2 and M5 monoclonal antibodies in immunoprecipitation, Western blotting, and immunofluorescence microscopy. The F7 tag did not interfere with Ras stability, EGF stimulation of Ras activation, and downstream phosphorylation of MAPK Erk1/2. Unlike the full FLAG sequence, the F7 tag had minimal effect on the growth properties of H-Ras in a colony-forming assay. The F7 tag may be useful when minimizing the effect of tagging on protein function is an important criterion in the selection of an N-terminal epitope tag.

INTRODUCTION

A common approach to studying a specific cellular protein is to transfect the gene of interest into cultured mammalian cells and examine functional properties of the expressed protein. The incorporation of an epitope tag on the protein provides a convenient handle for isolating and studying the exogenous protein. In principle, any short oligopeptide may serve as an epitope tag. However, oligopeptides that serve as efficient immunogens are more practical for obtaining peptide-specific monoclonal antibodies of high affinity (6). In addition, the epitope tag must localize to the surface of the attached protein to be accessible to antibodies and should not interfere with the normal function of the target protein. Occasionally, protein tags alter the conformation and function of the fusion protein. For instance, the 11-amino-acid substance P tag has been shown to interfere with the normal structure of intermediate filament proteins (2). In the study of the small G-protein Ras, the GST and His tags were found to

Short Technical Reports

reduce the biological activity of Ras in microinjection studies (10), while GFP was found to increase its stability (9). Thus, caution should be exercised in selecting a protein tag for Ras and perhaps other small G-proteins. One way to minimize obtrusive properties of epitope tags is to minimize the length of the tag. Recently, the smaller EE and Myc tags appended to Ras proteins were used in the study of its functional properties (7,14). However, the impact of epitope tagging on Ras function was not the point of these studies, and how tagging affects its function remains unclear.

A frequently used epitope tag is the FLAG[®] sequence (DYKDDDDK) (6). The monoclonal antibodies, M1 and M2, react with this sequence, while the M2 and M5 antibodies react with its N-terminal methionine variant (MDYKDDDDK). Although the FLAG sequence is often chosen as an epitope tag, the highly charged nature of this epitope

occasionally induces deleterious effects on fusion proteins altering their normal characteristics. For example, fusion of the FLAG sequence to the membrane-binding subunit of the influenza virus haemagglutinin (HA2) causes the protein to localize to the cytoplasm of cells rather than the membrane (3). The FLAG sequence likely increased the hydrophilicity of HA2, reducing its ability to localize to the cell membrane. Epitope tag interference of fusion proteins incorporating the FLAG sequence and other larger tags indicates a need for a less obtrusive epitope tag for the study of sensitive proteins.

In recent experiments, we transfected NIH3T3 cells with FLAG-H-Ras fusion constructs and observed a substantial population increase during G418 selection as compared to cells transfected with H-Ras or vector alone, suggesting alteration in H-Ras activity caused by the FLAG sequence. In addi-

tion, the mobility of FLAG-H-Ras during SDS-PAGE was retarded relative to H-Ras, possibly indicating structural alterations. For this reason, we sought to develop a smaller, less interfering epitope tag. We describe the use of the F7 heptapeptide, MDYKDDD, to reduce the interference of normal H-Ras activity while maintaining reactivity with M2 and M5 antibodies in Western blotting, immunoprecipitation, and immunofluorescent staining of cells.

MATERIALS AND METHODS

Cell Culture

NIH3T3 (CRL 1658) and COS-1 (CRL 1650) cells were obtained from ATCC (Manassas, VA, USA) and cultured in DMEM supplemented with 10% FBS, 100 U/mL penicillin, and 100 µg/mL streptomycin (Invitrogen,

Carlsbad, CA, USA). Cells were grown in an atmosphere of 5% CO₂ at 37°C and passaged every 3–4 days using 0.05% trypsin, 0.53 mM EDTA (trypsin-EDTA; Invitrogen) to detach cells from tissue culture plastic.

Epitope-Tagged H-Ras Constructs and Cell Transfections

To generate FLAG-tagged Ras, we first constructed the pcF vector by inserting the FLAG epitope (MDYKD DDDK) into the pcDNA3 vector (Invitrogen) by ligating annealed, complementary DNA strands containing the coding sequence for FLAG and overhangs compatible with *Hind*III and *Kpn*I restriction endonucleases (5'-AGCTTATGGACTACAAGGACGACGATGAC-AAGGTAC-3' and 5'-CTTGTCATCGTCGTCCTGTAGTCCATA-3'). H-Ras cDNA was cloned using PCR from placental mRNA and primers flanked by *Kpn*I and *Bam*HI into pcF. Shorter FLAG variants (F6 and F7) were constructed by PCR of Ras sequences using primers that incorporated the required epitope sequence and *Hind*III and *Bam*HI restriction sites (F6-1-H-Ras forward primer: 5'-CCCAAGCTTATGGACTACAAGGACGACACTGAATAAACTG-3'; F6-3-H-Ras forward primer: 5'-CCCAAGCTTATGGACTACAAGGACGACTATAAACTGGTTCG-3'; F7-1-H-Ras forward primer: 5'-CCCAAGCTTATGGACTACAAGGACGACGATACTGAATATAAACTG-3'; F7-3-H-Ras forward primer: 5'-CCCAAGCTTATGGACTACAAGGACGACGATTATAAACTGGTGG-3'; H-Ras reverse primer: 5'-CCCGGGATCCTCAGGAGAGCACACACTTGCAGC-3'). The PCR products were digested with *Hind*III and *Bam*HI and ligated into similarly digested pcDNA3. For transient transfection, NIH3T3 and COS-1 cells were plated at approximately 60% confluence in 35-mm plates. One day after plating, each plate of cells was transfected with 6 µL LIPOFECTAMINE® reagent (Invitrogen) and 2 µg cDNA in serum-free medium. After 18 h, the transfection solution was replaced with complete medium and incubated for an additional 24 h before analysis. For epidermal growth factor (EGF) stimulation, cells were rendered quiescent by incubating for an additional 24 h in DMEM, 0.5% FBS.

Immunoprecipitation

Transfected cells were washed with PBS and then lysed in TBS-TDS (10 mM Tris-HCl, pH 7.4, 140 mM NaCl, 1% Triton® X-100, 0.5% sodium deoxycholate, 0.1% SDS) on ice for 30 min. The cell lysates were cleared by centrifugation at 16000×g for 5 min. The supernatants were incubated with 1 µg antibody and 10 µL 50% agarose-Protein G slurry (Sigma, St. Louis, MO, USA) for 2 h at 4°C. The antibodies used were Y13-259 anti-pan-Ras (CRL 1742; ATCC), Y13-238 anti-H-Ras (CRL 1741; ATCC), M2 anti-FLAG and M5 anti-FLAG (Sigma). The immunoprecipitates were collected by centrifugation at 16000×g for 1 min and washed three times with TBS-TDS. The pellet was resuspended in 10–20 µL sample buffer (125 mM Tris-HCl, pH 6.8, 1% SDS, 5% 2-mercaptoethanol, 0.001% bromophenol blue) and analyzed by Western blotting as described below.

Western Blotting

Protein samples were separated on 12.5% polyacrylamide SDS-denaturing mini gels (Bio-Rad Laboratories, Hercules, CA, USA) and then electroblotted onto PVDF membranes (Millipore, Bedford, MA, USA). Membranes were blocked in 5% skim milk (Difco, Detroit, MI, USA) in PBST (10 mM NaPO₄, pH 7.4, 140 mM NaCl, 0.5% Tween® 20) for 30 min and incubated overnight at 4°C with 50 ng/mL antibody in PBST, 1% BSA, followed by goat anti-mouse conjugated to HRP (Sigma) at a dilution of 1:20 000 in PBST, 1% BSA for 1 h at room temperature. The antibodies used were M2, M5, and LA069 anti-H-Ras (Quality Biotech, Camden, NJ, USA). Membranes were developed using ECL® (Amersham Biosciences, Piscataway, NJ, USA), and reactive proteins were detected by exposure to X-ray film (Pierce Chemical, Rockford, IL, USA).

Immunofluorescence Microscopy

NIH3T3 cells in 35-mm dishes were transfected with pcDNA3 and constructs containing H-Ras and epitope-tagged H-Ras as described above. Cells were washed with PBS, fixed with

3.7% formaldehyde for 15 min, washed with PBS, permeabilized with PBS, 0.1% Triton X-100 for 10 min, washed with PBS, and then blocked with PBS, 1% BSA for 20 min. Control cells remained in blocking solution. Test cells were incubated with 5 mg/mL monoclonal antibody (LA069, M2, or M5) in 1% BSA-PBS for 30 min, washed with PBS, and then incubated with 25 mg/mL goat anti-mouse conjugated to rhodamine (Jackson ImmunoResearch, West Grove, PA, USA) in PBS, 1% BSA for 30 min. The secondary antibody was also applied to control cells to test for background immunofluorescence. The cells were then washed with PBS, post-fixed with 3.7% formaldehyde for 15 min, rinsed with PBS, and mounted in 10% PBS, 90% glycerol under coverslips. Fluorescence was viewed using a Zeiss Axioplan fluorescence microscope (Carl Zeiss, Thornwood, NY, USA) with a 546-nm excitation filter and a 590-nm barrier.

Ras Activation Assay

COS-1 cells were transfected with H-Ras, FLAG-H-Ras, and F7-1-H-Ras constructs as described above and then serum-starved for 24 h. Cells were stimulated for 5 min in DMEM, 0.05% BSA with or without 50 ng/mL EGF (Collaborative Research, Bedford, MA, USA). EGF-stimulated cells were lysed for 20 min on ice in 1 mL co-precipitation lysis solution (30 mM HEPES, pH 7.5, 1% Triton X-100, 10 mM sodium chloride, 10% glycerol, 1 mM EGTA, 25 mM sodium fluoride, 1 mM sodium orthovanadate, 10 mM β-glycerophosphate, 10 mM benzamide, 0.1 mM PMSF, 10 µg/mL aprotinin, 10 µg/mL leupeptin, 10 µg/mL trypsin inhibitor). Lysates were cleared by centrifugation at 4°C for 5 min at 16000×g. To prepare a Ras activation probe (5), cDNA corresponding to the Ras-binding domain (RBD) of B-Raf (amino acids 149–226) was generated by reverse transcription-coupled PCR (forward primer: 5'-CATATGGGATCCAAGTCAACACAAAACCTATC-3'; reverse primer: 5'-CTCGAGAATTCCTCCAACTTCCACATGCAAT-3') of placental mRNA that incorporated flanking *Bam*HI and *Xho*I restriction sites. The PCR product was ligated into

Short Technical Reports

appropriately digested pGEX-4T-1 (Amersham Biosciences). The glutathione S-transferase/B-Raf RBD (GST-RBD) fusion protein was expressed in *E. coli* strain BL-21 (Stratagene, La Jolla, CA, USA), and the GST-RBD was purified by binding to glutathione covalently linked to agarose (GSH-agarose). The amount of GST-RBD bound to GSH-agarose was determined by quantification of Coomassie®-stained polyacrylamide gels using BSA as a standard. Ras was co-precipitated with 10 µg GST-RBD bound to GSH-agarose beads for 1 h at 4°C. The Ras/GST-RBD complexes were washed four times in co-precipitation wash buffer (50 mM HEPES, pH 7.5, 100 mM NaCl, 0.1% Triton X-100, 10% glycerol, 20 mM sodium fluoride). The remaining inactivated Ras proteins were recovered from the supernatant by immunoprecipitating with for 2 h at 4°C with 1 µg Y13-238 and 10 µL 50% slurry of Protein G conjugated to agarose beads. Precipitates were washed three times in co-precipitation wash buffer. Activated (Ras/GST-RBD co-precipitation) and inactivated (Y13-238 recovery immunoprecipitation) Ras precipitates were solubilized in sample buffer and analyzed by Western blotting as described above using M2 and LA069 antibodies. To monitor Erk activation in the same lysates used for the Ras activation assay, 100 µL of the cell lysate was removed before the addition of GST-RBD and adjusted to contain sample buffer. The lysates were analyzed by Western blotting as described above using rabbit polyclonal Erk (p44/42 MAP kinase) and phospho-Erk (phospho-p44/42 MAP kinase) antibodies (Cell Signaling, Beverly, MA, USA), followed by HRP-conjugated goat anti-rabbit IgG secondary antibody (Sigma). For quantification, images of blots were digitized using a flatbed scanner (Umax S-12; Umax Datasystems, Taiwan, R.O.C.) and analyzed using Image 1.60 densitometry software (National Institutes of Health, Bethesda, MD, USA).

Ras Stability

NIH3T3 cells were transfected with pcDNA3 and H-Ras constructs as described above. Cells were then treated with 10 µg/mL cycloheximide (Sigma)

for 24 h. Cells were detached with trypsin-EDTA and pooled with any unattached cells from the culture medium. Cells were rinsed in PBS and lysed in TBS-TDS for 30 min on ice. Lysates were cleared by centrifugation at 4°C for 5 min at 16000× g. H-Ras was immunoprecipitated with 1 µg Y13-238 and 10 µL 50% slurry of agarose-protein G for 4 h at 4°C. Precipitates were washed three times in TBS-TDS, dissolved in sample buffer, and analyzed by Western blotting using LA069 and M2 antibodies. Results were quantified by densitometry as described above.

Colony-Forming Assay

NIH3T3 cells were transfected as described above. A mock control was

Table 1. Construction of Epitope Tags

Tag	Sequence
FLAG	DYKDDDDK VP MEYKLV
F7-1	MDYKDD DT EYKLV
F7-3	MDYKDD DD EYKLV
F6-1	MDYKDD DT EYKLV
F6-3	MDYKDD DD EYKLV

Bolded residues represent N-terminal H-Ras amino acids.

included that omitted DNA in the transfection procedure. Twenty-four hours after transfection, cells were re-plated into four 100-mm plates in growth medium containing 400 µg/mL G418 (Mediatech, Herndon, VA, USA) and

Figure 1. Immunoprecipitation and Western blot analysis of tagged Ras. NIH3T3 cells were transfected with pcDNA3-H-Ras (H/H-Ras), pcDNA3-FLAG-H-Ras (FLG/FLAG), pcDNA3-F7-1-H-Ras (F7-1), pcDNA3-F7-3-H-Ras (F7-3), pcDNA3-F6-1-H-Ras (F6-1), or pcDNA3-F6-3-H-Ras (F6-3). (A) Western blotting. Ras proteins were immunoprecipitated from cell lysates with Y13-259 (anti-pan Ras), separated on 12.5% polyacrylamide gels and transferred to PVDF membranes. Membranes were probed with LA069 anti-H-Ras, M2 and M5 antibodies, followed by HRP-conjugated anti-mouse IgG. Reactive proteins were discerned using ECL. (B) Immunoprecipitation. Ras proteins were immunoprecipitated from cell lysates using Y13-259, M2, and M5 antibodies. Western blotting was performed as above using LA069. Data are representative of three independent experiments.

Figure 2. Immunofluorescence microscopy of tagged Ras. NIH3T3 cells were transfected as in Figure 1. Twenty-four hours after transfection, cells were fixed and stained with LA069, M2, and M5 monoclonal antibodies, followed by goat anti-mouse IgG conjugated to rhodamine. Stained cells were imaged using a fluorescence microscope. Cells representative of staining observed for each population are presented.

G418 selection continued for 10 days. Colonies were stained with 0.2% methylene blue, 50% ethanol for 30 min, washed three times in PBS, and individual colonies were counted.

RESULTS AND DISCUSSION

Epitope Tagging of H-Ras

To study the biology of exogenously expressed Ras proteins, Ras was tagged with the FLAG epitope (6). During the course of our studies, it was observed that compared with cells transfected with vector or H-Ras, FLAG-H-Ras increased the number of cells during G418 selection (data not shown). This increase in cell growth suggested that the FLAG tag might be interfering with normal H-Ras activity. It was reasoned that a shorter FLAG sequence appended to H-Ras might mini-

mize its interference with normal function while retaining reactivity with the M2 and M5 antibodies. Previously, it was demonstrated that certain amino acid residues within the FLAG sequence are not essential for reactivity with anti-FLAG antibodies (12). Within the N-terminal methionine FLAG sequence, MDYKDDDDK, it was found that residues D2, Y3, K4, and D7 are the most critical for reactivity with M2, while residues D2, Y3, D5, and D6 are most critical for reactivity with M5. These data indicate that residues D8 and K9 are not critical for reactivity with the M2 and M5 anti-FLAG antibodies. In contrast, the studies of Knappik and Plückthun (8) examined the reactivity of M2 with variants of the FLAG sequence fused to the N-terminus of a single-chain antibody. In their studies, M2 failed to react with the sequence DYKDED, which maintains the D2, Y3, K4, and

D7 residues found to be important in the studies of Slootstra et al. (12). Moreover, M2 recognized the sequence DYKDEVKD, which lacks the critical residue D7. The contrasting results of these two studies may relate to the fact that one used an ELISA of peptides immobilized to polyethylene, while the other used Western blotting of tags fused to proteins. Thus, we decided to take an empirical approach to determining a minimal FLAG sequence.

A series of fusion proteins were designed that have six- or seven-amino-acid truncated versions of the FLAG sequence (Table 1). To minimize potential interference due to the length of overhang of the N-terminus, the tags were embedded one or three amino acids into the N-terminus of H-Ras. Based on the 3-D structure of Ras, the first three amino acids (MTE) play a minor structural role. Most notably, threonine-2 interacts with β -strand-3

Short Technical Reports

and may contribute to stability. It was reasoned that it might be possible to alter these N-terminal residues without significant consequences to the structure and function of the protein. The seven-amino-acid tags F7-1 and F7-3 both consisted of residues MDYKDDD and were embedded one or three residues, respectively, into the N-terminus of Ras (Table 1). The six-amino-acid tags F6-1 and F6-3 both consisted of residues MDYKDD and were embedded one or three residues, respectively, into the N-terminus of H-Ras (Table 1). It should be noted that removal of the last amino acid (K9) effectively removes the enterokinase recognition sequence (DDDDK) engineered into the FLAG tag (6). These constructs were tested for their ability to retain reactivity to anti-FLAG antibodies in several commonly used procedures.

Western Blotting

First, we sought to determine the usefulness of the smaller FLAG variants in Western blotting analysis. Immunoprecipitated Ras was immunoblotted using LA069 (anti-H-Ras), M2, or M5. While both endogenous and exogenous Ras proteins were immunoprecipitated, only the exogenous, tagged proteins will produce a positive signal on the M2 and M5 Western blots. The amount of Ras loaded onto each gel is the same as shown in the top panel of Figure 1A (LA069). The results shown in Figure 1A demonstrate that the F7 tag retained reactivity with M2 and M5 in immunoblot analysis. In contrast, the F6 tag showed weak to negligible reactivity with M2 and M5, indicating the importance of the D7 residue. A significant reduction in H-Ras mobility (apparent mass of H-Ras = 22.0 kDa) during SDS-PAGE was seen with the FLAG epitope (also see Figure 4). The mobility reduction corresponds to a 3.3-kDa mass shift, greater than 1.2 kDa ($\Delta = 2.1$ kDa), as predicted by the mass increase due to the addition of the FLAG peptide. This shift was reduced with the abbreviated tags. The F7 tag migrated with an apparent mass shift of 1.4 kDa compared to its predicted increase of mass of 0.8 kDa ($\Delta = 0.6$ kDa). It is not clear whether the decreased mobility caused

by the FLAG tag is due to structural alteration of H-Ras or is simply an artifact of SDS-PAGE. We have previously observed a similar reduction of mobility of Ras proteins during SDS-PAGE following ADP-ribosylation by exoenzyme S (13). The epitope tags that were more deeply embedded into the Ras sequence, F7-3 and F6-3, showed reduced reactivity as compared to F7-1 and F6-1, respectively, suggesting that embedding of the tag reduced its accessibility.

Immunoprecipitation

To examine the ability of M2 and M5 to react with the FLAG-derived sequences in solution, H-Ras fusion proteins were immunoprecipitated from lysates of transiently transfected NIH3T3 cells using Y13-259, M2, and M5 antibodies. The immunoprecipitated proteins were immunoblotted with LA069 anti-H-Ras (Figure 1B). The data demonstrate that the F7 epitope is sufficient for immunoprecipitation by ei-

Figure 3. Effect of FLAG and F7 tags on H-Ras activity. COS-1 cells were transfected with pcDNA3-H-Ras (H), pcDNA3-FLAG-H-Ras (FLAG) and pcDNA3-F7-1-H-Ras (F7-1) and rendered quiescent. Cells were stimulated with 50 ng/mL EGF (+) or mock stimulated (-) for 5 min. (A) Activated Ras was co-precipitated with GST-RBD and (B) inactivated Ras recovered from the GST-RBD co-precipitation supernatants by immunoprecipitation with Y13-238 anti-H-Ras antibody. Activated and unactivated Ras were detected by Western blotting as described in Figure 1 using M2 antibody. Blots were then re-probed with LA069 antibody. (C) Erk activation was determined by Western blotting of cell lysates used to determine Ras activation. Erk blots were probed with phospho-Erk specific antibodies (P-Erk) and re-probed with Erk-specific antibodies (Erk). (D) The protein blots were analyzed by densitometry and are displayed graphically. The amount of Ras or Erk activated by EGF is presented as the percent activated relative to mock-stimulated control. The data are the mean (columns) and standard deviation (error bars) from three independent experiments. The line indicates the relative amount of activated Ras and phosphorylated Erk in unstimulated cell populations.

Short Technical Reports

ther M2 or M5 antibodies. In contrast, the F6 tags only weakly reacted with the M2 and M5 antibodies. Based on the intensity of the Western blots, the M5 antibody appears to be less efficient than M2 in immunoprecipitating tagged H-Ras.

Immunofluorescence Microscopy

We next examined the usefulness of the F6 and F7 tags in immunofluorescence analysis. For this study, NIH3T3 cells were transiently transfected with the H-Ras constructs as indicated (Figure 2). Cells were then stained with LA069, M2, or M5, followed by a rhodamine-conjugated secondary antibody. The staining pattern of H-Ras and tagged H-Ras is similar to that observed previously for overexpressed H-

Ras (1,4), indicating that the epitope tag does not interfere with the localization of the fusion protein. Images of positively stained cells indicate that the F7 tags are accessible to both M2 and M5 anti-FLAG antibodies. Although the staining patterns of F6-tagged H-Ras were similar, the signal was greatly reduced as compared to the F7 signals. The intensity of the fluorescent signal is the greatest in those cells transfected with the F7-1 construct, indicating that this tag is the most compatible immunofluorescence analysis.

Ras Activation

Of the four variant tags tested (F7-1, F7-3, F6-1, and F6-3), F7-1 displayed the strongest reactivity to M2 and M5

antibodies and thus appeared to be the most useful. To evaluate whether appending the F7 tag to H-Ras interfered with its biological activity, the effect of tagging on EGF stimulation of H-Ras activation was examined. The Ras binding domain of B-Raf was used as a Ras-GTP-dependent probe (5). As can be seen in Figure 3A, H-Ras, FLAG-H-Ras, and F7-1-H-Ras were each activated in response to EGF with comparable increase in stimulation levels relative to unstimulated. To ensure that downstream signaling of Ras remained intact, we examined the phosphorylation status of the MAPK, Erk. Ras signals through Raf and Mek1/2 to activate Erk1/2. A portion of the cell extract used to monitor Ras activation was used to examine Erk phosphorylation. Figure 3B demonstrates that FLAG and F7-1 tagged H-Ras do not interfere with Erk phosphorylation. These data indicate that EGF/Ras/Erk signaling remains intact in cells expressing FLAG-H-Ras and F7-1-H-Ras.

Ras Stability

It has previously been observed that GFP fused to the N-terminus of Ras protein increases its stability by interfering with its normal turnover (9). It

Figure 4. Effect of FLAG and F7-1 tags on Ras stability. NIH3T3 cells were transfected as described in Figure 1. After allowing exogenous proteins to express, protein synthesis was inhibited by treatment with (+) 10 μ g/ μ L cycloheximide (CHX) for 24 h. Control cells were left untreated (-). (A) The amount of H-Ras was determined by Western blotting as described in Figure 1 using M2 antibody and reprobing the blot with LA069 antibody. Note that the two protein bands seen in the FLAG and F7-1 lanes of the re-probed membrane correspond to tagged (upper band) and endogenous (lower band) H-Ras. (B) The protein blots were analyzed by densitometry and the data displayed graphically. The data are presented as the percent of Ras remaining in cycloheximide-treated cells relative to untreated cells. The data are the mean (column) and standard deviation (error bars) from three independent experiments.

Figure 5. Effects of epitope tags on colony formation. NIH3T3 cells were transfected as described in Figure 1. A mock control was included that omitted DNA in the transfection procedure (Mock). Twenty-four hours after transfection, each plate of cells was re-plated at low density. Plasmid-containing colonies were selected with 400 μ g/ μ L of G418 for 10 days. Plates were stained with methylene blue, and individual colonies were counted. Data represent the mean and standard error of two independent experiments performed in quadruplicate.

Short Technical Reports

was of interest to determine whether epitope tagging of H-Ras altered its stability. To test this, NIH3T3 cells were transfected with vector (for comparison with endogenous H-Ras), H-Ras, FLAG-H-Ras, and F7-1-H-Ras. Following expression of exogenous Ras proteins, further protein synthesis was halted by the addition of cycloheximide. As shown in Figure 4, overexpressed H-Ras displayed a decreased turnover when compared to endogenous H-Ras (88% compared to 53%, respectively). This is likely due to the accumulation of H-Ras in the ER/Golgi complex during overexpression (1,4). Exogenous FLAG-H-Ras and F7-1-H-Ras displayed decreased stability relative to exogenous H-Ras but increased stability relative to endogenous H-Ras. The decreased turnover of exogenous Ras proteins likely reflects a combination of changes in rates of processing in the ER/Golgi and normal protein turnover determined by cellular levels of H-Ras. Nonetheless, epitope-tagged H-Ras was no less stable than endogenous H-Ras.

Colony-Forming Assay

The initial indication that the FLAG tag alters wild-type H-Ras function was an increase in cell number during G418 selection (data not shown). The effect of the F7-1 tag on normal H-Ras function was determined in a colony-forming assay. As can be seen in Figure 5, compared to the vector control, untagged H-Ras increased colony formation by 20%, F7-1-H-Ras by 60%, and FLAG-H-Ras by 217%. Wild-type H-Ras was not expected to have a large effect on colony formation. The F7 tag had only a moderate effect on wild-type H-Ras activity, while the full FLAG sequence caused a substantial increase in the number of G418-resistant colonies. The mechanism by which the FLAG tag affects H-Ras function in this assay is unknown. Its increased activity suggests either increased nucleotide exchange rate or decreased GTPase activity. However, neither FLAG-H-Ras nor F7-1-H-Ras displayed changes in EGF-stimulated activity (Figure 3). This suggests that tagging does not affect Ras-GTP levels. It remains possible that the FLAG epitope increases the interaction of H-Ras

with one of its effectors (e.g., PI3 kinase, RalGDS) or interferes with Ras-GAP stimulation of its GTPase activity.

These studies indicate that the F7 epitope tag can be useful in minimizing the effect of tagging on protein function. The F7 tag is applicable to immunoprecipitation, Western blotting, and immunofluorescence analysis. Based on previous studies (8,12) and the results presented here, it may be possible to use the sequence DYKDDD as a tag for secretory proteins following cleavage of the signal peptide. Although the lack of enterokinase sequence precludes the removal of the F7 tag, the reduced effect on biological activity may compensate for this by making it an unnecessary step. In the future, it may be possible to develop smaller N-terminal epitopes similar to the very small Glu-Glu-Phe C-terminal epitope (11). This is limited by the minimal epitope size required to obtain a useful monoclonal antibody. In the meantime, the F7 heptapeptide may be of value where both N-terminal tagging is possible and retention of normal protein function is required.

REFERENCES

1. Apolloni, A., I.A. Prior, M. Lindsay, R.G. Parton, and J.F. Hancock. 2000. H-ras but not K-ras traffics to the plasma membrane through the exocytic pathway. *Mol. Cell Biol.* 20:2475-2487.
2. Browning, E.T. and M. Ruina. 1984. Glial fibrillary acidic protein: norepinephrine stimulated phosphorylation in intact C-6 glioma cells. *J. Neurochem.* 42:718-726.
3. Chen, J., J.J. Skehel, and D.C. Wiley. 1998. A polar octapeptide fused to the N-terminal fusion peptide solubilizes the influenza virus HA2 subunit ectodomain. *Biochemistry* 37:13643-13649.
4. Choy, E., V.K. Chiu, J. Silletti, M. Feoktistov, T. Morimoto, D. Michaelson, I.E. Ivanov, and M.R. Phillips. 1999. Endomembrane trafficking of Ras: the CAAX motif targets proteins to the ER and Golgi. *Cell* 98:69-80.
5. de Rooij, J. and J.L. Bos. 1997. Minimal Ras-binding domain of Raf1 can be used as an activation-specific probe for Ras. *Oncogene* 14:623-625.
6. Hopp, T.P., K.S. Prickett, V.L. Price, R.T. Libby, C.J. March, D.P. Cerretti, D.L. Urdal, and P.J. Conlon. 1988. A short polypeptide marker sequence useful for recombinant protein identification and purification. *Biotechnology* 6:1204-1210.
7. Jones, M.K. and J.H. Jackson. 1998. Ras-GRF activates Ha-Ras, but not N-Ras or K-Ras 4B, protein *in vivo*. *J. Biol. Chem.* 273:1782-1787.
8. Knappik, A. and A. Plückthun. 1994. An improved affinity tag based on the FLAG peptide for the detection and purification of recombinant antibody fragments. *BioTechniques* 17:754-761.
9. Niv, H., O. Gutman, Y.L. Hennis, and Y. Kloog. 1999. Membrane interactions of a constitutively active GFP-Ki-Ras 4B and their role in signaling. *J. Biol. Chem.* 274:1606-1613.
10. Rodríguez-Viciana, P., P.H. Warne, A. Khwaja, B.M. Marte, D. Pappin, P. Das, M.D. Waterfield, A. Ridley, and J. Downward. 1997. Role of phosphoinositide 3-OH kinase in cell transformation and control of the actin cytoskeleton by Ras. *Cell* 89:457-467.
11. Skinner, R.H., S. Bradley, A.L. Brown, N.J.E. Johnson, S. Rhodes, D.K. Stammers, and P.N. Lowe. 1991. Use of the Glu-Glu-Phe C-terminal epitope for rapid purification of the catalytic domain of normal and mutant ras GTPase-activating proteins. *J. Biol. Chem.* 266:14163-14166.
12. Sloodstra, J.W., D. Kuperus, A. Plückthun, and R. Melen. 1997. Identification of new tag sequences with differential and selective recognition properties for the anti-FLAG monoclonal antibodies M1, M2 and M5. *Mol. Divers.* 2:156-164.
13. Vincent, T.S., J.E. Fraylick, E.M. McGuffie, and J.C. Olson. 1999. ADP-ribosylation of oncogenic Ras proteins by *Pseudomonas aeruginosa* exoenzyme S *in vivo*. *Mol. Microbiol.* 32:1054-1064.
14. Voice, J.K., R.L. Klemke, A. Le, and J.H. Jackson. 1999. Four human Ras homologs differ in their abilities to activate Raf-1, induce transformation, and stimulate cell motility. *J. Biol. Chem.* 274:17164-17170.

Address correspondence to Dr. Timothy S. Vincent, Department of Pathology and Laboratory Medicine, Medical University of South Carolina, 165 Ashley Avenue, Suite 309, Charleston, SC 29425, USA. e-mail: vincent@musc.edu

Received 8 December 2000; accepted 17 January 2002.

Kristy Y. Johnson, Liya Liu,
and Timothy S. Vincent
Medical University of South
Carolina
Charleston, SC, USA

For reprints of this or
any other article, contact
Reprints@BioTechniques.com