Short Technical Reports

cleic acid hybrids at the maximum concentration tested [6% (w/v)]. Like NaCl, SDS stabilizes hybrids, with 1% (w/v) SDS being equivalent to approximately 8 mM NaCl in its effects on hybrid melting temperature (rather than to 35 mM if fully ionized).

Under our conditions, the T_m^* for related but mismatched DNA/DNA hybrids was reduced by only 0.6°C/% nonhomology, less than the figure of 1.2°C adopted previously).

Nevertheless, mismatching significantly reduced the extent of hybridization: at 69% homology, the extent of hybridization was reduced by a factor of 5 compared with perfect hybrids and fell by a factor of 10 at 57% homology (not presented). This accords with earlier reports (3,11) that T_m reduction due to sequence divergence significantly reduces the rate of association. Thus, probe selectivity for perfect matches versus distantly related sequences (e.g., Southern blotting) is most probably due to a reduction in the rate/extent of hybridization (even after 16 h incubation) rather than to reduced hybrid stability per se.

A significant effect of SDS on hybridization background on nylon filters was observed. Low (0.6%) or high (6%) SDS concentrations equally reduced the background, as did NaCl (>100 mM). The mechanism is not known. We also noted that the sharpness of the temperature versus dissociation profile was influenced by salt concentration. At high salt, a temperature range of only 10°C could span fully hybridized to complete dissociation; at the lowest salt concentrations, temperature-dependent dissociation was gradual, spanning 25°C or more (data not presented).

We suggest that it is advantageous to perform hybridization in the presence of SDS, as proposed previously (6,13), calculating the T_m using the equivalence figure above (1% SDS = 8 mM NaCl) and selecting the hybridization temperature to be approximately 20°C below the T_m . The increased stability of RNA/DNA hybrids should allow stringency conditions (temperature or equivalent salt) to be increased by 10°C without loss of signal, although caution is advocated if the T_m^* is, as we suggest, markedly (15°C–17°C) below the calculated T_m (Reference 10 and this work).

REFERENCES

1. **Anderson, M.L.M. and B.D. Young.** 1985. Quantitative filter hybridization, p. 73-111. *In* B.D. Hames and S.J. Higgins (Eds.), *Nucleic Acid Hybridisation: A Practical Approach*. IRL Press, Oxford.
2. **Beltz, G.A., K.A. Jacobs, T.H. Eickbush, P.T. Cherbas, and F. Kafatos.** 1983. Isolation of multigene families and determination of homologies by filter hybridization methods. *Methods Enzymol.* 100B:266-285.
3. **Bonner, T.L., D.J. Brenner, B.R. Neufeld, and R.J. Britten.** 1973. Reduction in the rate of DNA reassociation by sequence divergence. *J. Mol. Biol.* 81:123-135.
4. **Britten, R.J., D.E. Graham, and B.R. Neufeld.** 1974. Analysis of repeating DNA sequences by reassociation. *Methods Enzymol.* 29E:363-418.
5. **Britten, R.J. and E.H. Davidson.** 1985. Hybridization strategy, p. 3-15. *In* B.D. Hames and S.J. Higgins (Eds.), *Nucleic Acid Hybridisation: A Practical Approach*. IRL Press, Oxford.
6. **Church, G.M. and W. Gilbert.** 1984. Genomic sequencing. *Proc. Natl. Acad. Sci. USA* 81:1991-1995.
7. **Lathe, R.** 1985. Synthetic oligonucleotide probes deduced from amino acid sequence data. Theoretical and practical considerations. *J. Mol. Biol.* 183:1-12.
8. **Richardson, M., D. Redmond, C.J. Watson, and J.O. Mason.** 1999. Mouse Wnt8B is expressed in the developing forebrain and maps to chromosome 19. *Mamm. Genome* 10:923-925.
9. **SantaLucia, J., Jr., H.T. Allawi, and P.A. Seneviratne.** 1996. Improved nearest-neighbor parameters for predicting DNA duplex stability. *Biochemistry* 35:3555-3562.
10. **Schildkraut, C. and S. Lifson.** 1965. Dependence of the melting temperature of DNA on the salt concentration. *Biopolymers* 3:195-208.
11. **Sutton, W.D. and M. McCallum.** 1971. Mismatching and the reassociation rate of mouse satellite DNA. *Nat. New Biol.* 232:83-85.
12. **Thomas, C.A.J. and B.M. Dancis.** 1973. Ring stability. *J. Mol. Biol.* 77:43-55.
13. **Virca, G.D., W. Northemann, B.R. Shiels, G. Widera, and S. Broome.** 1990. Simplified northern blot hybridization using 5% sodium dodecyl sulfate. *BioTechniques* 8:370-371.
14. **Wang, J. and G.M. Shackleford.** 1996. Murine Wnt10a and Wnt10b: cloning and expression in developing limbs, face and skin of embryos and in adults. *Oncogene* 13:1537-1544.
15. **Williams, A.P., C.E. Longfellow, S.M. Freier, R. Kierzek, and D.H. Turner.** 1989. Laser temperature-jump, spectroscopic, and thermodynamic study of salt effects on duplex formation by dGCATGC. *Biochemistry* 28:4283-4291.
16. **Wood, W.L., J. Gitschier, L.A. Lasky, and R.M. Lawn.** 1985. Base composition-independent hybridization in tetramethylammonium chloride: a method for oligonucleotide screening of highly complex gene libraries. *Proc. Natl. Acad. Sci. USA* 82:1585-1588.

We would like to thank Melville Richardson for advice and assistance. This work was supported by the BBSRC and DEFRA. Address correspondence to Dr. Richard Lathe, Dept. Biomedical Sciences, Faculty of Medicine, University of Edinburgh, Hugh Robson Building, George Square, Edinburgh EH8 9XD, UK. e-mail: rlathe@ed.ac.uk

Received 9 January 2002; accepted 13 March 2002.

Ken Rose, John O. Mason, and Richard Lathe
University of Edinburgh, Edinburgh, UK

RT-PCR Heteroduplex Analysis Permits Differentiation of Transgene and Host Gene Expression in a Transgenic Animal Model

BioTechniques 33:58-66 (July 2002)

ABSTRACT

In transgenic animal models, the conservation of DNA sequences between the transgene and the host wild-type gene can complicate the evaluation of the expression of each gene. The potential for gene silencing may complicate matters further. Here we report the use of RT-PCR heteroduplex analysis to differentiate the expression of a transgene and its homologous wild-type, even when these genes are very similar in their respective DNA sequences. We designed RT-PCR primers to amplify identically sized 243-bp fragments within the DNA binding domain of the p53 gene from both human and mouse mRNA samples. Ten samples from human p53 (273H) transgenic mice and 10 samples from wild-type controls were tested. Heteroduplex bands were formed in all transgenic samples but were absent from all wild-type samples. In addition, RT-PCR heteroduplex analysis was

Short Technical Reports

able in one sample to differentiate a silenced transgene from its wild-type allele, without the assistance of sequencing or labeling. In summary, the RT-PCR heteroduplex analysis is easy to use and has the ability to screen a large number of samples in a short time. The RT-PCR heteroduplex analysis is especially useful for the detection of expression when a transgene and the host homologous endogenous allele are too conserved in sequence to design species-specific RT-PCR primers.

INTRODUCTION

The development of animal models through transgenic technology is one of the most frequently used techniques to study the relationship between human disease and gene function and regulation in vivo. The technique consists of injecting the target gene into the pronucleus of an animal fertilized oocyte,

where it integrates randomly in the genome. The transgene replicates together with the host endogenous chromosome during mitosis and meiosis (6), and its subsequent expression produces a functional protein that may contribute to the phenotype of the animal. In many cases, the host animal has a wild-type gene that is homologous to the exogenous transgene. A nucleotide sequence that is highly conserved between a transgene and a wild-type gene may cause difficulty in the detection of the expression of the transgene. Evaluating for the presence of the final product of the transgene, the protein is able in many instances to circumvent this problem. However, species- and gene-specific antibodies capable of identifying the target protein are unavailable for many genes.

Another common method used to detect gene expression is to examine the presence of mRNA, the transcriptional product of a gene. Northern blot RNA

analysis or RT-PCR amplification can be used to detect the presence of the particular mRNA species. However, if the DNA sequence homology between the transgene and the host wild-type gene is very high, then neither Northern blotting nor RT-PCR would reliably differentiate expression of the host gene from the transgene. In this case, the probe used in the Northern blot analysis would hybridize with both the mRNA from the transgene and from the host wild-type alleles. Similarly, the primers used for RT-PCR would amplify both the transgene and the host wild-type alleles.

Heteroduplex analysis has been used to detect mismatches in dsDNA between a normal strand and one with an altered nucleotide sequence (5,7,11). This technique, which is based on DNA hybridization, has been used to detect gene mutations (4,8,12) and has been successfully applied for diagnostic purposes (1). Despite the sequence similarities

between the human transgene sequence and the host wild-type allele, the two gene sequences are not identical. Therefore, in theory, a set of primers used to amplify a cDNA fragment from both the transgene and the host wild-type gene can be used for heteroduplex amplification. If two DNA fragments with similar sequence are mixed, then heteroduplex products can be obtained by a denaturing and re-naturing process. When the products are run on a gel matrix, the homoduplexes (the original DNA fragments) migrate as a function of length, while the heteroduplex products (the transgene/ wild-type hybrid) migrate at a different rate through a gel (3,7,13) because the region of mismatch forms a "kink" in the DNA. Heteroduplex appears on the gel as one or two distinct bands that migrate separately from the homoduplex. Therefore, if both the transgene and the wild-type gene are expressed, then heteroduplex products that contain a one-stranded transgene and a one-stranded wild-type gene can be obtained by the denaturing and re-naturing process. Here we refer to this method as the RT-PCR heteroduplex method. Furthermore, using the transgene RNA and wild-type RNA as probes, this technique can also be used to detect and identify silenced transgenes without the need for any labeling process. Because the heteroduplex procedure is very simple to perform and can be done during PCR amplification, it is an efficient method to screen a large number of samples within a transgenic colony.

The *p53* gene, which is one of the most well-studied genes both in vivo and in vitro (2,10), has a very conserved binding domain. The protein and mRNA sequences for the domain are identical in length in mice and humans (Figure 1). We used a *p53* mutant (273H) transgenic mouse line to demonstrate the application of the RT-PCR heteroduplex technique in the detection of expression and silencing of a transgene and the detection of expression and silencing of the host wild-type homologous gene.

MATERIALS AND METHODS

RNA Isolation

A transgenic line of mice with lung-

specific expression of human mutant *p53* (273H) was established by microinjection of SP-C-*p53* (273H) fragments into a pronucleus of wild-type FVB/N mouse zygotes (unpublished data). Total RNA was isolated from the lungs of transgenic and wild-type mice. The lung tissue was wrapped with sterilized foil and frozen in liquid nitrogen. The frozen tissue was mechanically fragmented, and the RNA was isolated with Trizo[®] RNA isolation reagent (Invitrogen, Carlsbad, CA, USA), according to the manufacturer's protocol.

Mutant human *p53* RNA was prepared from a transfected human lung cancer cell line (H1299). Cells (H1299, deficient in *p53*) were transfected with human mutant *p53* (273H) constructs. Total RNA was isolated from the transfected cells with the Trizo isolation reagent.

RT-PCR Heteroduplex Analysis

Primers for the RT-PCR heteroduplex analysis were designed after examination of the mouse *p53* and the human mutant *p53* cDNA sequences (GenBank[®] accession nos. AF151353 and K03199, respectively). Primers E5F located in exon 5 (5'-GTCCGCGCC-ATGCCATCTA-3') and E7R located in exon 7 (5'-CATGTASTTGTAGTGATGGTGGT-3') were used to amplify identically sized 243-bp fragments from both human and mouse mRNA samples (Figure 1). Because the human and murine fragments are not identical in nucleotide sequence (87% identity), it is possible to form heteroduplex complexes between the transgene and the host wild-type gene. In addition, because primers amplify a 243-bp cDNA fragment between exons 5 and 7, DNA contamination would result in a larger product that includes introns 5 and 6.

The amplification was conducted in 25 μ L using the OneStep RT-PCR kit (Qiagen, Valencia, CA, USA). Each reaction contained 400 μ M each dNTP, 1 $\times$ RT-PCR buffer, 0.4 μ M each primer, 1 μ L OneStep RT-PCR enzyme mixture, and 200 ng total RNA. Conditions were at 50°C for 30 min and 95°C for 15 min, followed by 35 cycles of 94°C for 30 s, 64°C for 30 s, and 72°C for 1 min. A negative control, which included all reagents except the RNA sample,

Short Technical Reports

was used in each set of amplifications. The product of the RT-PCR was then diluted with same amount of 1× Tris-EDTA buffer and followed by a heteroduplex procedure. Heteroduplex conditions were at 94°C for 5 min, 70°C for 30 min, 60°C for 1 h, 50°C for 1 h, 45°C for 1 h, and 38°C for 5 h.

Ten microliters of RT-PCR heteroduplex product were loaded in 10% pre-cast NOVEX® polyacrylamide gels (Invitrogen) with 1× TBE running buffer. One lane was loaded with 4 µL 1-kb marker mixture. The marker mixture consists of a 2-µL 1 Kb DNA Ladder™ (Invitrogen), 6 µL water, and 2 µL loading buffer. Gels were run for approximately 1.5 h at 160 V and then stained in 1× TBE with 1 µg/mL EB for 10 min. The banding patterns were visualized by exposing the gel to UV light.

RESULTS

Using primers E5F and E7R, we amplified a 243-bp cDNA fragment from both transgenic and wild-type mice. The amplified products appeared as a single band in the 10% polyacrylamide gel (Figure 2, lanes 1 and 2). After the RT-PCR products were processed with denaturing and re-naturing steps and the RT-PCR heteroduplex products were separated via 10% PAGE, the products amplified from the transgenic samples showed three distinct bands, two originating from the heteroduplex products and one from the homoduplex products (Figure 3, lanes 4 and 6). In contrast, the product amplified from wild-type samples appeared as a single band (Figure 2, lanes 3 and 5). To confirm these results, we compared 10 additional samples each from transgenic mice and wild-type littermates. The heteroduplex products were formed in all transgenic samples but were absent in all non-transgenic samples. Sequencing of a heteroduplex band revealed one strand of transgene and one of murine *p53*.

Because gene silencing is a common mechanism of gene inactivation in transgenic animal models (9), we decided to evaluate this technique as a tool to study this phenomenon. We extracted total RNA from eight lung adenocarcinomas from the *p53* (273H) transgenic mice. One of the eight RT-

PCR heteroduplex products appeared as a single band using 10% PAGE. These results indicated that either the transgene or the host wild-type *p53* gene was silenced in that tumor. When we mixed the tumor RNA with an RNA sample extracted from a wild-type mouse, the RT-PCR heteroduplex product remained as a single band. However, when we mixed the tumor RNA with RNA extracted from a human lung cell line that contains the mutant *p53* (273H), heteroduplex bands were formed. This indicated that the transgene was silenced in the tumor sample, whereas murine wild-type *p53* was expressed (Figure 3). Furthermore, DNA sequencing of four clones that contain RT-PCR products amplified from the tumor sample showed that in each case the products were murine *p53*.

DISCUSSION

RT-PCR has been used widely to detect gene expression. In animal models, the functional domains of some genes are highly conserved across different species. It is technically challenging to design species-specific RT-PCR primers to detect the expression of a transgene. In this situation, the RT-PCR heteroduplex method is an adequate method to detect transgene and host wild-type gene expression. The major

advantages of the heteroduplex technique include the capacity to screen large number of samples in a short time and to differentiate a transgene and a host endogenous allele when they are very conserved in DNA sequence.

Although most host animals used in transgenic studies are pure inbred lines, heterozygous animals are often used for various purposes. In these cases, the heteroduplex products can be obtained by denaturing and re-naturing RT-PCR products amplified from a single diploid heterozygous individual. However, the position of a heteroduplex band formed between heterozygous alleles should differ from the position of the heteroduplex band formed between the transgenic and wild-type alleles in a gel. In this regard, a non-transgenic control is very important.

Heteroduplex products might be detectable using agarose gel electrophoresis. In our experience, the heteroduplex band was detected in 2% agarose gels in some transgenic samples (data not shown). However, agarose gels could not detect all heteroduplex bands efficiently. For this reason, we recommend using 10%–20% PAGE if the RT-PCR heteroduplex method is to be used to detect gene expression.

Preferably, the animals used should be as young as possible, since the risk of spontaneous mutations increases when animals get older. In addition, somatic

Figure 1. Alignment of the human *p53* mRNA fragment and the murine *p53* mRNA fragment. Primers E5F located in exon 5 (5'-GTCCGCGCCATGGCCATCTA-3') and E7R located in exon 7 (5'-CATGTASTTGTAGTGGATGGTGGT-3') were designed based on human and murine *p53* mRNA sequences to amplify identically sized 243-bp fragments from both human and mouse mRNA samples. Nucleotide sequence identity is 87% between the human fragment and the murine fragment. HUM, human *p53* sequence. MUR, murine *p53* sequence.

Short Technical Reports

mutation is a common event in malignant cells, and heteroduplex bands may form between the somatic mutant and wild-type allele or between the somatic mutant and the transgenic allele. However, the position of a heteroduplex band formed between the transgene and the wild-type gene should differ from the position of a heteroduplex band formed between the somatic mutant and wild-type gene. Therefore, we recommend using a transgenic positive control sample in the RT-PCR heteroduplex amplification.

It is of interest to note that the RT-PCR heteroduplex method can be used to detect if a gene has been silenced and to identify which gene has been silenced, even if the transgene and the host wild-type homologous gene are

very conserved in their DNA sequence. The probes (or drivers) used in the test are the host wild-type RNA samples and the transgene RNA samples, which are easily obtained. Since the heteroduplex method has been used to detect point mutations (13), it is possible that the RT-PCR heteroduplex method can be used to detect the expression of a transgene, even if there is only a single base difference between a transgene and the host wild-type alleles. If so, then a high-quality polyacrylamide gel or MDE gel (BMA, Rockland, ME, USA) should be utilized.

In conclusion, the RT-PCR heteroduplex technique is a suitable method to differentiate the expression of a transgene and the wild-type gene in transgenic animals, if the DNA se-

quence between the transgene and host wild-type gene is too conserved to design gene- and species-specific RT-PCR primers. The low cost, speed, simplicity, and the ability to screen a large number of samples are major advantages of this method.

REFERENCES

1. Cherny, S., S. Mraz, L. Su, J. Harvell, and S. Kohler. 2001. Heteroduplex analysis of T-cell receptor λ gene rearrangement as an adjuvant diagnostic tool in skin biopsies for erythroderma. *J. Cutan. Pathol.* 28:351-355.
2. Chumakov, P.M. 2000. Function of the *p53* gene: choice between life and death. *Biochemistry (Mosc.)* 65:28-40.
3. Delwart, E.L., H.W. Sheppard, B. Walker, J. Goudsmit, and J.I. Mullins. 1994. Human immunodeficiency virus type 1 evolution in vivo tracked by DNA heteroduplex mobility assays. *J. Virol.* 68:6672-6683.
4. Delwart, E.L., E.G. Shpaer, J. Louwagie, F.E. McCutchan, M. Grez, H. Rubsamen-Waigmann, and J.I. Mullins. 1993. Genetic relationships determined by a DNA heteroduplex mobility assay: analysis of HIV-1 *env* genes. *Science* 262:1257-1261.
5. Freel, S.A., J.M. Williams, J.A. Nelson, L.L. Patton, S.A. Fiscus, R. Swanstrom, and D.C. Shugars. 2001. Characterization of human immunodeficiency virus type 1 in saliva and blood plasma by V3-specific heteroduplex tracking assay and genotype analyses. *J. Virol.* 75:4936-4940.
6. Jackson, I.J. and C.M. Abbott. 2000. *Mouse Genetics and Transgenics*. Oxford University Press, Oxford.
7. Keen, J., D. Lester, C. Inglehearn, A. Curtis, and S. Bhattacharya. 1991. Rapid detection of single base mismatches as heteroduplexes on Hydrolink gels. *Trends Genet.* 7:5.
8. Larsen, L.A., P.S. Andersen, J.K. Kanters, J.R. Jacobsen, J. Vuust, and M.A. Christiansen. 1999. Single strand conformation polymorphism/heteroduplex (SSCP/HD) method for detection of mutations in 15 exons of the KVLQT1 gene, associated with long QT syndrome. *Clin. Chim. Acta* 280:113-125.
9. Lung, H.Y., I.S. Meeus, R.S. Weinberg, and G.F. Atweh. 2000. In vivo silencing of the human λ -globin gene in murine erythroid cells following retroviral transduction. *Blood Cells Mol. Dis.* 26:613-619.
10. May, P. and E. May. 1999. Twenty years of *p53* research: structural and functional aspects of the *p53* protein. *Oncogene* 18:7621-7636.
11. Resch, W., N. Parkin, E.L. Stuelke, T. Watkins, and R. Swanstrom. 2001. A multiple-site-specific heteroduplex tracking assay as a tool for the study of viral population dynamics. *Proc. Natl. Acad. Sci. USA* 98:176-181.
12. Shih, H.A., K.L. Nathanson, S. Seal, N. Collins, M.R. Stratton, T.R. Rebbeck, and

Figure 2. The RT-PCR heteroduplex method was used to analyze the expression of the transgene (human *p53* 273H) and host wild-type homologous gene (the murine wild-type *p53*) in *p53* transgenic and wild-type mice. A 243-bp fragment was amplified. The RT-PCR products amplified from transgenic and non-transgenic samples without heteroduplex processing appeared as a single band in a 10% polyacrylamide gel (lanes 1 and 2). After the RT-PCR products were processed with denaturing and re-naturing treatment, the products amplified from transgenic samples contained three distinct bands, two heteroduplex and one homoduplex (lanes 4 and 6), whereas the non-transgenic mice have the homoduplex product only, which appears as a single band (lanes 3 and 5). Lane M, 1-kb marker.

Figure 3. RT-PCR heteroduplex technique was used to detect transgene (human *p53* 273H) and/or murine wild-type *p53* gene silencing. Lane 1 is a RT-PCR heteroduplex product amplified from a wild-type mouse, and lane 2 is a RT-PCR heteroduplex product amplified from a transfected human lung cancer cell line that contains the human mutant *p53* (273H). Lane 3 is a RT-PCR heteroduplex product amplified from a lung adenocarcinoma collected from a transgenic mouse in which gene silencing was suspected. Lane 4 is the RT-PCR heteroduplex product amplified from a mixture sample [the RNA mixture contains RNA extracted from the human cancer cell line containing the human mutant gene (lane 2) and the RNA extracted from the mouse lung tumor (lane 3)]. Lane 5 is a RT-PCR heteroduplex product amplified from another mixture of samples [the RNA mixture contains RNA extracted from a wild-type mouse (lane 1) and RNA extracted from a mouse lung tumor (lane 3)]. Lane 6 is the RT-PCR heteroduplex product amplified from a transgenic sample used as a positive control. The matrix was a 10% polyacrylamide gel.

Short Technical Reports

B.L. Weber. 2000. BRCA1 and BRCA2 mutations in breast cancer families with multiple primary cancers. *Clin. Cancer Res.* 6:4259-4264.

13. **White, M.B., M. Carvalho, D. Derse, S.J. O'Brien, and M. Dean.** 1992. Detecting single base substitutions as heteroduplex polymorphisms. *Genomics* 12:301-306.

The work was supported by National Cancer Institute grant no. K01 CA76970 to M.A.V.-C. and no. P30 CA16058 to The Ohio State University Comprehensive Cancer Center. Address correspondence to Dr. Miguel A. Villalona-Calero, Arthur G. James Cancer Hospital, Ohio State University, B406 Starling-Loving Hall, 320 W10th Avenue, Columbus, OH 43210-1240, USA. e-mail: villalona-1@medctr.osu.edu

Received 19 October 2001; accepted 19 February 2002.

W. Duan, H. Ding, W.-G. Zhu, K. Srinivasan, G.A. Otterson, and M.A. Villalona-Calero

*The Ohio State University
College of Medicine and
Public Health
Columbus, OH, USA*

cAMP Binding Protein Assay for Widespread Use in Cell Signaling Studies

BioTechniques 33:66-72 (July 2002)

ABSTRACT

cAMP plays a critical role in intracellular signaling pathways that regulate proliferation or differentiation. The cAMP binding protein assay, using a naturally derived cAMP binding protein, is one of the most widely used methods for cAMP determination. The major steps of this binding assay include purification of the binding protein, cAMP extraction from samples, and quantification of the cAMP. Most purification methods of the cAMP binding protein were published before 1975, and many of the materials and methods are outdated. Here

we describe an updated method of purification of cAMP binding protein from bovine skeletal muscle with the advantages of simplicity, low cost, and high yield. The isolation procedures can be completed in two days using commercially available materials and equipment. The cAMP binding properties of the isolated protein can be utilized for more than two years. Binding protein isolated from 1 kg bovine muscle is sufficient for at least 3×10^4 assay tubes. Furthermore, we describe the techniques of cAMP extraction and quantification that have been used successfully in studying parathyroid hormone signaling as an example of a G protein-linked seven transmembrane domain receptor that signals through the protein kinase A pathway.

INTRODUCTION

cAMP was discovered and recognized as a second messenger in intracellular signaling pathways more than 40 years ago (15). Since that time, cAMP-dependent signaling has been under intense scrutiny in physiological systems such as the nervous (10,16) and skeletal systems (1,2), and in pathologic conditions such as carcinoma (3,9). In the last year alone, more than 5000 cAMP-related publications were listed in Medline.

Many hormones exert their specific functions through binding to seven transmembrane domain G-protein coupled receptors. The cAMP pathway is downstream of receptors that couple to Gs, one of the heterotrimeric GTP-binding proteins. Activation of Gs releases G α_s , which activates adenylate cyclase and subsequently leads to cAMP generation. The increase in cAMP activates cAMP-dependent protein kinase A (PKA), which subsequently phosphorylates specific substrates to generate cellular responses. Thus, the determination of cAMP levels plays a critical role in evaluating PKA-mediated signal transduction. The most widely employed techniques to quantify cAMP levels are based on the principle of competition binding, such as radioimmunoassay (RIA) (4,5), enzymatic immunoassay (EIA) (8), and binding protein assay (13). In immunoassays, a cAMP-specific antibody is used for competitive binding between radiolabeled cAMP (RIA) or enzyme-conjugated cAMP (EIA) and

the cAMP in samples or standards. In contrast, the binding protein assay involves the competitive binding between cAMP in samples or standards and [3 H]cAMP to a naturally derived cAMP binding protein. Disadvantages of RIA or EIA are that the antisera used for immunoassays are difficult to prepare. Further, although commercial immunoassay kits are available with high sensitivity, the unit price/sample is high; hence, it becomes quite expensive for repetitive or large-scale experiments. Finally, the shelf life of commercial assays that utilize iodinated cAMP is short and can limit the design of experiments to optimize the activity of the iodinated tracer. For routine determination of cAMP levels, the binding protein assay is likely a better choice. The binding protein used is easy to prepare with high yield and is stable for months to years. In addition, the binding protein assay possesses the advantages of simplicity, low cost, and high specificity. The binding protein assay has been successfully utilized for determination of cAMP levels in various tissue samples and cell cultures.

The major steps of the cAMP binding protein assay include purification of the binding protein, cAMP extraction from samples, and quantification of cAMP. The major obstacle of using a straightforward cAMP binding protein assay is the availability of cAMP binding protein. Most cAMP binding protein purification protocols were published more than 25 years ago (7,14,17). They were either complicated or used outdated materials and equipment, which ren-

Figure 1. DEAE-cellulose chromatograph. Potassium phosphate buffer (as described in Materials and Methods) was added at time point A to elute the cAMP binding protein. cAMP binding protein was collected between time points B and C. Time points B and C were 1 and 2.2 h after time point A, respectively. Flow rate was 135 mL/h.