surging receptor mediated changes in intracellular calcium levels. *Anal. Biochem.* 286:231-237.

6. Haas, J., E.C. Park, and B. Seed. 1996. Codon usage limitation in the expression of HIV-1 envelope glycoprotein. *Curr. Biol.* 6:315-324.
7. Kolodziej, P.A. and R.A. Young. 1991. Epitope tagging and protein surveillance. *Methods Enzymol.* 194:508-519.
8. Pear, W.S., G.P. Nolan, M.L. Scott, and D. Baltimore. 1993. Production of high-titer helper-free retroviruses by transient transfection. *Proc. Natl. Acad. Sci. USA* 90:8392-8396.
9. Shaw, B.D., O. Kozlova, N.D. Read, B.G. Turgeon, and H.C. Hoch. 2001. Expression of recombinant aequorin as an intracellular calcium reporter in the phytopathogenic fungus *Phyllosticta ampellicida*. *Fungal Genet. Biol.* 34:207-215.
10. Yang, T.T., L. Cheng, and S.R. Kain. 1996. Optimized codon usage and chromophore mutations provide enhanced sensitivity with the green fluorescent protein. *Nucleic Acids Res.* 24:4592-4593.
11. Yao, F. and E. Eriksson. 1999. A novel tetracycline-inducible viral replication switch. *Hum. Gene Ther.* 10:419-427.
12. Yao, F., T. Svensjo, T. Winkler, M. Lu, C. Eriksson, and E. Eriksson. 1998. Tetracycline repressor, tetR, rather than the tetR-mammalian cell transcription factor fusion derivatives, regulates inducible gene expression in mammalian cells. *Hum. Gene Ther.* 9:1939-1950.

Address correspondence to Dr. John A. Printen, AstraZeneca Pharmaceuticals, Enabling Science and Technology, Mereside Alderley Park, Macclesfield, Cheshire SK10 4TG, UK. e-mail: john.printen@astrazeneca.com

Received 4 April 2002; accepted 29 May 2002.

Wendy I. Vernon and John A. Printen
AstraZeneca Pharmaceuticals
Cheshire, UK

For reprints of this or any other article, contact
Reprints@BioTechniques.com

Accelerated Apoptotic DNA Laddering Protocol

BioTechniques 33:734-736 (October 2002)

Apoptosis is a natural "self-destruct" process that enables multicellular organisms to eliminate unwanted cells during development and homeostasis. Interference with apoptosis, such as by mutation of the tumor suppressor gene *p53*, may contribute to carcinogenesis. Apoptosis, also known as programmed cell death, is a genetically determined series of events involving de novo protein synthesis and resulting in the death of apoptosing cells. The orderly realignment of biochemical events in an apoptosing cell offers many targets for the detection and monitoring of the apoptotic process: membrane blebbing, phosphatidylserine externalization, mitochondria cytochrome C release, activation of caspases, poly(ADP-ribose) polymerase cleavage, compromise of membrane cleavage, chromatin condensation, DNA fragmentation, and the formation of apoptotic bodies (1,2,4).

Among the many markers of apoptosis, DNA fragmentation is considered a hallmark in many, if not all, apoptotic pathways (3,5). During apoptosis, nucleases are activated, cleaving internucleosomal DNA strands unprotected by histones to generate multiples of 198-bp DNA fragments that will run electrophoretically as a characteristic DNA ladder in agarose gels. Thus, a demonstrable presence of DNA laddering in a given cell population is indicative of ongoing apoptosis and has generally been regarded as a distinguishing feature of cell death by apoptosis rather than necrosis.

Isolation of apoptotic DNA fragments is commonly accomplished by lysing apoptotic cells with a cell-lysing buffer, followed by the digestion of contaminating RNA with a high concentration of RNase A in the presence of SDS, elimination of contaminating proteins by protease digestion, and finally agarose gel electrophoresis (1). While being straightforward, this procedure and variations of it suffer from long incubation times (4 h or more) and contaminating RNA fragments that cannot be totally digested by RNase A.

RNase A cleaves the 3' of single-stranded C and U residues and will not result in the complete digestion of RNA to mononucleotides. To circumvent these problems, here we describe an efficient and inexpensive apoptotic DNA laddering protocol that can be completed in less than 1 h.

As an illustration, 10^6 U937 cells (up to 5×10^6 cells can be used without scaling up the apoptotic DNA extraction protocol) growing in RPMI 1640 with 5% FCS were treated with 2 $\mu\text{g}/\text{mL}$ camptothecin for 3 h. Cells were recovered by centrifugation, and the cell pellets were lysed by gentle, repeated pipetting in a 2-mL microcentrifuge tube with 0.5 mL cell lysing buffer (50 mM Tris-HCl, pH 7.5, 1% Nonidet™ P-40, 20 mM EDTA). After centrifugation at $1600 \times g$ for 5 min to remove cell debris as well as intact nuclei of non-apoptosing cells (1), the supernatant was subject to one round of phenol:chloroform:isoamyl alcohol (25:24:1; pH 7.4; 0.5 mL) extraction. Apoptotic DNA fragments and contaminating RNA present in the liquid phase

Figure 1. Isolation of apoptotic DNA fragments. Apoptotic DNA fragments were isolated from camptothecin-treated U937 cells as described in the text and were run on a 1.8% TAE agarose gel. Lane 1, Hi-Lo DNA markers (Bionexus, Berkeley, CA, USA); lane 2, control U937 not treated with camptothecin (10^6 cells); lane 3, camptothecin-treated U937 (10^6 cells); lane 4, camptothecin-treated U937 (1.5×10^6 cells); and lane 5, camptothecin-treated U937 (10^6 cells). Lanes 2-4 were obtained from the RNaseOne digestion protocol described in this paper; lane 5 was from the RNase A/proteinase K double digestion method (1).

Benchmarks

were precipitated by adding 50 μ L 3 M sodium acetate, pH 5.2, 1 μ L nuclease-free glycogen (Roche Applied Science, Indianapolis, IN, USA) and 0.6 mL isopropanol. After incubation on ice for 5 min, the precipitated nucleic acids were pelleted by centrifugation at 12 900 $\times$ g for 5–10 min at 40°C and followed by a brief 70% ethanol wash. The pellet was resuspended in 30 μ L 1 $\times$ RNaseOne® buffer (Promega, Madison, WI, USA). One microliter (5–10 U) RNaseOne was added to the sample, followed by incubation at 37°C for 30 min or longer. Apoptotic DNA fragments were resolved by 1.8% TAE agarose gel electrophoresis. For comparison, another identically camptothecin-treated U937 sample was extracted with the conventional RNase A-proteinase K double digestion protocol for isolating apoptotic DNA fragments (1).

Figure 1, lane 3, shows a typical apoptotic DNA ladder achievable with our protocol. It compares favorably with the traditional RNase A/proteinase K DNA laddering protocol (Figure 1, lane 5), where incompletely digested RNA could sometimes be seen accumulating at the bottom of the DNA ladder. In contrast, the protocol described here rendered apoptotic DNA ladders apparently free of contaminating undigested RNA (Figure 1, lane 3) and exhibited similar, if not more, sensitivity

as compared to the RNase A/proteinase K double digest method (Figure 1, lane 5). Further, smearing due to contaminating genomic DNA was insignificant (Figure 1, lane 2). With our protocol, the extent of DNA fragmentation can be compared among different samples if the samples are treated and extracted identically: there was more DNA laddering in lane 4 (1.5×10^6 cells) than in lane 3 (1×10^6 cells) in Figure 1. Similarly, when a set of identical cultures of U937 cells (1×10^6 cells each) treated with 2 μ g/mL camptothecin for 3 h was followed with time, our protocol revealed quantitatively the surge in DNA fragmentation 3 h after camptothecin treatment (Figure 2, lane 5).

The protocol documented here represents a fast method to isolate efficiently apoptotic DNA fragments from apoptosing tissue culture cells. From lysing cell pellets to the end of RNaseOne digestion, the whole procedure takes less than 1 h. The merits of our protocol over previous protocols include increased speed and efficiency, a more complete reduction of genomic DNA contamination, and the elimination of using proteinase K. In our protocol, the use of RNaseOne, which cleaves the phosphodiester bond between any two ribonucleotides, ensures a more com-

plete digestion of RNA to monomers, and hence avoidance of RNA contamination that might mask apoptotic DNA fragments during gel electrophoresis. As with conventional apoptotic DNA fragment isolation methods, our accelerated protocol still allows effective comparison of the extent of apoptosis (DNA fragmentation) among different samples. Finally, we have successfully used our protocol for other cell lines including Jurkat, HeLa, L-cells, MCF-10A, and MCF-7 (data not shown).

REFERENCES

1. Herrmann, M., H.M. Lorenz, R. Voll, M. Grunke, W. Woith, and J.R. Kalden. 1994. A rapid and simple method for the isolation of apoptotic DNA fragments. *Nucleic Acids Res.* 22:5506-5507.
2. Kaufmann, S.H. and M.O. Hengartner. 2001. Programmed cell death: alive and well in the new millennium. *Trends Cell Biol.* 11:526-534.
3. Nagata, S. 2000. Apoptotic DNA fragmentation. *Exp. Cell Res.* 256:12-18.
4. Olasode, B.J. 1997. Dying by default, the biology of apoptosis: a review. *East Afr. Med. J.* 74:108-111.
5. Zhang, J.H. and M. Xu. 2000. DNA fragmentation in apoptosis. *Cell Res.* 10:205-211.

Address correspondence to Dr. Michael C. Yeung, Snyder Medical Research Institute, 1407 Wheat Road, Winfield, KS 67156, USA. e-mail: myeung@snydermri.org

Received 5 April 2002; accepted 30 May 2002.

Michael C. Yeung
*Snyder Medical Research
Institute
Winfield, KS, USA*

**For reprints of this or
any other article, contact
Reprints@BioTechniques.com**

Figure 2. Detection of apoptotic DNA fragments over time. Four identical cultures of U937 cells (1×10^6 cells each) were treated with 2 μ g/mL camptothecin for up to 3 h and extracted for apoptotic DNA fragments. Lane 1, Hi-Lo DNA markers; lane 2, untreated control; lanes 3–5, cells treated with camptothecin for 1, 1.5, and 3 h, respectively.