Short Technical Reports

only and *ADRB2* SNP 2 by PolyPhred only. These findings emphasize the importance of sequencing in both directions and that an alternative method of SNP confirmation may be required.

Initially, SNPs were reported for all sets of sequences analyzed, but many of these were false-positive SNPs located in stretches of consecutive Ts. A filter was included in the SNPCEQer program to flag these SNPs as suspect, except where the SNP was reported at the first T. No genuine SNPs were reported in the regions sequenced from *REV1L*, *CREB1*, *TRF4*, and *MAP4K3*.

SNPCEQer offers a convenient method of compiling a list of potential polymorphic positions for further examination. SNPCEQer provides a functionality not currently available as part of the CEQ2000 software analysis package; since it can run in a Windows NT environment, it offers convenience and familiarity to the typical user.

REFERENCES

1. Beckman Coulter. 2000. CEQ™2000 DNA Analysis System User's Guide. Beckman Coulter, Fullerton, CA.
2. Collins, F.S., M.S. Guyer, and A. Chakravarti. 1997. Variations on a theme: cataloging human DNA sequence variation. *Science* 278:1580-1581.
3. Dobbs, M. and N. Gee. 2001. Detecting Heterozygotes with the CEQ™2000XL. Beckman Coulter Application Information A-1892A. Beckman Coulter, Fullerton, CA.
4. Ewing, B. and P. Green. 1998. Base-calling of automated sequencer traces using Phred. II. Error probabilities. *Genome Res.* 8:186-194.
5. Ewing, B., L. Hillier, M.C. Wendt, and P. Green. 1998. Base-calling of automated sequencer traces using Phred. I. Accuracy assessment. *Genome Res.* 8:175-185.
6. Gordon, D., C. Abajian, and P. Green. 1998. Consed: a graphical tool for sequence finishing. *Genome Res.* 8:195-202.
7. Hall, T. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids. Symp. Ser.* 41:95-98.
8. Marth, G., L. Korf, M.D. Yandell, R.T. Yeh, Z. Gu, H. Zakeri, N.O. Stitzel, L. Hillier, et al. 1999. A general approach to single-nucleotide polymorphism discovery. *Nat. Genet.* 23:452-456.
9. Nickerson, D.A., V.O. Tobe, and S.L. Taylor. 1997. PolyPhred: automating the detection and genotyping of single nucleotide substitutions using fluorescence-based resequencing. *Nucleic Acids Res.* 25:2745-2751.
10. Notredame, C., D.G. Higgins, and J. Heringa. 2000. T-Coffee: a novel method for fast and accurate multiple sequence alignment. *J. Mol. Biol.* 302:205-217.
11. Picoult-Newberg, L., T.E. Ideker, M.G. Pohl, S.L. Taylor, M.A. Donaldson, D.A. Nickerson, and M. Boyce-Jacino. 1999. Mining SNPs from EST databases. *Genome Res.* 9:167-174.
12. Rieder, M.J., S.L. Taylor, V.O. Tobe, and D.A. Nickerson. 1998. Automating the identification of DNA variations using quality-based fluorescence re-sequencing: analysis of the human mitochondrial genome. *Nucleic Acids Res.* 26:967-973.
13. Taillon-Miller, P., Z. Gu, Q. Li, L. Hillier, and P.Y. Kwok. 1998. Overlapping genomic sequences: a treasure trove of single-nucleotide polymorphisms. *Genome Res.* 8:748-754.
14. Walther, D., G. Bartha, and M. Morris. 2001. Basecalling with LifeTrace. *Genome Res.* 11:875-888.

This work was supported by the National Institutes of Health (Programs for Genomic Applications grant no. 1U01-HL66880 and grant no. T32-HL07360), the State of Texas Advanced Technology Program (grant no. 010019-0212), the National Cancer Institute (Specialized Programs of Research Excellence grant no. P50 CA70907), and via a collaboration with Beckman Instruments. We wish to thank Andy Vuong for assisting in the development of the computer code. We would also like to thank Profs. R.V. Shohet, G.E. O'Keefe, and J. D. Minna for genomic DNA samples. Address correspondence to Dr. Elizabeth Flood, Eugene McDermott Center for Human Growth and Development, The University of Texas Southwestern Medical Center, Dallas, TX 75390-8591, USA. e-mail: elizabeth.flood@utsouthwestern.edu

Received 4 December 2001; accepted 20 June 2002.

E.M. Flood, F. Tang, M.M. Horvath, A. Pertsemlidis, and H.R. Garner
The University of Texas Southwestern Medical Center Dallas, TX, USA

For reprints of this or any other article, contact
Reprints@BioTechniques.com

SNPkit: An Efficient Approach to Systematic Evaluation of Candidate Single Nucleotide Polymorphisms in Public Databases

BioTechniques 33:822-830 (October 2002)

ABSTRACT

There is widespread interest in devising genotyping methods for SNPs that are robust, inexpensive, and simple to perform. Although several high-throughput SNP genotyping technologies have been developed, including the oligonucleotide ligation assay, real-time PCR, and mass spectrometry, the issues of simplicity and cost-effectiveness have not been adequately addressed. Here we describe the application of a novel computer software package, SNPkit, which designs SNP genotyping assays based on a classical approach for discriminating alleles, restriction enzyme digestion. SNPkit can be used in genotyping assays for almost any SNPs including those that do not alter "natural" restriction sites. Using this method, 164 SNPs have been evaluated in DNA samples from 48 immortalized cell lines of randomly selected Chinese subjects. Sixty-two (37.8%) of the SNPs appeared to be common (frequencies of the minor alleles are $\geq 5\%$) and were subsequently applied to a larger population-based sample. Overall, by using SNPkit, we have been able to validate and genotype accurately a large fraction of publicly available SNPs without sophisticated instrumentation.

INTRODUCTION

SNPs are the most abundant type of DNA variation, with an estimated frequency of 1 in every 300–500 bp (7), and they provide important physical landmarks for identifying genes important in human disease and drug responses (13). Because of rigorous efforts in SNP discovery, more than 2 million SNPs have been identified and cataloged (19).

Several publicly available SNP data-

Short Technical Reports

bases have been created, including the National Center for Biotechnology Information (NCBI) dbSNP (<http://www.ncbi.nlm.nih.gov/SNP/>), Human Gene Mutation Database (HGMD®), Human Genic Bi Allelic Sequences (HGBASE) (<http://hgbase.cgr.ki.se/>), ENSEMBL (<http://www.ensembl.org/>), and the SNP consortium database (<http://snp.cshl.org/>), which provide the most comprehensive resource for human variation studies. This ever-expanding data set is a rich source of gene-associated and anonymous SNPs (5).

However, many of the SNPs deposited in the public domain are identified using *in silico* screening through comparing sequences from overlapping genomic clones (20). SNPs identified in this way are only “candidate” SNPs, and typically only 50%–63% are validated by experimentation (13). Furthermore, some SNPs display ethnic specificity, and their frequencies in one population could not be directly applied to another ethnic group (10,26). Thus, the emerging era of SNP-based genetic analysis has underscored the need for simple and accurate genotyping methods that can accommodate thousands of loci with economies of cost and consumption of sample DNA.

RFLP is a classical enzymatic approach that is applied in a standard laboratory for genotyping SNPs with natural alternations of restriction recognition sites (25), with low false-negative and false-positive rates. The method relies on the exquisite selectivity of restriction endonucleases for short stretches of DNA sequence that act as recognition sites for DNA strand cleavage. Because of the inherent catalytic specificity of enzymes for their substrates, RFLP generally possesses a high degree of assay fidelity. The cost of an assay is generally low, especially when we depend primarily on the common robust restriction enzymes, which is possible in most cases of mutagenic PCR-RFLP.

If a SNP fails to alter a known restriction site, then one can artificially generate a mutagenic PCR primer. This is a well-known method (9,11,12,21), but virtually all these assays are designed in a nonstandardized fashion. Here we report a novel SNP approach, SNPkit, which combines the classical PCR-RFLP with a newer *in silico*

methodology that engineers an artificial restriction site through the design of mutagenic primers in an automated mode. Furthermore, we explore the extent of linkage disequilibrium in various genomic segments examined.

MATERIALS AND METHODS

Software Algorithm of Oligonucleotide Design

To ensure the success of PCR, the algorithm keeps at least two bases immediately upstream or downstream from the SNP in perfect match with the DNA template. The site of the SNP, along with its unaltered 5′ and 3′ flanking sequences (0-mutagenic model), is searched exhaustively against a comprehensive restriction endonuclease database to see if the SNP creates or eliminates a restriction site. If such an 0-mutagenic model fails, then one (1-mutagenic model), two (2-mutagenic model), or three (3-mutagenic model) nucleotides within the 5–10 bp flanking region spanned by the primer sequence are randomly permuted, such that the targeted SNP in conjunction with these flanking artificially engineered mutagenics are automatically and thoroughly searched to see if they result in any change of an endonuclease recognition site. The software package (written by Dr. Tianhua Niu) will typically produce several choices under these different mutagenic models that often correspond to different restriction enzymes. Then, the best mutagenic primer is selected based on (i) the number of mismatches introduced and (ii) the price and robustness of the restriction enzyme. The opposing primer is designed with the Primer3 software (http://www-genome.wi.mit.edu/cgi-bin/primer/primer3_www.cgi). Examples are provided in Table 1.

Sample DNA

Forty-eight lymphoblastoid cell lines were established from individuals randomly ascertained in Anqing, China, where our linkage and association study samples were collected. Genomic DNA was isolated from harvested cell using standard protocols (23).

PCR and Restriction Endonuclease Digestion

A uniform touchdown PCR protocol was implemented for all PCR assays. Each reaction was carried out in a standard, 10 µL cocktail solution containing 30 ng genomic DNA, 2.5 mM MgCl₂, 200 µM each dNTP, 200 nM each primer, 0.5 U *Taq* DNA polymerase (Applied Biosystems, Foster City, CA, USA), and 1× reaction buffer containing 10 mM Tris-HCl, pH 8.3, 50 mM KCl. The PCR profile was as follows: an initial denaturation at 94°C for 10 min, followed by 14 touchdown cycles of 94°C for 30 s, 61°C–54°C for 30 s (a steady decrease of 0.5°C each cycle), and 72°C for 30 s, followed by 30 amplification cycles of 94°C for 30 s, 54°C for 30 s, 72°C for 30 s, and a final 7-min extension at 72°C. The PCR product was digested with corresponding restriction endonuclease, according to the manufacturer’s instructions (New England Biolabs, Beverly, MA, USA).

Agarose Gel Electrophoresis

Digestion products were electrophoresed on 2.5% SeaKem® LE (BioWhittaker Molecular Applications, Rockland, ME, USA) agarose gels at 300 V for 60 min and visualized with ethidium-bromide staining and UV illumination. The gel image was saved electronically using the Gel-Doc™ 2000 Gel Documentation System (BioRad Laboratories, Hercules, CA, USA).

Analysis

Allele frequencies were calculated by gene counting methods, and linkage disequilibrium was calculated by the LDMAX software (<http://www.sph.umich.edu/csg/abecasis/GOLD/docs/ldmax.html>).

RESULTS

A total of 189 SNPs downloaded from publicly available databases (i.e., NCBI dbSNP) were subject to validation using the SNPkit. The SNPkit software package successfully designed mutagenic PCR-RFLP assays for all these markers. Overall, 25 of the 189

Table 1. Mutiple Mutagenic PCR-RFLP Assays Designed on the Same SNPs

SNP	Chromosome	First RFLP Enzyme	Second RFLP Enzyme
rs3899404	19	<i>Hpy188I</i>	<i>BsoBI</i>
rs435499	19	<i>Fnu4HI</i>	<i>BbvI</i>
rs344594	19	<i>MwoI</i>	<i>Fnu4HI</i>
rs344593	19	<i>StyI</i>	<i>MnII</i>
rs453462	19	<i>BanI</i>	<i>KpnI</i>
rs1915390	19	<i>BstI</i>	<i>BsrBI</i>
rs1915377	19	<i>BstUI</i>	<i>MluI</i>
ACEintron9	17	<i>AluI</i>	<i>MseI</i>

SNPs failed the PCR step. Among the successfully amplified 164 SNPs, 94 (57.3%) were monomorphic in our sample. In the remaining 70 (42.7%) SNPs, 8 (11.4%) were rare SNPs (minor allele frequency < 5%) and 62 (88.6%) were common (minor allele frequency ≥ 5%). These SNPs were located in more than 10 genes from human chromosomes 2, 8, 15, 19, and 22. Among these 70 SNPs, 25 (35.7%) and 26 (37.1%) were G→A and C→T transitions, respectively; 6 (8.6%) and 4 (5.7%) were A→C and G→T transversions, respectively; and 6 (8.6%) and 3 (4.3%) were G→C and A→T transversions, respectively. The diagnostic fragments for each SNP were unambiguously separated by the agarose gel electrophoresis. Since several RFLPs used the same restriction enzyme, we tested a multiplex PCR-RFLP design. Duplex RFLP can provide clear fragment separation patterns for both SNPs.

Figure 1 shows the distribution of

the rare allele frequencies for these 70 SNPs, 41 (58.6%) of which are located in CpG or GpC dinucleotides. We found that almost every pair of SNPs within each gene is in strong linkage disequilibrium, with the disequilibrium from 0.5 to 0.8. SNPs in different genes on the same chromosome show only weak linkage disequilibrium.

DISCUSSION

Because of the surge in interest in using SNPs for genotyping, a facile and affordable method for this is an absolute necessity. Thanks to the wealth of sequence information in public databases, candidate SNPs can be identified by in silico methods. Previous studies showed that 55% of candidate SNPs found by the computer program SNPfinder (<http://lpgws.nci.nih.gov:82/perl/snp/snp.cgi.pl>) can be validated using experimental procedures (8). Here we pre-

Figure 1. Frequency of the minor allele of 70 SNPs validated in a Chinese population.

Short Technical Reports

Table 2. Comparison of Major Genotyping Technologies for SNP Evaluation or Genotyping

Genotyping Technology	Estimated Cost of SNP Genotyping for Moderate Sample Size (per SNP) ^a	Low Initial Investment on Instrumentation ^b	Assays Robustness ^c	Summary ^d
Mutagenic PCR-RFLP	\$0.30–\$0.50	Yes	Yes	Robust and cost-effective
Real-Time PCR	\$0.50–\$1.00	No	Yes	Robust
Oligonucleotide Ligation Assay	\$0.50–\$1.00	No	Yes	Robust
Allele-Specific Amplification	\$0.30–\$0.50	Yes	No	Cost-effective
Single Base Extension-Fluorescent Polarization	\$0.30–\$0.50	No	No	Difficult
Mass Spectrometry	\$0.30–\$0.50	No	Yes	Robust

These are estimated costs for a typical molecular biology laboratory. The costs are estimated without consideration of multiplexing, which can achieve a lower cost for mutagenic PCR-RFLP and other listed technologies.

^aThe oligonucleotide ligation and real-time PCR assays need specific fluorescence-tagged oligonucleotides. Single base extension-fluorescent polarization involves a dye-labeled dideoxy terminator that is incorporated allele-specifically in the presence of a matching ssDNA template.

^bRFLP and allele-specific amplification could be performed using inexpensive agarose gel electrophoresis. However, the rest of the assays all need sophisticated instrumentation for allele detection. For example, fluorescence-based oligonucleotide ligation requires an ABI PRISM® 377 Semi-Automated DNA Sequencer (Applied Biosystems); single base extension-fluorescent polarization requires solution-phase fluorescence polarization detection systems; and the mass spectrometry method requires the SpectroJET spotting device and the time-of-flight mass spectrometer (both from Sequenom).

^cAccording to our and other researchers' experiences, allele-specific amplification and single base extension-fluorescent polarization are particularly sensitive to primer designs and assay conditions (1). As a consequence, time-consuming protocol optimization (e.g., adjustment of Mg²⁺ concentration, annealing temperature, etc.) is frequently needed, which constitutes a significant obstacle for large-scale SNP interrogations.

^dMutagenic PCR-RFLP is the most suitable method for SNP validation or genotyping across the board for a typical molecular genetics laboratory.

sented a novel systematic, mutagenic PCR-RFLP system for streamlined SNP validation and genotyping.

Stephens et al. (26) have investigated 3899 polymorphic sites in 313 human genes in four (Caucasian, Asian, Hispanic, and African American) populations of diverse ancestry. Only 21% of those polymorphisms are found to be "cosmopolitan" (observed in all four populations). Lee et al. (18) interrogated 300 published coding SNPs, and found that 23 of them were monomorphic in Koreans (18). Our results showed that only about 43% of the published SNPs are polymorphic in this Chinese population, which is consistent with previous SNP validations (27). We also found that over 70% of these validated SNPs were C→T or G→A (a mirror image of C→T) transitions, which is consistent with previous literature (4). More than half of the SNPs were located in CpG or GpC dinucleotides, strongly supporting the idea that

CpG or GpC is a mutation hotspot (30).

SNP genotyping is cost driven, and it is important to choose the most reliable and economic genotyping technology. A range of high-throughput methods for SNP typing are currently available, such as allele-specific amplification (24), real-time PCR (15), oligonucleotide ligation assay (3), single base extension-fluorescent polarization (6), and mass spectrometry (14). We compared the estimated cost, initial investment, and assay robustness for mutagenic PCR-RFLP, real-time PCR, oligonucleotide ligation assay, allele-specific amplification, single base extension fluorescent polarization, and mass spectrometry (Table 2). Overall, the cost of mutagenic PCR-RFLP assay is generally low (\$0.30–\$0.50/SNP), especially when we primarily depend on the low-cost restriction enzymes (less than \$0.05/U; *NcoI*, for example), and the estimated cost is comparable with other large-scale SNP genotyping methods [e.g.,

the mass spectrometry technology of Sequenom (San Diego, CA, USA) costs \$0.30–\$0.50/SNP, with no consideration of the software license and installation/maintenance costs for the sophisticated instrumentation].

Recently, DNA pooling has become an interesting strategy for two general applications: (i) linkage disequilibrium estimation from pooled DNA samples and (ii) allele frequency estimation from pooled DNA samples. In regard to linkage disequilibrium estimation, for medium to large (e.g., 10³–10⁵) sample sizes, mini-pooling could be an attractive cost-saving method. However, individual genotype information is partially missing because of the DNA pooling, which may constitute a critical loss of information. Therefore, one should exercise caution in considering the pooling strategy in linkage disequilibrium-based association studies. For allele frequency estimation, one needs to obtain very accurate measurements

Short Technical Reports

of the DNA concentration using a very sensitive and sophisticated detection method [e.g., PicoGreen® DNA quantitation instrumentation using the Picofluor™ Handheld Dual Channel Fluorometer (Turner Biosystems, Sunnyvale, CA, USA) (22)] to reduce the DNA quantification error, and the allele frequency estimation can be calculated according to the peak heights shown on pyrograms using Pyrosequencing™ methods (22,28). Because we suggested the SNP validation to be done on a small-scale sample (e.g., DNA from the 48 cell lines presented here), it is not difficult to obtain individual genotypes and then estimate the allele frequency by allele counting. Therefore, allele frequency estimation can be conveniently inferred from individual genotyping, although pooling can be considered when sample size is considerably large.

Direct sequencing is a popular SNP discovery and validation approach.

However, it is quite expensive to be used as a standard tool for SNP validation for a large sample size. Another disadvantage of sequencing is that after a SNP is validated, another genotyping assay has to be developed for further association studies, which turns out to be a nontrivial task when the number of SNPs is only moderate. By contrast, the mutagenic PCR-RFLP method can be easily applied to both SNP validation and genotyping.

The SNPkit mutagenic PCR-RFLP system provides a method with advantages in terms of complexity, time, and cost. This method is particularly attractive for application in conducting genome scan studies based on SNP maps (17) or SNP-based association analyses (2). The SNPkit package can provide multiple mutagenic PCR-RFLP designs, which gives us ample opportunity for quality control. Because the genotype is determined by

the specificity of the restriction endonuclease, RFLPs using different enzymes can be considered as independent genotyping essays. By repeating eight SNPs with alternative restriction enzymes, we confirmed the SNP validation and genotyping results (Table 1). From the software output, many SNPs have multiple RFLPs, and we selected these eight for demonstration purposes. The result showed 100% concordance between two RFLP assays on the same SNP. Incomplete digestion is a common problem with RFLP genotyping. Because our strategy allows us to select an economical and robust restriction endonuclease, we could either increase enzyme concentration or confirm it with alternative RFLPs to avoid possible incomplete digestions.

In summary, the SNPkit and mutagenic PCR-RFLP system is an easy and reliable tool for SNP validation and genotyping, which is essential for dissecting the genetic underpinnings of human diseases, molecular diagnoses, and predictive medicine. For example, the P-450 SNPs thought to be involved in drug response variance (16), can be designed for a SNPkit panel with multiplex mutagenic RFLP assays for rapid genotyping. This system can also be applied in genome-wide screening using high-density SNP maps for species other than *Homo sapiens*, such as the *Caenorhabditis elegans* (29). The SNPkit software package with documentation is available by written request from the author.

REFERENCES

1. **Ahmadian, A. and J. Lundeberg.** 2002. A brief history of genetic variation analysis. *BioTechniques* 32:1122-1128.
2. **Bader, J.S.** 2001. The relative power of SNPs and haplotype as genetic markers for association tests. *Pharmacogenomics* 2:11-24.
3. **Brinson, E.C., T. Adriano, W. Bloch, C.L. Brown, C.C. Chang, J. Chen, F.A. Eggerding, P.D. Grossman, et al.** 1997. Introduction to PCR/OLA/SCS, a multiplex DNA test, and its application to cystic fibrosis. *Genet. Test.* 1:61-68.
4. **Brookes, A.J.** 1999. The essence of SNPs. *Gene* 234:177-186.
5. **Brookes, A.J.** 2001. HGBASE—a unified human SNP database. *Trends Genet.* 17:229.
6. **Chen, X., L. Levine, and P.Y. Kwok.** 1999. Fluorescence polarization in homogeneous nucleic acid analysis. *Genome Res.* 9:492-498.
7. **Collins, F.S.** 1999. Shattuck lecture—medical

Short Technical Reports

- and societal consequences of the Human Genome Project. *N. Engl. J. Med.* 341:28-37.
8. Cox, D., C. Boillot, and F. Canzian. 2001. Data mining: efficiency of using sequence databases for polymorphism discovery. *Hum. Mutat.* 17:141-150.
9. Gallagher, G., J. Eskdale, H.H. Oh, S.D. Richards, D.A. Campbell, and M. Field. 1997. Polymorphisms in the TNF gene cluster and MHC serotypes in the West of Scotland. *Immunogenetics* 45:188-194.
10. Halushka, M.K., J.B. Fan, K. Bentley, L. Hsie, N. Shen, A. Weder, R. Cooper, R. Lipschutz, et al. 1999. Patterns of single-nucleotide polymorphisms in candidate genes for blood-pressure homeostasis. *Nat. Genet.* 22:239-247.
11. Hattori, M., S. Nanko, X.Y. Dai, R. Fukuda, and H. Kazamatsuri. 1994. Mismatch PCR RFLP detection of DRD2 Ser311Cys polymorphism and schizophrenia. *Biochem. Biophys. Res. Commun.* 202:757-763.
12. Hiratsuka, M., T. Inoue, F. Omori, Y. Agatsuma, Y. Kishikawa, and M. Mizugaki. 2000. Detection assay of rare variants of the *thiopurine methyltransferase* gene by PCR-RFLP using a mismatch primer in a Japanese population. *Biol. Pharm. Bull.* 23:1090-1093.
13. Irizarry, K., V. Kustanovich, C. Li, N. Brown, S. Nelson, W. Wong, and C.J. Lee. 2000. Genome-wide analysis of single-nucleotide polymorphisms in human expressed sequences. *Nat. Genet.* 26:233-236.
14. Jurinke, C., D. van den Boom, C.R. Cantor, and H. Koster. 2001. Automated genotyping using the DNA MassArray technology. *Methods Mol. Biol.* 170:103-116.
15. Kalinina, O., I. Lebedeva, J. Brown, and J. Silver. 1997. Nanoliter scale PCR with TaqMan detection. *Nucleic Acids Res.* 25:1999-2004.
16. Kato, R., Y. Yamazoe, and T. Yasumori. 1990. Pharmacogenetics and polymorphism of human P-450 which activates and detoxicates xenobiotics and carcinogens. *Princess Takamatsu Symp.* 21:45-53.
17. Kruglyak, L. 1997. The use of a genetic map of biallelic markers in linkage studies. *Nat. Genet.* 17:21-24.
18. Lee, S.G., S. Hong, Y. Yoon, I. Yang, and K. Song. 2001. Characterization of publicly available SNPs in the Korean population. *Hum. Mutat.* 17:281-284.
19. Marth, G., R. Yeh, M. Minton, R. Donaldson, Q. Li, S. Duan, R. Davenport, R.D. Miller, and P.Y. Kwok. 2001. Single-nucleotide polymorphisms in the public domain: how useful are they? *Nat. Genet.* 27:371-372.
20. Marth, G.T., I. Korf, M.D. Yandell, R.T. Yeh, Z. Gu, H. Zakeri, N.O. Stitzel, L. Hillier, et al. 1999. A general approach to single-nucleotide polymorphism discovery. *Nat. Genet.* 23:452-456.
21. Nakashima, H., T. Otsuka, Y. Ohba, M. Akahoshi, S. Nagano, E. Ogami, Y. Arinobu, K. Miyake, et al. 1999. Two polymorphisms within interleukin-3 (hIL3) gene detected by mismatch PCR/RFLP. *Genes Immun.* 1:156-158.
22. Neve, B., P. Froguel, L. Corset, E. Vaillant, V. Vatin, and P. Boutin. 2002. Rapid SNP allele frequency determination in genomic DNA pools by Pyrosequencing™. *BioTechniques* 32:1138-1142.
23. Niu, T., C. Chen, J. Yang, B. Wang, Z. Wang, N. Schork, Z. Fang, and X. Xu. 1999. Blood pressure and the T174M and M235T polymorphisms of the angiotensinogen gene. *Ann. Epidemiol.* 9:245-253.
24. Okayama, H., D.T. Curiel, M.L. Brantly, M.D. Holmes, and R.G. Crystal. 1989. Rapid, nonradioactive detection of mutations in the human genome by allele-specific amplification. *J. Lab. Clin. Med.* 114:105-113.
25. Parsons, B.L. and R.H. Hefflich. 1997. Genotypic selection methods for the direct analysis of point mutations. *Mutat. Res.* 387:97-121.
26. Stephens, J.C., J.A. Schneider, D.A. Tanquay, J. Choi, T. Acharya, S.E. Stanley, R. Jiang, C.J. Messer, et al. 2001. Haplotype variation and linkage disequilibrium in 313 human genes. *Science* 293:489-493.
27. Taillon-Miller, P., E.E. Piernot, and P.Y. Kwok. 1999. Efficient approach to unique single-nucleotide polymorphism discovery. *Genome Res.* 9:499-505.
28. Wasson, J., G. Skolnick, L. Love-Gregory, and M.A. Permut. 2002. Assessing allele frequencies of single nucleotide polymorphisms in DNA pools by Pyrosequencing™ technology. *BioTechniques* 32:1144-1152.
29. Wicks, S.R., R.T. Yeh, W.R. Gish, R.H. Waterston, and R.H. Plasterk. 2001. Rapid gene mapping in *Caenorhabditis elegans* using a high-density polymorphism map. *Nat. Genet.* 28:160-164.
30. Youssoufian, H., H.H. Kazazian, Jr., D.G. Phillips, S. Aronis, G. Tsiftis, V.A. Brown, and S.E. Antonarakis. 1986. Recurrent mutations in haemophilia A give evidence for CpG mutation hotspots. *Nature* 324:380-382.

We thank Dr. Mark Seielstad for careful reading of the manuscript. This study is supported in part by the National Institute of Arthritis and Musculoskeletal and Skin Diseases grant no. RO1 AR45651-OA1 and the National Institute of Environmental Health Sciences Kresge Center grant no. ES-00002. Address correspondence to Dr. Xiping Xu, Program for Population Genetics, Harvard School of Public Health, Boston, MA, USA. e-mail: xu@hsph.harvard.edu

Received 1 May 2002; accepted 18 June 2002.

K. Hao¹, T. Niu¹, C. Sango-koya², J. Li¹, and X. Xu^{1,3,4}
¹Harvard School of Public Health
Boston, MA

²Stanford University
Stanford, CA, USA

³Beijing Medical University,
Beijing, China

⁴Brigham and Women's Hospital
Harvard Medical School
Boston, MA, USA

Directional Genome Walking Using PCR

BioTechniques 33:830-834 (October 2002)

ABSTRACT

We describe here a PCR-based "directional genome walking" protocol. The basic procedure for the amplification consists of two rounds of PCR. A primary PCR was performed, on the genomic DNA using a biotinylated primer specific to a known sequence in the genome along with four universal walker primers that were designed with partial degeneracy. The biotinylated primary PCR products were immobilized on streptavidin-linked paramagnetic beads. This step removed all nonspecific amplification products, and the purified template was used for the second PCR using a nested primer and the walker primer-2 to increase specificity. This technique is potentially useful for cloning promoter regions and has been successfully used to isolate 5'-flanking genomic regions of many cDNA clones previously isolated by us.

INTRODUCTION

The traditional approach for "walking" from regions of known sequence into flanking DNA sequences involved the successive probing of libraries with clones obtained from prior screenings. This method, besides being laborious, was also time consuming. Though PCR is an effective method for selectively amplifying specific DNA segments, conventional PCR only allows the amplification of sequences within known boundaries. Therefore, sequence information at the extremities of the DNA fragment to be amplified is a prerequisite for selective amplification of specific DNA, thus posing a major limitation on the use of the PCR in the amplification and isolation of these unknown regions.

Numerous modifications have been made to existing protocols for the amplification of an unknown DNA sequence that flank regions of known sequences, by PCR (5). They include inverse PCR (13), panhandle PCR (17),