RNeasy-purified RNA yields reverse transcription PCR (RT-PCR) products of the expected size and renders contaminating gDNA undetectable by a PCR assay. Accordingly, this method allows reliable RT-PCR analyses of both genes exhibiting retropseudogenes and genes without introns. There have been reports of primer pairs for *GAPDH* (8) and *ACTB* (16) that were specially designed to exclude amplification of retropseudogenes. However, this method is time-consuming, and appropriate primer pairs have to be designed for every new gene analyzed. By contrast, the procedure described in this report is efficient and can be readily used with any existing primer pairs.

ACKNOWLEDGMENTS

This work was supported by a grant from the Deutsche Forschungsgemeinschaft (Bu 592/4-3). We wish to thank Ms. Renate Francke for excellent technical help.

REFERENCES

1. Vanin, E.F. 1985. Processed pseudogenes: characteristics and evolution. *Annu. Rev. Genet.* 19:253-272.
2. Goncalves, I., L. Duret, and D. Mouchiroud. 2000. Nature and structure of human genes that generate retropseudogenes. *Genome Res.* 10:672-678.
3. Johnson, K.R., D.A. Lehn, and R. Reeves. 1989. Alternative processing of mRNAs encoding mammalian chromosomal high-mobility-group proteins HMGI and HMGI-Y. *Mol. Cell. Biol.* 9:2114-2123.
4. Rogalla, P., Z. Borda, K. Meyer-Bolte, K.H. Tran, S. Hauke, R. Nimzyk, and J. Bullerdiek. 1998. Mapping and molecular characterization of five HMGI-related DNA sequences. *Cytogenet. Cell Genet.* 83:124-129.
5. Blank, C., P. Rogalla, K.H. Tran, and J. Bullerdiek. 2000. A novel high mobility group protein gene is a candidate for Xp22 abnormalities in uterine leiomyomas and other benign tumors. *Cancer Genet. Cytogenet.* 121:172-180.
6. Ng, S.Y., P. Gunning, R. Eddy, P. Ponte, J. Leavitt, T. Shows, and L. Kedes. 1985. Evolution of the functional human beta-actin gene and its multi-pseudogene family: conservation of noncoding regions and chromosomal dispersion of pseudogenes. *Mol. Cell. Biol.* 5:2720-2732.
7. Hanauer, A. and J.L. Mandel. 1984. The glyceraldehyde 3 phosphate dehydrogenase gene family: structure of a human cDNA and of an X chromosome linked pseudogene; amazing complexity of the gene family in mouse. *EMBO J.* 3:2627-2633.
8. Lehmann, M.H., J. Weber, O. Gastmann, and H.H. Sigusch. 2002. Pseudogene-free amplification of human GAPDH cDNA. *Bio-Techniques* 33:766-770.
9. Gentles, A.J. and S. Karlin. 1999. Why are human G-protein-coupled receptors predominantly intronless? *Trends Genet.* 15:47-49.
10. Sutton, D.H., G.L. Conn, T. Brown, and A.N. Lane. 1997. The dependence of DNase I activity on the conformation of oligodeoxynucleotides. *Biochem. J.* 321:481-486.
11. Muller, S., P. Scaffidi, B. Degryse, T. Bonaldi, L. Ronfani, A. Agresti, M. Beltrame, and M.E. Bianchi. 2001. New EMBO members' review: the double life of HMGB1 chromatin protein: architectural factor and extracellular signal. *EMBO J.* 20:4337-4340.
12. Chiappetta, G., V. Avantaggiato, R. Visconti, M. Fedele, S. Battista, F. Trapasso, B.M. Merciai, V. Fidanza, et al. 1996. High level expression of the HMGI (Y) gene during embryonic development. *Oncogene* 13:2439-2446.
13. Williams, A.J., W.L. Powell, T. Collins, and C.C. Morton. 1997. HMGI(Y) expression in human uterine leiomyomata. Involvement of another high-mobility group architectural factor in a benign neoplasm. *Am. J. Pathol.* 150:911-918.
14. Bandiera, A., D. Bonifacio, G. Manfioletti, F. Mantovani, A. Rustighi, F. Zancanati, A. Fusco, L. Di Bonito, and V. Giancotti. 1998. Expression of HMGI(Y) proteins in squamous intraepithelial and invasive lesions of the uterine cervix. *Cancer Res.* 58:426-431.
15. Flohr, A.M., P. Rogalla, U. Bonk, B. Puettmann, H. Buerger, J. Packeisen, W. Wosniok, et al. 2003. High mobility group protein HMGA1 expression in breast cancer reveals a positive correlation with tumour grade. *Histol. Histopathol.* 18:999-1004.
16. Raff, T., M. van der Giet, D. Endemann, T. Wiederholt, and M. Paul. 1997. Design and testing of beta-actin primers for RT-PCR that do not co-amplify processed pseudogenes. *BioTechniques* 23:456-460.

Received 8 May 2003; accepted 12 August 2003.

Address correspondence to Joern Bullerdiek, Center for Human Genetics, University of Bremen, Leobenerstr. ZHG, D-28359 Bremen, Germany. e-mail: bullerdiek@uni-bremen.de

A rapid assay for the quantitation of γ -glutamyl hydrolase using a fluorogenic peptide as substrate

Erin L. Volk^{1,2}, Jessica J. Pankuch³, Karen J. Chave², James K. Coward³, and Erasmus Schneider^{1,2}

¹University at Albany, Albany, ²Wadsworth Center, Albany, NY, and

³University of Michigan, Ann Arbor, MI, USA

BioTechniques 35:926-932 (November 2003)

Gamma-glutamyl hydrolase (γ -GH) is a lysosomal enzyme that catalyzes the cleavage of natural folylpoly- γ -glutamates, as well as that of the polyglutamates of several antifolate chemotherapeutic drugs (1). Cleavage of polyglutamate residues from antifolates, such as methotrexate (MTX), results in the generation of "free" drug, which is less effectively retained within the cell, thereby reducing its cytotoxicity (2). Increased γ -GH activity has therefore been associated with lower sensitivity to, or resistance against, multiple an-

tifolate drugs (3). These observations and further in vitro (4-6) and in vivo (7-9) studies have suggested that γ -GH activity may be clinically relevant as a possible prognostic marker for the efficacy of chemotherapy in several cancers such as childhood leukemia, for which antifolate therapy is widely used.

Several methods have been developed for the determination of γ -GH activity in cells (10-12). Current protocols measure γ -GH activity by incubating cell extracts in the presence of (radiolabeled) MTX polyglutamates

and quantitatively analyzing the cleavage products using high-performance liquid chromatography (10) or capillary electrophoresis (11). However, while these methods are adequate for end point analysis of enzymatic activity in a small number of samples, they are relatively cumbersome and slow, and ill suited for the analysis of clinical samples. A colorimetric assay (12) has also been described, but it requires three enzymatic steps. Like the assays using radiolabeled substrates, this technique does not allow real-time monitoring of enzyme activity.

Here, we describe a novel method for measuring γ -GH activity in whole cell lysates in real time that is both rapid and effective. By using a newly synthesized, internally quenched fluorogenic peptide, N-Me-pAB-Glu- γ -Glu- γ -Tyr(3-NO₂) (13), as a specific γ -GH substrate, the enzymatic activity of multiple samples can be determined continuously by monitoring the increase in fluorescence from the hydrolysis of the peptide. The activity can then be quantitated with the aid of a standard curve. The substrate is specific for γ -GH in so far that it is not cleaved by glutamate carboxypeptidase II/N-

acetylated- α -linked acidic dipeptidase (NAALADase) (J.J. Pankuch and J.K. Coward, unpublished observations).

To determine γ -GH enzymatic activity, 85 μ L of assay buffer consisting of 50 mM sodium acetate, pH 5.5, with 50 mM β -mercaptoethanol, and 10 μ L cell lysate (containing 50 μ g protein) or purified enzyme (50–400 ng protein) were added to each well of a 96-well microtiter plate. The reaction was initiated by the addition of 5 μ L of the 100 μ M fluorogenic substrate N-Me-pAB-Glu- γ -Glu- γ -Tyr(3-NO₂) (dissolved in 50 mM sodium acetate, 50 mM β -mer-

Figure 1. Time and concentration dependence of fluorogenic substrate cleavage by γ -glutamyl hydrolase. Various amounts of human baculovirus-produced γ -GH enzyme were incubated for up to 50 min and fluorescence readings taken every 5 min. (A) Fluorescence produced as a function of enzyme concentration. (B) Fluorescence produced as a function of incubation time. (C) Standard curve based on the rate of change in fluorescence between 0 and 40 min. (D) Standard curve based on the fluorescence at 30 min (end point method). Data for panels A and B are representative from a single experiment performed in triplicate. For panels C and D, the values shown are the mean ($\bar{x}$) from five individual experiments, each assayed either in duplicate or triplicate, with standard errors. The linear regression line (—) and 95% confidence intervals (-----) were determined with the aid of the Prism3® software package (Graphpad, San Diego, CA, USA). The specific activity of the purified γ -GH used was 0.23 nM/min/ μ g of MTX generated from MTX-G2 as determined by capillary electrophoresis.

captoethanol, pH 5.5) to each well. Fluorescence measurements were started immediately and were repeated at 5-min intervals over a period of 50 min at room temperature on an LS50B luminescence spectrometer (Perkin Elmer, Boston, MA, USA) with the following settings: 325-nm excitation wavelength, 415-nm emission wavelength, 10-nm bandpass, and 0.1 s/well read time. To generate a standard curve, pure human baculovirus-produced recombinant γ -GH enzyme (14) (generously provided by Dr. Thomas Ryan, Wadsworth Center) was used. The activity of various amounts of the enzyme was determined by incubating in the presence of the fluorogenic substrate and measuring fluorescence at 5-min intervals for 50 min. The raw fluorescence data were corrected for nonspecific background fluorescence by subtracting the average fluorescence values obtained from replicate wells that contained the complete assay mixture except for enzyme. No changes of background fluorescence were detected over time. As demonstrated in Figure 1, fluorescence increased linearly as a function of enzyme concentration (Figure 1A) and time (Figure 1B) up to the limits shown in the graphs. Standards containing the largest amounts of γ -GH protein produced the greatest levels of fluorescence as demonstrated by the increasing slopes in Figure 1B. At higher enzyme

concentrations (>400 ng) or longer incubation times (>40–50 min, depending on enzyme concentration), however, the results started to deviate from linearity and exhibited saturation behavior, presumably due to substrate depletion (data not shown). By plotting the change in fluorescence over time (i.e., the slopes from Figure 1B) or the fluorescence at a single time point (end point method) as a function of γ -GH protein quantity (Figure 1, C and D, respectively), a standard curve can be generated that shows a linear response over a broad range of enzyme activities and can therefore be used to quantitate γ -GH activity in whole-cell lysates. Assessments of inter-assay variability (Figure 1, C and D) revealed small standard errors and 95% confidence intervals, showing that the assay is highly reproducible. Together, these data indicate that the assay is suitable for the quantitative assessment of γ -GH activity.

To evaluate its practical usefulness, this method was used to screen for γ -GH transgene expression in whole-cell lysates from MCF7 cells (ATCC, Manassas, VA, USA) that had been stably transfected with a human γ -GH expression construct (16 clones) or the corresponding empty control vector (2 clones) (E.L. Volk and E. Schneider, unpublished observations). Lysates were prepared from cells that

cell lysates yielded a linear increase in fluorescence detected as a function of time. The fluorescence at 30 min or the slopes of the curves (i.e., the rates of fluorescence production) were used to calculate the amount of enzyme in the lysates from appropriate standard curves that had been established in parallel. Both methods gave similar results. Each cell lysate was analyzed three times in triplicate on different days. The results were reproducible with an average 14% coefficient of variation (CV) across the three experiments. To compare the new assay with an established assay, each cell lysate was also analyzed by capillary electrophoresis as previously described (15). As shown in Figure 2, there was a good correlation between the results obtained with the two assays. Furthermore, the three samples with the highest activity in both assays were from cells from three individual clones that had been transfected with γ -GH, indicating that the transfection had been successful. Thus, the assay proved to be useful as a rapid screening method to identify transfected clones that exhibited γ -GH overexpression.

In conclusion, we have expanded upon the current methodology for the measurement of γ -GH activity in whole-cell lysates. The recently synthesized γ -GH-specific fluorogenic substrate has allowed the development of a rapid detection assay that quantitates the continuous rate of enzymatic γ -linked polyglutamate cleavage using fluorescence measurements. The results are comparable to those obtained with a more conventional assay, and, when used with a standard curve, allow the calculation of absolute amounts of γ -GH protein present in a cell extract. The use of this easy method in a microplate format allows rapid, simultaneous, and reproducible analysis of multiple experimental and clinical samples to further enhance our understanding of the role of γ -GH in chemotherapeutic efficacy.

ACKNOWLEDGMENTS

We thank Dr. Thomas Ryan for providing the purified human γ -GH enzyme and the Wadsworth Center's Biochem-

Figure 2. γ -Glutamyl hydrolase activity in whole cell lysates. Cell lysates obtained from untransfected and γ -GH transfected MCF7 cells were prepared and assayed for γ -GH activity either by the method described herein or by capillary electrophoresis. Each lysate was measured three times in triplicate with the present method and once in triplicate by capillary electrophoresis. The results from both methods were normalized to those from untransfected MCF7 cells and the relative values from each method graphed against each other. $r^2=0.869$; P value for correlation <0.0001 .

istry Core Facility for the use of the fluorescence plate reader. The work in the authors' labs was supported by NIH grants CA72455 (E.S.) and CA28097 (J.K.C.), and grant C017924 (E.S.) from the NYS Health Research Science Board. For questions concerning the fluorogenic substrate, contact Dr. J. Coward at jkcoward@umich.edu.

REFERENCES

- McGuire, J.J. and J.K. Coward. 1984. Pteroylpolylglutamates: biosynthesis, degradation, and function, p. 135-190. In R.L. Blakley and S.J. Benkovic (Eds.), *Folates and Pteridines*. Chemistry and Biochemistry of Folates. John Wiley, New York.
- Galivan, J., T.J. Ryan, K. Chave, M. Rhee, R. Yao, and D. Yin. 2000. Glutamyl hydrolase. Pharmacological role and enzymatic characterization. *Pharmacol. Ther.* 85:207-215.
- Li, W., M. Waltham, W. Tong, B. Schweitzer, and J. Bertino. 1993. Increased activity of γ -glutamyl hydrolase in human sarcoma cell lines: a novel mechanism of intrinsic resistance to methotrexate. *Pharmacol. Ther.* 85:207-215.
- Rhee, M.S., Y. Wang, M.G. Nair, and J. Galivan. 1993. Acquisition of resistance to antifolates caused by enhanced γ -glutamyl hydrolase activity. *Cancer Res.* 53:2227-2230.
- Yao, R., M.S. Rhee, and J. Galivan. 1995. Effects of γ -glutamyl hydrolase on folyl- and antifolylpolyglutamates in cultured H35 hepatoma cells. *Mol. Pharmacol.* 48:505-511.
- Pizzorno, G., B.A. Moroson, A.R. Cashmore, O. Russello, J.R. Mayer, J. Galivan, M.A. Bunni, D.G. Priest, and G.P. Beardsley. 1995. Multifactorial resistance to 5,10-dideazatetrahydrofolic acid in cell lines derived from human lymphoblastic leukemia CCRF-CEM. *Cancer Res.* 55:566-573.
- Longo, G.S., R. Gorlick, W.P. Tong, S. Lin, P. Steinherz, and J.R. Bertino. 1997. γ -Glutamyl hydrolase and folylpolyglutamate synthetase activities predict polyglutamylation of methotrexate in acute leukemias. *Oncol. Res.* 9:259-263.
- Rots, M., R. Pieters, G. Peters, P. Noordhuis, C. van Zantwijk, G. Kaspers, K. Hahlen, U. Creutzig, A. Veerman, and G. Jansen. 1999. Role of folylpolyglutamates synthetase and folylpolyglutamate hydrolase in methotrexate accumulation and polyglutamylation in childhood leukemia. *Blood* 5: 1677-1683.
- Whitehead, M., D. Rosenblatt, M.-J. Vuchich, J. Shuster, A. Witte, and D. Beaulieu. 1990. Accumulation of methotrexate and methotrexate polyglutamates in lymphoblasts at diagnosis of childhood acute lymphoblastic leukemia: a pilot prognostic factor analysis. *Blood* 76:44-49.
- Samuels, L., L. Goutas, D. Priest, J. Piper, and F. Sirotnak. 1986. Hydrolytic cleavages of methotrexate gamma-polyglutamates by folylpolyglutamyl hydrolase derived from various tumors and normal tissues of the mouse. *Cancer Res.* 46:2230-2235.
- Waltham, M., S. Lin, W. Li, E. Goker, H. Gritsman, W. Tong, and J. Bertino. 1997. Capillary electrophoresis of methotrexate polyglutamates and its application in evaluation of gamma-glutamyl hydrolase activity. *J. Chromatogr.* 689:387-392.
- Wagh, P.V. and T.I. Kalman. 1992. A rapid colorimetric assay for gamma-glutamyl hydrolase (conjugase). *Anal. Biochem.* 207:1-5.
- Pankuch, J. and J. Coward. 2001. N-Me-pAB-Glu- γ -Glu- γ -Tyr(3-NO₂): an internally quenched fluorogenic γ -glutamyl hydrolase substrate. *Bioorg. Med. Chem. Lett.* 11: 1561-1564.
- Rhee, M.S., B. Lindau-Shepard, K.J. Chave, J. Galivan, and T.J. Ryan. 1998. Characterization of human cellular γ -glutamyl hydrolase. *Mol. Pharmacol.* 53:1040-1046.
- Chave, K.J., J. Galivan, and T.J. Ryan. 1999. Site-directed mutagenesis establishes cysteine-110 as essential for enzyme activity in human gamma-glutamyl hydrolase. *Biochem. J.* 343:551-555.

Received 6 June 2003; accepted 19 August 2003.

Address correspondence to Erasmus Schneider, Wadsworth Center, Empire State Plaza, Albany, NY 12201, USA. email: schneid@wadsworth.org

High yield preparation of genomic DNA from *Streptomyces*

Jasmina Nikodinovic¹, Kevin D. Barrow¹ and Jo-Anne Chuck²

¹University of New South Wales, Sydney and ²University of Western Sydney, Penrith South, Australia

BioTechniques 35:932-936 (November 2003)

Streptomyces species produce important drugs such as antibiotics, immunosuppressants, and antitumor compounds. The isolation of genomic DNA is imperative for the understanding of the biosynthesis of these compounds and has led to the rational design of new analogs (1-5). *Streptomyces* are Gram-positive bacteria, making DNA isolation difficult due to their resistance to cell lysis (6,7). Most methods use lysozyme and sodium dodecyl sulfate (SDS) for cell disruption. To further increase lysis, glycine is often incorporated into media to minimize peptidoglycan cross-linking; muramidases such as mutanolysin or grinding of mycelia are also commonly used (1,8-10). Compared with DNA isolation methods for *Escherichia coli*, most methods are time-consuming or low yielding, or give low-quality DNA (1). This report details an improved method for DNA isolation from *Streptomyces* species using achromopeptidase, lysozyme, and SDS for cell lysis that results in higher yield compared with current standard methods.

The addition of achromopeptidase was prompted by its use in protoplast generation in *Streptomyces*, suggesting that it interacts with the cell wall of the bacterium (11). It has also been used in the lysis of other Gram-positive organisms (12,13). Presumably the mode of action of the protease in *Streptomyces* is to cause disruption of the peptidoglycan layer through cleavage of *N*-acetylmuramoyl-L-alanine amide bonds together with D-Ala-Gly and Gly-Gly bonds as reported for *Staphylococcus aureus* (14). When incubated simultaneously with lysozyme disrupting glycosidic linkages in the polymer, the resulting bacterial structures are more susceptible to SDS lysis. The increase in cell lysis would lead to an increase in DNA concentration for purification in the later stages of the protocol.

Streptomyces nodosus (ATCC 14899; American Type Culture Collection, Manassas, VA, USA), *S. noursei* (ATCC 11455), *S. avermitilis* (NRRL 3165; Agricultural Research Service Culture Collection, Peoria, IL, USA)