9q and 20, where, respectively, chromosomal loss and gain is frequently detected in SCA III ovarian tumors (7,8). After selecting the genes on these respective chromosomes, hierarchical clustering is performed. Sorting the genes on chromosomal order reveals particular chromosomal regions with under- or overexpressed genes in subsets of SCA III tumors. Possible retention or amplification on chromosome 9p in all tumors and predominant chromosomal loss on chromosome 9q in a number of SCA III tumors is observed. When analyzed in the same way, chromosome 20 reveals potential amplified regions in the majority of tumors (Figure 1, B and C).

Application of this method gives direct insight into the chromosomal distribution of up- or down-regulated genes. It reveals continuous regions with gene expression biases. The visualization of these biases is a first step in better understanding of the relations between physical loss or gain of chromosomal regions and the effects of gene expression of genes located in these regions. This observation may also affect the significance of microarray expression data. Up-regulating of a gene within a region of other up-regulated genes may be due to genomic instability and can be less significant than a solitary up-regulated gene. The macro and the user manual are available from the authors at <http://cms-dev.bioasp.nl/applicationservices/extra/genesort?view=MoreDetail> and also as supplementary material at the BioTechniques web site at <http://www.BioTechniques.com/April04/vanEijkSupplementary.html>.

ACKNOWLEDGMENTS

This work was supported by the Leiden University Medical Center. We thank Dr. Han Rauwerda and the Bioinformatics Application Service Provider (BioASP) for making the guide available through the BioASP.

REFERENCES

1. Kallioniemi, A., O.P. Kallioniemi, D. Sudar, D. Rutovitz, J.W. Gray, F. Waldman, and D. Pinkel. 1992. Comparative genomic hybridization for molecular cytogenetic analysis of solid tumors. *Science* 258:818-821.
2. Crawley, J.J. and K.A. Furge. 2002. Identification of frequent cytogenetic aberrations in hepatocellular carcinoma using gene-expression microarray data. *Genome Biol.* 3:RESEARCH0075.
3. Caron, H., B. van Schaik, M. van der Mee, F. Baas, G. Riggins, P. van Sluis, M.C. Hermus, R. van Asperen, et al. 2001. The human transcriptome map: clustering of highly expressed genes in chromosomal domains. *Science* 29:1289-1292.
4. Pollack, J.R., T. Sorlie, C.M. Perou, C.A. Rees, S.S. Jeffrey, P.E. Lonning, R. Tibshirani, D. Botstein, et al. 2002. Microarray analysis reveals a major direct role of DNA copy number alteration in the transcriptional program of human breast tumors. *Proc. Natl. Acad. Sci. USA* 99:12963-12968.
5. Hughes, T.R., C.J. Roberts, H. Dai, A.R. Jones, M.R. Meyer, D. Slade, J. Burchard, S. Dow, et al. 2000. Widespread aneuploidy revealed by DNA microarray expression profiling. *Nat. Genet.* 25:333-337.
6. Harding, M.A., K.C. Arden, J.W. Gildea, J.J. Gildea, E.J. Perlman, C. Viars, and D. Theodorescu. 2002. Functional genomic comparison of lineage-related human bladder cancer cell lines with differing tumorigenic and metastatic potentials by spectral karyotyping, comparative genomic hybridization, and a novel method of positional expression profiling. *Cancer Res.* 62:6981-6989.
7. Hu, J., V. Khanna, M.W. Jones, and U. Surti. 2003. Comparative study of primary and recurrent ovarian serous carcinomas: comparative genomic hybridization analysis with a potential application for prognosis. *Gynecol. Oncol.* 89:369-375.
8. Bayani, J., J.D. Brenton, P.F. Macgregor, B. Beheshti, M. Albert, D. Nallainathan, J. Karaskova, B. Rosen, et al. 2002. Parallel analysis of sporadic primary ovarian carcinomas by spectral karyotyping, comparative genomic hybridization, and expression microarrays. *Cancer Res.* 62:3466-3476.
9. Irizarry, R.A., B. Hobbs, F. Collin, Y.D. Beazer-Barclay, K.J. Antonellis, U. Scherf, and T.P. Speed. 2003. Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* 4:249-264.

Received 14 November 2003; accepted 11 February 2004.

Address correspondence to Ronald van Eijk, Department of Pathology, Leiden University Hospital, Albinusdreef 2, Building 1, P1-41, P.O. Box 9600, 2300 RC Leiden, The Netherlands. e-mail: R.van_Eijk@lumc.nl

Production and assay of a transcription activator CRP in coupled in vitro transcription and translation

Shumo Liu¹ and Albert Libchaber²

¹NEC Laboratories America, Princeton, NJ and ²Rockefeller University, New York, NY, USA

BioTechniques 36:596-600 (April 2004)

We describe a method to produce an *Escherichia coli* transcription activator and assay its function in vitro in a coupled transcription and eukaryotic translation (CTT) reaction. The model transcription activator used here is CRP (also called CAP), a well-known *E. coli* catabolic regulator protein. In the presence of cyclic-AMP (cAMP), the CRP-cAMP complex binds to a CRP-dependent promoter, interacts with RNA polymerase and stimulates transcription at the promoter (1-4).

In the experiments described here, we use T3 RNA polymerase to produce CRP in vitro from a plasmid, pT3CRP, which has the CRP gene, *crp*, transcribed from a T3 promoter (Figure 1). The function of CRP is tested with a reporter plasmid, p42crp-luc, which has a synthetic CRP-dependent promoter and the firefly luciferase gene, *luc*. The synthetic promoter has the following features: (i) it has a -10 site of *E. coli* consensus promoter to ensure a high promoter activity upon activation; (ii)

598 BioTechniques

vided to the four individual reactions to which other ingredients specific to each reaction are added.

At the time of experiment, the reactions are prepared in clear 0.2-mL PCR tubes at 0°C. The temperature of the tubes is raised to room temperature (22°–23°C) to start CTT. Although the CTT reaction temperature can be raised to at least 30°C, we choose room temperature for convenience. Each reaction is monitored with a photomultiplier tube (PMT) of a four-channel luminometer. The relative luminescence unit (RLU) is recorded with a computer. The transcription level of *luc* can be estimated from RLU values. The variation of RLU is around 10% among repeated reactions.

Functional CRP can be produced and assayed either in two separate CTT reactions (referred to as pre-made CRP) or in a combined reaction (referred to as in situ activation). Premade CRP is produced in 10 μ L CTT reaction with 2% (v/v) T3 RNA polymerase (Promega; included in the CTT kit) and 0.15 nM pT3CRP DNA. The reaction lasts for 100 min at room temperature, and it is terminated by adding 5 μ L glycerol and stored at -20°C as premade CRP preparation. In the following experiment, we use 3% (v/v) premade CRP preparation and 2 μ M cAMP (Sigma, St. Louis, MO, USA) in an assay reaction (Figure 2A). For in situ activation, we add T3RNA polymerase (2%, v/v) and pT3CRP DNA (0.075 nM) directly to another assay reaction. As a reference, we include a positive control reaction with pConsensus-*luc*. This plasmid has a constitutively active consensus promoter in place of the CRP-dependent promoter on p42crp-*luc* (Figure 1). Also included is a negative control reaction with premade CRP but without cAMP.

Figure 2A shows promoter activity of reactions as reported by *luc* expression. In the presence of cAMP and CRP, the CRP-dependent promoter activity is as active as the consensus promoter. CRP can be supplied to the promoter either in situ or premade. In the case of in situ activation, there is an additional lag of *luc* expression, which reflects the time needed to produce enough CRP. In contrast to the active state, in the ab-

sence of cAMP (negative control) CRP-dependent promoter gives very low *luc* expression. Therefore, the transcription at the CRP-dependent promoter is clearly cAMP dependent, confirming the identity of CRP as the activator protein. The low level of *luc* expression in negative control also shows that wheat germ extract has no detectable amount of either endogenous cAMP or cAMP-independent activator.

Next, we tested in situ activation dependence on pT3CRP DNA concentration (Figure 2B). The *luc* expression is approximately proportional to the concentration of pT3CRP. Evidently CRP-dependent promoter activity is proportional to CRP concentration, which in turn, is proportional to pT3CRP concentration. The very low *luc* expression without pT3CRP indicates that wheat germ extract has no detectable amount of endogenous CRP-like activator. Finally, we observe CRP activation at various concentrations of cAMP (Figure 2C). The activity at 2 μ M cAMP is near maximum (higher than 2 μ M cAMP not shown), and the half-activation concentration of cAMP is between 0.6 and 0.8 μ M. The cAMP dose response is comparable to what has been previously reported (8,9).

In the experiments, we apply CTT to produce CRP from its DNA coding sequences and to observe CRP transcription activation in near real-time. We use wheat germ extract to avoid the interference from endogenous CRP and cAMP, which are usually present in *E. coli* 30-S extract. Combining activator production and assay in one, the reaction mimics a simple in vivo gene expression circuit.

REFERENCES

1. Kolb, A., S. Busby, H. Buc, S. Garges, and S. Adhya. 1993. Transcriptional regulation by cAMP and its receptor protein. *Annu. Rev. Biochem.* 62:749-795.
2. Busby, S. and R.H. Ebright. 1999. Transcription activation by catabolite activator protein (CAP). *J. Mol. Biol.* 293:199-213.
3. Harman, J.G. 2001. Allosteric regulation of the cAMP receptor protein. *Biochim. Biophys. Acta* 1547:1-17.
4. Savery, N., V. Rhodius, and S. Busby. 1996. Protein-protein interactions during transcription activation: the case of the *Escherichia coli* cyclic AMP receptor protein. *Philos. Trans. R Soc. Lond. B Biol. Sci.* 351:543-550.
5. Ebright, R.H. 1993. Transcription activation at Class I CAP-dependent promoters. *Mol. Microbiol.* 8:797-802.
6. Ushida, C. and H. Aiba. 1990. Helical phase dependent action of CRP: effect of the distance between the CRP site and the -35 region on promoter activity. *Nucleic Acids Res.* 18:6325-6330.
7. Kolb, V.A., E.V. Makeyev, and A.S. Spirin. 1994. Folding of firefly luciferase during translation in a cell-free system. *EMBO J.* 13:3631-3637.
8. Nisseley, S.P., W.B. Anderson, M.E. Gottesman, R.L. Perlman, and I. Pastan. 1971. In vitro transcription of the gal operon requires cyclic adenosine monophosphate and cyclic adenosine monophosphate receptor protein. *J. Biol. Chem.* 246:4671-4678.
9. Takahashi, M., B. Blazy, A. Baudras, and W. Hillen. 1989. Ligand-modulated binding of a gene regulatory protein to DNA. Quantitative analysis of cyclic-AMP induced binding of CRP from *Escherichia coli* to non-specific and specific DNA targets. *J. Mol. Biol.* 207:783-976.

Received 2 September 2003; accepted 11 February 2004.

Address correspondence to Shumo Liu, Department of Physics, University of California at San Diego, 9500 Gilman Drive, MC0374, La Jolla, CA 92093, USA. e-mail: sliu@physics.ucsd.edu