A parallel purification of the ssDNA was done by using the conventional method, which took about 1.5 h, compared with about 15 min for the above method. An equivalent amount of phages was resuspended in 0.5 mL TE and subjected to RNase A digestion at 37°C for 30 min, followed by phenol, phenol/chloroform extractions until no white floccule-like residue was seen at the interface between the organic and aqueous phase after centrifugation. After chloroform extraction followed by ethanol-precipitation and vacuum-drying, the centrifuged DNA was resuspended in 30 μ L TE. Three microliters of the sample were analyzed by electrophoresis on 1% agarose. The result of analysis showed no chromosomal DNA contamination present in the sample prepared by the above-described method, whereas this contamination was present in the sample prepared by the conventional method (Figure 1). Contamination was especially great when the yield of secretion of phagemids was low and when a large volume of culture was needed to enrich the phages (data not shown).

Figure 1. Comparison of DNA purified by two different methods. (A) Chromosomal DNA, (B) ssM13KO7 DNA and (C) ssPhagemid DNA. Lane 1, DNA purified by the conventional method; and lane 2, DNA purified by the new method described in this paper.

This method results in a preparation of single-stranded phagemid DNA free of chromosomal DNA contamination, and it eliminates several steps, including phenol and phenol/chloroform extraction, ethanol precipitation and vacuum drying, during which loss of DNA may occur. The method is also economical and comparable in price to the conventional one.

REFERENCES

1. Boom, R., C.J. Sol, M.M. Salimans, C.L. Jansen, P.M. Wertheim-van Dillen and J. Vander Noordaa. 1990. Rapid and simple method for purification of nucleic acids. *J. Clin. Microbiol.* 28:495-503.
2. Lin, J.-J., M. Smith, J. Jessee and F. Bloom. 1992. DH11S: An *Escherichia coli* strain for preparation of single-stranded DNA from phagemid vectors. *BioTechniques* 12:718-721.
3. Marko, M.A., R. Chipperfield and H.C. Birnboim. 1982. A procedure for the large-scale isolation of highly purified plasmid DNA using alkaline extraction and binding to glass powder. *Anal. Biochem.* 121:382-387.
4. Ohlsson, R., C.C. Hentschel and J.G. Williams. 1978. A rapid method for the isolation of circular DNA using an aqueous two-phase partition system. *Nucleic Acids Res.* 5:583-590.
5. Titus, D.E. 1991. Single-stranded DNA template preparation from M13 vectors or pGEM-Zf phagemid, p.111. *Promega Protocols and Applications Guide*, 2nd ed. Promega Corporation, Madison, WI.

This work was supported by a grant from Neurex, Inc. (Menlo, CA, USA). Address correspondence to Victor Gurewich, Vascular Research Laboratory, Institute for the Prevention of Cardiovascular Disease, Deaconess Hospital, Harvard Medical School, Boston, MA 02215, USA.

Received 10 October 1995; accepted 12 January 1996.

Wei Tang, Jian-Ning Liu and Victor Gurewich
*Harvard Medical School
 Boston, MA, USA*

RNA/DNA Mini-Prep From a Single Sample of Orchid Tissue

BioTechniques 21:54-56 (July 1996)

Initial analysis of genetically transformed plant materials generally includes both Southern and Northern blot analyses to verify transgene integration and transcriptional activation, respectively. Isolation of high-quality DNA and RNA for analysis from small amounts of tissue is a critical step in this process. For many plants, callus tissue or regenerated plantlets grow sufficiently fast to provide adequate amounts of tissue for nucleic acid isolation and analysis in a short period of time. Unfortunately, plants that grow more slowly have an extensive delay between the time the transgene is introduced to the time sufficient tissue is available for analysis. The orchid, *Orchidaceae*, is one such family of plants.

Figure 1. Ethidium bromide-stained 1.5% formaldehyde-agarose gel of total RNA. Lane M, Markers 2.3-kb synthetic RNA (Life Technologies, Gaithersburg, MD, USA) and 1.9-kb luciferase control RNA (Promega, Madison, WI, USA); lanes B1 and B2, *Brassia rex* "Sakata" $\times$ *B. verucosa*; lanes P1 and P2, (*Phalaenopsis* Su's Red Lip "Ching Hua II" $\times$ *Doritaenopsis* Amour Rojo "Bonnie Vasquez") $\times$ *Phalaenopsis stuartiana*.

Benchmarks

Two recent reports for gene transfer into *Dendrobium* orchids describe DNA isolation from tissues 8–21 months after they were transformed (2,4). RNA was isolated from a second, independent sample of tissue for Northern analysis (2) after 8 months. Here, we report a rapid mini-prep method for isolating both RNA and DNA from a very small (20–40 mg fresh weight), single sample of orchid tissue, allowing both Southern and Northern analyses of transformed tissues in as little as 4 months after transformation and selection.

Nucleic acids were isolated by the hexadecyltrimethylammonium bromide (CTAB) method of Doyle and Doyle (3) as modified by Rowland and Nguyen (5) but with the polyethylene glycol (PEG) precipitation step omitted. Briefly, fresh tissue (protocorms, plantlets or leaves) was placed in a microcentrifuge tube and ground to a powder in liquid nitrogen with a heat-sealed 1000- μ L pipet tip. Three hundred microliters of extraction buffer (6) were added (3% wt/vol CTAB, 1.42 M NaCl, 20 mM EDTA, 100 mM Tris-HCl, pH 8.0, 2% wt/vol polyvinylpyrrolidone [PVP-40; Sigma Chemical, St. Louis, MO, USA], 5 mM ascorbic acid), and the tissue was further homogenized for 2 min. Samples were

heated at 65°C for 15 min, then extracted with chloroform. One-fifth volume of 5% CTAB solution (5% CTAB, 0.7 M NaCl) was added to the aqueous phase, and the samples were heated again and extracted with chloroform. RNA was isolated from the aqueous phase by the addition of 1/2 volume 7.5 M NH_4OAc (1), incubated on ice 15 min and centrifuged 10 min at 16000 $\times$ g in an Eppendorf® Model 5415 Microcentrifuge (Brinkmann Instruments, Westbury, NY, USA). DNA was precipitated from the supernatant by the addition of two volumes of ethanol, incubated at -20°C for 15 min and centrifuged for 15 min. RNA was quantitated by absorbance at 260 nm, and DNA was quantitated by fluorometry. Yields from this method are 15–50 μ g DNA and 20–150 μ g RNA depending on fresh weight of starting material.

Figure 1 shows the high-quality profile for total RNA isolated from two diverse types of orchids using this method. These RNAs have been used successfully on Northern blots (data not shown). Figure 2 shows the high-quality total genomic DNAs recovered by using this procedure after digestion with *Bam*HI. Each purified DNA sample was easily digested with all the restriction enzymes tested and also used successfully on Southern blots (data not shown). This method has also been used successfully for the isolation of DNA from roots of a number of woody species including blueberry (93 μ g/mg dry weight), rhododendron (38 μ g/mg dry weight) and oak (68 μ g/mg dry weight) with typical yields indicated in parentheses. This suggests that this method may have wide application to many plant species.

REFERENCES

1. **Anonymous.** 1994. A simple method for RNA purification from *in vitro* transcription reactions. *Epicentre Forum* 1:8.
2. **Chia, T-F., Y-S. Chan and N-H. Chua.** 1994. The firefly luciferase gene as a non-invasive reporter for *Dendrobium* transformation. *The Plant J.* 6:441-446.
3. **Doyle, J.J. and J.L. Doyle.** 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull.* 19:11-15.
4. **Kuehnle, A.R. and N. Sugii.** 1992. Transformation of *Dendrobium* orchid using particle bombardment of protocorms. *Plant Cell Rep.* 11:484-488.
5. **Rowland, L.J. and B. Nguyen.** 1993. Use of polyethylene glycol for purification of DNA from leaf tissue of woody plants. *BioTechniques* 14:734-736.
6. **Stewart, C.N. Jr. and L.E. Via.** 1993. A rapid CTAB DNA isolation technique useful for RAPD fingerprinting and other PCR applications. *BioTechniques* 14:748-750.

This is paper No. 3182 of the Journal series of the Rhode Island Agricultural Experiment Station, Kingston. Address correspondence to Joel M. Chandlee, University of Rhode Island, Department of Plant Sciences, Kingston, RI 02881, USA. Internet: mtu101@uriacc.uri.edu

Received 11 October 1995; accepted 11 January 1996.

Jane E. Knapp and Joel M. Chandlee
*University of Rhode Island
Kingston, RI, USA*

Figure 2. Ethidium bromide-stained 1% agarose gel of *Bam*HI-restricted total genomic DNAs. Lane M, lambda *Hind*III molecular weight markers; lanes B1, B2, P1 and P2 as in Figure 1.