Benchmarks

protocol resulted in the highest fidelity sequence, with a 100% sequence accuracy over a 200-nucleotide read length for the sequence NL4-3 (Accession No. M19921) (7). In the absence of a blocking agent, dilution of the template failed to improve the clarity of the sequence obtained from either magnetic bead preparation. However, the template could be titrated further on Type A beads blocked with 1% BSA rather than on beads alone (data not shown).

The incorporation of Nonidet® P-40 (NP40) into the annealing reaction to improve the integrity of short, double-stranded sequences has been previously described (2). The incorporation of NP40 with the described blocking protocol did not affect the quality of the sequence obtained (data not shown).

This protocol demonstrates that the quality of a sequence derived from a template immobilized on streptavidin-coated magnetic beads is vastly improved by the incorporation of a simple blocking step before the annealing reaction. This modification also prevents the adsorption of sequencing reagents by a streptavidin-coated magnetic bead preparation (Type B) from which solid-phase sequencing has previously been unsuccessful.

Streptavidin-coated magnetic bead preparation Type B, in conjunction with blocking in 1% casein, has been successfully used to sequence the *env* region of DNA derived from several primary HIV-1 isolates (3).

REFERENCES

1. Adachi, A., H.E. Gendelman, S. Koenig, T. Folks, R. Willey, A. Rabson and M.A. Martin. 1986. Production of acquired immunodeficiency syndrome-associated retrovirus in human and non-human cells transfected with an infectious molecular clone. *J. Virol.* 59:284-291.
2. Bachman, B., W. Lüke and G. Hunsmann. 1990. Improvement of PCR amplified sequencing with the aid of detergents. *Nucleic Acids Res.* 18:1309.
3. Deacon, N.J., A. Tyskin, A. Solomon, K. Smith, M. Ludford-Menting, D.J. Hooker, D.A. McPhee, A.L. Greenway, et al. 1995. Genomic structure of an attenuated quasi species of HIV-1 from a blood transfusion donor and recipients. *Science* 270:988-991.
4. Hultman, T., S. Ståhl, E. Hornes and M. Uhlen. 1989. Direct solid phase sequencing of genomic and plasmid DNA using magnetic beads as solid support. *Nucleic Acids Res.* 17:4937-4946.
5. Mitchell, L.G. and C.R. Merril. 1989. Affinity generation of single-stranded DNA for dideoxy sequencing following the polymerase chain reaction. *Anal. Biochem.* 178:239-242.
6. Mitchell, L.G., R. Vaghamar and J.J. Muttigh, Jr. 1994. Biotin-enhanced sequencing of PCR products and generation of single-stranded DNA probes, p. 31-48. In M. Uhlen, B. Hornes and Ø. Olsirik (Eds.), *Advances in Biomagnetic Separation*. Eaton Publishing, Natick, MA.
7. Myer, G., B. Korber, J.A. Berzofsky, R.F. Smith and G.N. Pavlakis. 1992. *Human Retroviruses and AIDS. Theoretical Biology and Biophysics*. Los Alamos National Laboratory, Los Alamos.

This project was funded by the Commonwealth AIDS Research Grants Committee through both a project grant and in conjunction with the National Centres for HIV Virology Research and the research fund of the Macfarlane Burnet Centre for Medical Research. We thank Karen Coates-Fryer for providing the sequencing primer sp11. Address correspondence to Victoria A. Lawson, AIDS Cellular Biology Unit, Macfarlane Burnet Centre for Medical Research, P.O. Box 254, Fairfield, Victoria, 3078, Australia.

Received 30 May 1995; accepted 13 February 1996.

Victoria Alice Lawson, Dale Alan McPhee and Nicholas John Deacon
*Macfarlane Burnet Centre for Medical Research
Fairfield, Victoria, Australia*

Recovery of Unlabeled PCR Product from Polyacrylamide Gel for Sequencing

BioTechniques 21:358-362 (September 1996)

Following polymerase chain reaction (PCR) and gel analysis of the amplified product, it is often desirable to recover the amplified DNA fragments from the gel for further PCR manipulation or sequence determination. Agarose, instead of polyacrylamide gel, is routinely used to fractionate DNA fragments due to the ease with which DNA

can be recovered. However, agarose has several disadvantages. The major ones are poor resolution of closely migrating bands, less sensitivity and higher cost (1). Furthermore, agarose is known to contain substances that may be inhibitory to a number of enzymes (4), including AmpliTaq® DNA Polymerase (Perkin-Elmer, Norwalk, CT, USA), which may be used in experiments subsequent to DNA recovery. An alternative method, which allows higher resolution, greater sensitivity and easy recovery of DNA bands, is preferred.

We have modified a procedure previously described for recovering radio-labeled cDNA fragment in mRNA differential display (2). Our method was found to be very efficient in recovering unlabeled PCR-amplified DNA amenable to further manipulation.

After PCR, 5%–10% (5–10 µL) of the PCR product was heated at 65°C for 10 min and chilled on ice. After the addition of dyes, the sample was loaded onto a standard (15.5 cm × 18.5 cm) vertical 0.75-mm 3.5% polyacrylamide gel in 1× TBE, pH 8.3 (10×, 890 mM Tris base, 890 mM boric acid, 20 mM EDTA), and electrophoresed at 180 V for 1.5 h. After electrophoresis, the gel was stained in ethidium bromide solution (1 µg/mL) for 20 min, and the DNA bands were visualized under UV

Figure 1. Polyacrylamide gel electrophoresis of PCR product of hLHR. Lane S: Size marker (ϕX174 DNA digested with *Hae*III). Fragment sizes in decreasing order are: 1353 bp, 1078 bp, 872 bp, 603 bp, 310 bp, 271/281 bp, 234 bp and 194 bp, respectively. Lane P: PCR product. The size of the DNA fragments recovered using the described method are indicated along the side of the figure. Lane F1: Fragment 1, size 668 bp, recovered from P and further amplified by PCR. Lane F2: Fragment 2, size 217 bp, recovered from P and further amplified by PCR.

Benchmarks

illumination. The band to be recovered was overlaid with a piece of 3-mm filter paper cut to the size of the band. The entire gel was then covered with a large piece of 3-mm filter paper, wrapped on the other side with Saran® wrap and dried at 80°C for 1 h in a gel dryer. After drying, the gel is stuck to the paper. The band of interest is recovered sticking to the small piece of filter paper laid on the band.

There are a number of advantages of this procedure over the other recovery methods. The major advantage is the elimination of the time factor in recovering the DNA fragment for further analysis. Traditional methods, including electroelution, for recovering DNA fragment from gels require the processing of the excised gel band immediately after electrophoresis. DNA bands that are not excised from the gel are lost. On the other hand, marking of DNA bands, using overlaid paper strips as described, allows retrieval of these bands anytime, even after the entire gel has been dried down. DNA fragments in the dried gel are stable and can be stored at 4°C or recovered when desired. This is extremely useful, especially in cases when a number of bands is present and processing of all the bands are physically difficult or not desirable immediately after electrophoresis. Another advantage is the reduction of ultraviolet (UV) exposure time. UV light is damaging to eyes and exposed skin. Reducing the time of exposing laboratory personnel to UV light to the minimum is highly desirable. Overlaying of paper strips is a much faster process than excising gel bands. Therefore the procedure described reduces the amount of time required to work under UV light, and thus reduces the hazard of working with UV light. Furthermore, it is quite difficult to cut precisely the gel band corresponding to the DNA fragment, especially when this has to be done under UV illumination. On the other hand, the band can be very precisely located by trimming the filter paper to the exact size of the band in the current method. This will reduce the amount of polyacrylamide gel present, which may interfere with the recovery of the embedded DNA. This advantage is especially significant when two DNA fragments migrating closely together are to be

recovered separately. Cutting gel strips, corresponding to two closely migrating DNA bands, immediately after electrophoresis, can be extremely difficult because of the softness of the gel. Cross-contamination occurs easily under such circumstance. On the other

hand, manipulation with a dried gel is much easier.

To recover the DNA fragment, the small piece of filter paper, to which the dried gel stuck, was soaked in 100 µL of sterile deionized water for 10 min at room temperature in a 1.5-mL micro-

Figure 2. Autoradiograph of sequencing gel of fragment 1 (F1) and fragment 2 (F2). Both FIS and F1L represent sequence of fragment 1 (F1) determined by dideoxy chain termination method. The nucleotide sequence of FIS and F2 were determined using the 5'Ex8 primer (nucleotide 606–628) (5'-GGAGCTAAAGGAAAACGTACATCT-3') (3) while that of F1L was determined using the 5'Ex10 primer (nucleotide 867–891) (5'-ACAGAATTTTCACATTCCATTCT-3') (3). Lanes corresponding to the bases G, A, T and C were indicated. Deletion of exon 9 and the 5' portion of exon 11 in fragment 2 (F2) were indicated. The numbers indicate the nucleotides bordering the deletions.

centrifuge tube. The microcentrifuge tube was then placed in a boiling water bath for 15 min, and the gel/paper was pelleted by brief spinning for 2 min at 15 000× g. This procedure for releasing embedded DNA from polyacrylamide gel avoids the use of electroelution. It also allows cleaner separation of the gel from the supernatant than when fresh gel is used. This procedure has been used successfully in our laboratory to recover amplified DNA fragments ranging in size from 160 bp to 1.4 kb. Even though the yield appears to be better for smaller DNA fragments, no difference in the quality of DNA recovered has been observed.

The released DNA could be recovered by transferring the supernatant to a new microcentrifuge tube, and the DNA was precipitated by adding 10 µL of 3 M sodium acetate, 5 µL of glycogen (10 mg/mL) and 450 µL of 100% ethanol and keeping on dry ice (or at -70°C) for 30 min. To recover the precipitated DNA, the tube was spun at 15 000× g in a microcentrifuge for 10 min at 4°C. The DNA pellet was rinsed with 200 µL of prechilled 100% ethanol and then air-dried. The DNA could be resuspended in 10 µL of distilled deionized water or TE buffer (10 mM Tris-HCl, pH 7.5, 1 mM EDTA). As little as one-tenth of the DNA recovered can be used as template to yield satisfactory results in subsequent PCR or asymmetric PCR (APCR). If multiple amplified DNA fragments were observed after another round of PCR using the recovered band as a template, the procedure could be repeated until a single DNA fragment was obtained.

This method is extremely useful in cases where an individual displays genetic heterogeneity due to a possible deletion or insertion. Here, multiple bands will be observed following PCR amplification using genomic DNA or reverse transcription (RT)-PCR using total RNA as a template. To characterize the multiple amplified fragments, each DNA band must be recovered and analyzed. Using the method described above, the amplified DNA fragments can be recovered. APCR can be performed to generate single-stranded DNA (1), and the deletion/insertion of the interested gene is verified by nucleotide sequencing.

In our study of the splicing variations of the human luteinizing hormone receptor (hLHR) gene, cDNAs of the hLHR encoding exons 8 to 10 and portion of exon 11 (3) were prepared by RT-PCR using total RNA from a normal human adult testicle obtained from the National Disease Research Interchange, Philadelphia, PA, USA. Analysis of the PCR product using polyacrylamide gel electrophoresis revealed the presence of three prominent bands, one of the appropriate size (668 bp) and two smaller bands (Figure 1, lane P). Two of these bands, the 668-bp and the 217-bp bands, were selected for further analysis. The bands were recovered from the gel using the method described above. Analysis of the recovered DNA fragment using PCR and gel electrophoresis showed that they all contained a single species of DNA fragment (Figure 1, lanes F1 and F2). The recovered DNA bands were then used as templates for APCR (1), and the nucleotide sequence of the single-stranded DNAs generated was determined by dideoxy sequencing using [³²P]dATP (5). Parts of the sequencing results are shown in Figure 2. Sequences F1L and F1S represented nucleotide sequence of F1 (668 bp) determined using two different sequencing primers in order to reveal the sequence at different portions of the fragment. Comparison of the nucleotide sequence of F1 with the published sequence of hLHR cDNA showed that F1 was the normally spliced hLHR containing exon 8 to exon 11. On the other hand, the entire exon 9 and the 5' 266 bp of exon 11 (from nucleotide 948–1213) were deleted in F2 (Figure 2). Occurrence of the splice variants of the LHR had been demonstrated in gonads of rats, mice and macaque (6). This result demonstrates that splice variants of the hLHR also occur in normal adult human male gonads.

The procedure described in this report allows the subsequent analysis of amplified DNA fragments, which are present in small quantities even after PCR. The recovered DNA is free of possible inhibitory substances and allows nucleotide sequence determination after APCR. This enhances our capability of analyzing multiple DNA fragments generated by PCR.

REFERENCES

1. Ausubel, F.M., R. Brent, R.E. Kingston, D.D. Moore, J.G. Seidman, J.A. Smith and K. Struhl. 1994. Current Protocols in Molecular Biology. John Wiley & Sons, New York.
2. GenHunter Corporation. 1994. RNA Image™ mRNA Differential Display System Instruction Manual. GenHunter Corporation, Brookline, MA.
3. Minegishi, T., K. Nakamura, Y. Takakura, K. Miyamoto, Y. Hasegawa, Y. Ibuki and M. Igarashi. 1990. Cloning and sequencing of human LH/hCG receptor cDNA. Biochem. Biophys. Res. Commun. 172:1049-1054.
4. Sambrook, J., E.F. Fritsch and T. Maniatis. 1989. Molecular Cloning. A Laboratory Manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
5. Sanger, F., S. Nicklen and A.R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. Proc. Natl. Acad. Sci. USA 74:5463-5467.
6. Segaloff, D.L. and M. Ascoli. 1993. The lutropin/choriogonadotropin receptor...4 years later. Endocrine Rev. 14:324-346.

S.-M.W. was supported in part by NIH Grant P01 HD24633, and W.-Y.C. was supported in part by NIH Grants R01 HD31553 and P01 HD24633. Address correspondence to Wai-Yee Chan, Dept. of Pediatrics, Georgetown University Children's Medical Center, 3800 Reservoir Road, NW, Washington, DC 20007-2196, USA. Internet: wchan02@gumedlib.dml.georgetown.edu

Received 6 November 1995; accepted 13 February 1996.

Shao-Ming Wu, Le Ann Blomberg and Wai-Yee Chan
*Georgetown University
Children's Medical Center
Washington, DC, USA*

PCR-RFLP Genotyping of Murine MHC Haplotypes

BioTechniques 21:362-368 (September 1996)

The major histocompatibility complex (MHC) has a central role in the restriction of responses during graft acceptance and rejection, antibody production and antigen presentation (5). In the past, transplantation studies have purposefully generated mice of mixed H-2 (MHC) genotype by mating