ng thrombin (Sigma Chemical, St. Louis, MO, USA) in a total volume of 20 μ L for 1 h at 37°C. The buffer used was 50 mM sodium citrate, pH 6.5, 150 mM NaCl. Samples incubated with or without thrombin were analyzed by electrophoresis on a 15% polyacrylamide gel. After electrophoresis, a Western blot was performed to detect the sizes and thrombin susceptibilities of each GlpR (thrombin cleavage was also detected on the Coomassie blue-stained SDS gel). As shown in Figure 2, GlpR encoded by pGZ117 was cleaved by thrombin to yield a product that co-migrated with authentic GlpR. The GlpR variants encoded by pGZ118 or pGZ119 were slightly larger than authentic GlpR and were not cleaved by thrombin. Since the thrombin site encoded by pGZ117 is fused in the same reading frame predicted by the nucleotide sequence, the results prove that the reading frame predicted for *glpR* is correct as deduced from its DNA sequence and that cysteine is in fact the C-terminal amino acid residue of the *glp* repressor.

This method has proven to be an effective tool for verification of the translational reading frame predicted by DNA sequence analysis. The approach is adaptable to any gene. Other restriction enzyme pairs, such as *NarI* and *KasI*, can be used to create translational frameshifts. In most cases, the construct with the correct reading frame will yield a gene product that can be purified in a single step using immobilized metal affinity chromatography (2).

REFERENCES

1. Carty, S.M., A. Colbeau, P.M. Vignais and T.J. Larson. 1994. Identification of the *rpmF-plsX-fabH* genes of *Rhodobacter capsulatus*. FEMS Microbiol. Lett. 118:227-232.
2. Kasher, M.S., M. Wakulchick, J.A. Cook and M.C. Smith. 1993. One-step purification of recombinant human papillomavirus type 16 E7 oncoprotein and its binding to the retinoblastoma gene product. BioTechniques 14: 630-641.
3. Larson, T.J., S. Ye, D.L. Weissenborn, H.J. Hoffmann and H. Schweizer. 1987. Purification and characterization of the *glp* repressor of *Escherichia coli* K-12. J. Biol. Chem. 262:15869-15874.
4. Tabor, S. and C.C. Richardson. 1985. A bacteriophage T7 RNA polymerase/promoter system for controlled exclusive expression of specific genes. Proc. Natl. Acad. Sci. USA 82:1074-1078.

We thank Ali T. van Loo-Bhattacharya for skillful technical assistance. This work was supported by National Science Foundation Grant MCB-9118757. Address correspondence to Timothy J. Larson, Department of Biochemistry, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061-0308, USA. Internet: tilarson@vt.edu

Received 9 January 1996; accepted 12 April 1996.

Gang Zeng and Timothy J. Larson

*Virginia Polytechnic Institute
and State University
Blacksburg, VA, USA*

Preparation of Miniprep-DNA for Automated Non-radioactive Sequencing

BioTechniques 21:824-826 (November 1996)

Numerous rapid miniprep procedures for isolation of plasmid DNA have been described during the last years, mostly based on the standard alkaline lysis protocol (1,3,4,6). However, in our hands none of these techniques that we have tested so far resulted in DNA pure enough to be used for automated sequencing. Most of the preparations did contain contaminations like bacterial DNA, RNA or proteins, which inhibited automated sequencing of the DNA using an ALF DNA Sequencer™ (Pharmacia Biotech, Freiburg, Germany). Furthermore, commercially available kits were too expensive to be used in our project of differential display of myocardial RNA (2), where large numbers of cloned PCR products had to be screened by restriction digest and automated sequencing. Therefore, we had to modify the standard alkaline lysis protocol in such a way that the technique was accurate and reliable for the isolation of plasmid DNA suitable for automated sequencing. The procedure is described in Table 1.

In this procedure, RNAs are removed by treatment with RNases like RNase A or RNase-It™ (Stratagene, Heidelberg, Germany). The subsequent phenol-chloroform-isoamyl alcohol extraction removes all residual proteins including the RNases, whereas all other inhibiting contaminations seem to be excluded from the preparation by isopropanol precipitation. The average plasmid yield of this procedure is 15 μ g for high-copy number plasmids like pGEM® (Promega, Madison, WI, USA) or pBluescript® (Stratagene) at a concentration of about 0.5 μ g/ μ L and less for low-copy-number plasmids like the pBR series. The plasmid DNA is mostly supercoiled and can be used for further applications like restriction analysis (Figure 1), isolation of inserts and labeling for Northern or Southern blot analysis. Most important, the DNA isolated with this procedure is suitable for automated nonradioactive sequencing.

Benchmarks

Table 1. Protocol

1. Grow an overnight culture (15 h maximum) of the bacterial clone of interest in 3 mL LB medium supplemented with the appropriate antibiotics at 37°C with shaking at 250 rpm in a tube of 10 mL or more.
2. Pour about 1.5 mL in a microcentrifuge tube (store the rest at 4°C for further use) and centrifuge bacteria for 2 min at top speed in a microcentrifuge at room temperature.
3. Remove the supernatant completely using pipet tips and dry it by inverting for 5 min.
4. Resuspend the pellet in 150 µL of standard ice-cold GTE-buffer (50 mM glucose, 25 mM Tris-HCl, pH 8.0, 10 mM EDTA, pH 8.0, optional: lysozyme at 2 mg/mL) by pipetting up and down. Do not vortex mix!
5. Add 200 µL of freshly prepared lysis buffer (0.2 N NaOH, 1% sodium dodecyl sulfate) and mix by inverting tubes carefully. Do not vortex mix! Leave on ice for 5 min.
6. Add 150 µL of 3 M potassium acetate solution, prepared as described previously (3), mix by inverting tubes several times and centrifuge at top speed in a microcentrifuge for 10 min at room temperature.
7. Transfer supernatant in a new tube, add 0.5 volume of isopropanol (room temperature) and spin down for 10 min at top speed in a microcentrifuge at room temperature. Be careful not to disturb the pellet.
8. Discard supernatant, invert tube and dry pellet for 10–15 min. Do not let it dry down completely, otherwise the pellet will be difficult to resuspend.
9. Redissolve the pellet in 500 µL of STE (10 mM NaCl, 10 mM Tris-HCl, pH 8.0, 1 mM EDTA, pH 8.0) as described previously (5), add 10 µL of RNase A (10 mg/mL) or 6 µL of RNase-It (Stratagene) and incubate at 37°C for 30 min.
10. Extract twice with an equal volume of phenol-chloroform-isoamyl alcohol (25:24:1), then add 1/10 vol 3 M sodium acetate, pH 5.2, and 0.5 vol isopropanol at room temperature, invert tube several times and centrifuge the plasmid DNA at top speed in a microcentrifuge for 5 min at room temperature.
11. Finally, wash the DNA twice with 1 mL of 70% ethanol, dry under a vacuum, redissolve in 30 µL water (TE buffer seems to inhibit the sequencing reactions) and store at 4°C until further use.

ing using the ALF DNA Sequencer. We routinely use 8 µL of the DNA solution in standard sequencing reactions with 2 µL of fluorescein-dATP-labeled primers, leaving enough DNA available for

other applications. To facilitate binding of the primers to their target sequences on the template, the annealing temperature is lowered to 25°C and the annealing time increased up to 30 min. This

Figure 1. Isolation of miniprep plasmid DNA. Plasmid DNA was isolated as described in Table 1, digested with *Bss*HII and separated on a 0.7% agarose gel. Lanes 1, 3, 5, 7, 9, 11, 13 and 15 represent uncut miniprep DNA from different clones, whereas lanes 2, 4, 6, 8, 10, 12, 14 and 16 show the DNA digested with *Bss*HII. P and λ represent Φ X174/*Hae*III and λ DNA/*Hind*III markers, respectively.

change in the standard procedure worked for all primers tested so far, regardless of their melting temperature (T_m). Furthermore, it simplified the sequencing procedure when using a large number of primers with different T_m . All other reactions were performed according to the Autoread™ Sequencing Kit (Pharmacia Biotech). Using these minor modifications, sequences up to 500 bp in length with less than 2% sequence ambiguities could be read. The average length was about 290 bp with 2% sequence ambiguities or less. Therefore, our miniprep procedure offers an economical or time-saving alternative to commercial miniprep kits or maxi preps, respectively. This is especially suitable for screening a large number of plasmid clones derived from procedures like differential display or subtractive hybridization.

REFERENCES

1. Birnboim, H.C. and J. Doly. 1979. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res.* 7:1513-1523.
2. Liang, P. and A.P. Pardee. 1992. Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction. *Science* 257:967-971.
3. Sambrook, J., E.F. Fritsch and T. Maniatis. 1989. *Molecular Cloning: A Laboratory Manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
4. Tiesman, J. and A. Rizzino. 1991. A rapid and reliable method for the purification of high-quality plasmid DNA for double-stranded sequencing. *BioTechniques* 10:319-328.
5. Wang, L.-F., R. Voysey and M. Yu. 1994. Simplified large-scale alkaline lysis preparation of plasmid DNA with minimal use of phenol. *BioTechniques* 17:26-28.
6. Xiang, C., H. Wang, P. Shiel, P. Berger and D.J. Guerra. 1994. A modified alkaline lysis miniprep protocol using a single microcentrifuge tube. *BioTechniques* 17:30-32.

Address correspondence to René Zimmermann, Max-Planck-Institute, Department of Experimental Cardiology, Benekestrasse 2, D-61231 Bad Nauheim, Germany. Internet: rzimmer@kerckhoff.mpg.de

Received 26 December 1995; accepted 9 April 1996.

René Zimmermann, Claudia Ullmann and Wolfgang Schaper
*Max-Planck-Institute
 Bad Nauheim, Germany*