REFERENCES

1. Atcheson, C.L., B. DiDomenico, S. Frackman, R.E. Esposito and R.T. Elder. 1987. Isolation, DNA sequence, and regulation of a meiosis-specific eukaryotic recombination gene. *Proc. Natl. Acad. Sci. USA* 84:8035-8039.
2. Cha, J., W. Bishai and S. Chandrasegaran. 1993. New vectors for direct cloning of PCR products [published erratum appears in *Gene* 1994; 141:149]. *Gene* 136:369-370.
3. Clark, J.M. 1988. Novel non-templated nucleotide addition reactions catalyzed by prokaryotic and eukaryotic DNA polymerases. *Nucleic Acids Res.* 16:9677-9686.
4. Harrison, J., P.L. Molloy and S.J. Clark. 1994. Direct cloning of polymerase chain reaction products in an *XcmI* T-vector. *Anal. Biochem.* 216:235-236.
5. Ichihara Y. and Y. Kurosawa. 1993. Construction of new T-vectors for direct cloning of PCR products. *Gene* 130:153-154.
6. Kovacic, D., J.H. Kwak and B. Weisblum. 1991. General method for direct cloning of DNA fragments generated by the polymerase chain reaction. *Nucleic Acids Res.* 19:4560.
7. Mead, D.A., N.K. Pey, C. Herrstadt, R.A. Marciel and L.M. Smith. 1991. A universal method for the direct cloning of PCR amplified nucleic acid. *Bio/Technology* 9:657-663.
8. Mertz, L.M. and A. Rashtchian. 1994. PCR radioactive labeling system: a rapid method for synthesis of high specific activity DNA probes. *Focus* 16:45-48.
9. Testori, A., I. Listowsky and P. Sollitti. 1994. Direct cloning of unmodified PCR products by exploiting an engineered restriction site. *Gene* 143:151-152.

The authors wish to thank Margaret Malik for reviewing this protocol and the manuscript. This work was supported by NIH Grant P30-HD-27748 (B.C.S. and J.P.). Address correspondence to Brian C. Schutte, Department of Pediatrics, MRC 216, University of Iowa, Iowa City, IA 52242, USA. Internet: brian-schutte@uiowa.edu

Received 11 March 1996; accepted 21 June 1996.

Brian C. Schutte, Koustubh Ranade¹, Jonathan Pruessner and Nic Dracopoli²

University of Iowa
Iowa City, IA

¹Stanford University School
of Medicine
Stanford, CA

²Sequana Therapeutics
La Jolla, CA, USA

Universal Primers for Sequencing DNA Restriction Fragments

BioTechniques 22:44-46 (January 1997)

A common strategy for large-scale DNA sequencing is shotgun cloning. In this approach, the cloned DNA is fragmented into small pieces, which are first size-fractionated and then sub-cloned into a sequencing vector. From the collection of clones obtained, a random sample is chosen for sequencing. The nucleotide sequence information from individual sequence reads is then fed into a computer, which assembles sequence overlaps between contiguous clones (contigs), leading to the reconstruction of the sequence of the original cloned DNA. While this approach initially provides rapid sequence information, the utilization of more directed approaches is often needed for efficient gap closures and verification of contig assembly (1). In the present communication, we present a method for creating universal primers for sequencing restriction fragments, which can serve as an alternative directed approach to facilitate gap closure and assembly ver-

ification in large-scale DNA sequencing projects.

The method, as outlined in Figure 1, exploits the ability of the enzyme terminal deoxynucleotidyl transferase to add a string of T-residues at the 3' end of a restriction fragment. The T-tails, along with the neighboring remnant of the restriction enzyme recognition site, can then be used to direct the synthesis of specific primers for the sequencing of the modified restriction fragment.

To test this approach, we have used a unique *PstI*-*SphI* DNA fragment derived from plasmid pBR322. Both *PstI* and *SphI* enzymes produce 3' overhangs, which are the best substrates for the enzyme terminal deoxynucleotidyl transferase. Briefly, a 1315-bp fragment, produced after digestion of pBR322 DNA with both *PstI* and *SphI*, was extracted from an agarose gel using the Wizard™ DNA Clean-Up System (Promega, Madison, WI, USA). T-tailing of the 3' overhangs, present in the DNA fragment, was performed using terminal deoxynucleotidyl transferase (Promega) in a buffer supplied by Promega containing 1 pmol of DNA, 20 pmol dTTP and 2 pmol ddTTP. The tailed DNA product was then purified with the Wizard System. For the sequencing of the tailed restric-

Figure 1. Steps involved in the modification of restriction fragments for sequencing. The strategy used to sequence a *PstI*-*SphI* fragment derived from plasmid pBR322 is shown. UPstI and USphI denote universal primers for *PstI* and *SphI* restriction sites, respectively.

Benchmarks

tion fragment, universal primers UPstI (5'-AAAAAAAAAATGCAG-3') and USphI (5'-AAAAAAAAAACATGC-3') were synthesized. These primers contain ten adenine residues that have been added at the 5' end of the restriction remainder of enzyme recognition sequences for *Pst*I and *Sph*I, respectively. DNA sequencing of the modified restriction fragment was performed by cycle sequencing, using an *fmol*[®] DNA Sequencing System (Promega) and [γ -³²P]ATP end-labeled UPstI and USphI as primers. Figure 2 shows that both UPstI and USphI primers yield readable sequence information. More importantly, in each case, the priming site corresponds to the proper end of the restriction fragment. Our results therefore indicate that the approach described above to sequence restriction fragments is a feasible one.

Although we have so far only tested this methodology with restriction enzymes producing 3' overhangs, we assume that restriction fragments produced with blunt end cutters should also be amenable to this approach. Restriction fragments with 5' overhangs, on the other hand, would need to be filled with a Klenow fragment before tailing.

As indicated earlier, the sequencing of restriction fragments could serve as an alternative direct approach for re-

solving gap closure problems and in sequence assembly verification in DNA sequencing projects using a shotgun cloning approach. Our methodology could also be useful for the determination of sequence information from DNA regions that are refractive to cloning ("poison sequences") and for the amplification of these regions using polymerase chain reaction (PCR). Also, the simultaneous gathering of sequence information from distinct regions of a cloned DNA could facilitate the construction of probes from any particular region of this clone. Finally, we wish to emphasize that the primers required for sequencing a restriction fragment by our approach are universal in nature, since they are not dependent on the origin of the template DNA. Thus, any DNA fragment that has been generated with two particular restriction enzymes can, in principle, be sequenced with the universal primers targeted to the two restriction enzymes used to create the DNA fragment.

REFERENCE

1. Bodenteich, A., S. Chisoe, Y.F. Wang and B.A. Roe. 1994. p. 42-50. In M.D. Adams, C. Fields and J.C. Venter (Eds.), *Automated DNA Sequence and Analysis*. Academic Press, New York.

We thank G. Chaudhuri for critical comments. This work was supported in part by grants from the National Institutes of Health. Address correspondence to Manuel S. Valenzuela, Division of Biomedical Sciences, Meharry Medical College, 1005 D.B. Todd Blvd., Nashville, TN 37208, USA. Internet: valenz44@ccvax.mmc.edu

Received 15 March 1996; accepted 23 May 1996.

Yeeru J. Chen and Manuel S. Valenzuela
*Meharry Medical College
Nashville, TN, USA*

Figure 2. Nucleotide sequence of a *Pst*I-*Sph*I DNA fragment obtained with universal primers either to the *Pst*I (UPstI) or to the *Sph*I (USphI) end of the restriction fragment.