Benchmarks

9. **New England Biolabs**. 1996. Product Catalog, p. 214. Beverly, MA.
10. **Sambrook, J., E.F. Frisch and T. Maniatis**. 1989. Molecular Cloning: A Laboratory Manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
11. **Yanisch-Perron, C., J. Vieira and J. Messing**. 1985. Improved M13 phage cloning vectors and host strains: nucleotide sequence of the M13mp18 and pUC19 vectors. *Gene* 33:103-119.

This work was supported by grants from the National Eye Institute and Research to Prevent Blindness to A.S. Address correspondence to Abraham Spector, Department of Ophthalmology, Columbia University, 630 West 168th Street, New York, NY 10032, USA. Internet: as42@columbia.edu

Received 27 September 1996; accepted 19 November 1996.

Yinqing Yang and Abraham Spector
*Columbia University
New York, NY, USA*

PCR Strategy to Obtain the 5' and 3' Flanking Sequences of Partial Clones from λ ZAP cDNA Libraries

BioTechniques 22:1038-1044 (June 1997)

Determining the complete coding, and sometimes, the 5' and/or 3' untranslated regions from the available partial sequences is a common problem in molecular cloning. These partial clones can be derived either from polymerase chain reaction (PCR) amplifica-

tion of reverse-transcribed mRNA or genomic DNA using degenerate primers, or from screening cDNA libraries. Several approaches to obtain full-length clones have been developed. Screening a cDNA library is fairly dependable, but it is time-consuming, labor-intensive and costly. Rapid amplification of cDNA ends (RACE) (3) is a widely used, PCR-based method designed to clone full-length cDNAs. RACE is useful in analyzing a larger primary population of mRNAs than is possible in library screening, but it is technically demanding and can be limited by the availability of RNA (i.e., human RNA). An alternative PCR

Figure 1. Experimental procedure as detailed in Table 1.

Benchmarks

approach to cloning full-length sequences was described independently by Huang (4) and Zeiner and Gehring (10). This method involves reverse transcription of mRNA, second-strand synthesis, cDNA circularization and inverse PCR, in which the 3' ends of the two gene-specific primers from the known sequence are oriented away from each other (1,2,5,9). We describe a new strategy to clone the 5' and 3' flanking sequences of any cDNA when only a portion of the sequence is known. This method combines the utility of inverse PCR with the convenience of in vivo excision of λ ZAP cDNA libraries (which are commercially available or can be obtained from many laboratories) and comprises the following four steps: (i) in vivo excision of a λ ZAP cDNA library containing the gene of interest into a pBluescript[®] phagemid (Stratagene, La Jolla, CA, USA) cDNA library; (ii) inverse PCR of the excised library with two gene-specific primers to amplify fragments containing 5' and 3' flanking sequences and pBluescript vector sequence; (iii) a second round of standard PCR using a portion of the first amplification as template with nested 5' or 3' gene-specific and pBluescript vector primers; and (iv) subcloning and sequencing of the resulting fragments (Figure 1). Table 1 details the protocol for each step of the method.

We have tested this strategy on an excised turkey-blood cDNA library (8) to amplify the 5' and 3' flanking sequences of two genes, the P2Y₁ purinergic receptor (P2Y-R) and the G protein α_s subunit (G_{α_s}). Both of these genes were cloned previously from this library by traditional screening methods (Q. Li, C. Herold, S. Short and R.A. Nicholas, unpublished results). The sizes of the cDNA inserts, 3.6 kb for P2Y-R and 1.8 kb for G_{α_s} , were derived from single full-length isolates.

Inverse PCR of the excised library using the P2Y-R primers Y-1 and Y-2 would be predicted to amplify an approximately 6-kb fragment (Figure 2A). Although no specific band was observed in the expected region in an agarose gel stained with ethidium bromide (Figure 2B, left panel), a band of the expected size was observed when the fragments were blotted onto nylon

Table 1. Protocol for Cloning cDNA When Only a Portion of the Sequence is Known

1. In vivo mass excision of a λ ZAP cDNA library is carried out as described in the manual supplied with the ZAP[®]-cDNA Synthesis Kit (Stratagene). Excision of 1×10^9 plaque-forming units (pfu) of a turkey-blood cDNA library yielded 730 μ g of plasmid library, enough DNA for hundreds of amplifications.
2. Inverse PCR is used to amplify from the excised library the 5' and 3' flanking sequences of the gene of interest along with pBluescript vector sequence (2.96 kb). Because the efficiency of PCR with *Taq* DNA polymerase decreases markedly when amplifying fragments larger than 4 kb, use *Tth* DNA Polymerase, XL (Perkin-Elmer, Norwalk, CT, USA) in inverse PCR with an aliquot of the excised library (0.1–3 μ g of DNA) and the appropriate gene-specific primers (Figure 1).
3. To obtain sufficient amounts of 5' and 3' flanking sequences that can be more easily subcloned, carry out a second round of standard PCR with *Taq* DNA Polymerase (Life Technologies, Gaithersburg, MD, USA) using 1% of the inverse PCR products (from step 2) as template with pBluescript vector primers and nested gene-specific primers. Use a standard PCR protocol with glycerol and dimethyl sulfoxide (DMSO) to improve amplification of GC-rich templates (7).
4. Use standard protocols (6) to subclone and sequence the amplified products obtained from the second round of PCR. Once the sequences of the flanking regions are known, primers that will amplify the full-length cDNA from the excised library can be designed. To avoid errors introduced during amplification, a proofreading DNA polymerase such as *Pfu* (Stratagene) or Vent[™] (New England Biolabs, Beverly, MA, USA) is recommended.

membranes and probed with a ³²P-labeled P2Y-R cDNA insert (Figure 2B, right panel). The low yield of the P2Y-R sequence in inverse PCR may have been due to a decreased efficiency of amplification, since low yields of amplification were also observed in inverse PCR when the isolated cDNA clone was used as a template (data not shown). In the second round of PCR, 1% of the inverse PCR products was used as template with the nested primer Y-4 and reverse vector primer to amplify the 5' flanking sequence or with the nested primer Y-3 and T7 vector primer to amplify the 3' flanking sequence. The predicted sizes of the amplified 5' and 3' flanking regions from the second round of PCR were approximately 500 bp and 2 kb, respectively. As shown in Figure 2B, significant amounts of P2Y-R-specific fragments close to their predicted sizes were obtained. The fragments amplified from the 3' flanking region were much less heterogeneous than those amplified from the 5' flanking region. We suspect that the heterogeneity in the 5' flanking region is due to the existence in the excised library of

a population of clones that differ in size of their 5' ends. This is consistent with the results from library screening, which identified several P2Y-R cDNA clones that varied in the length of their 5' ends. Homogeneity of the amplified 3' flanking region would be expected because reverse transcription of turkey-blood mRNA was primed with oligo(dT) during library construction.

A similar set of experiments was carried out with turkey G_{α_s} cDNA as another test sequence (Figure 2A). Based on the size of the G_{α_s} clone isolated by library screening, inverse PCR of G_{α_s} with G-1 and G-2 would be expected to amplify an approximately 4-kb fragment. In contrast to the results with P2Y-R, inverse PCR with G-1 and G-2 primers resulted in highly efficient amplification of the G_{α_s} cDNA (Figure 2C). The large quantities of amplified product following inverse PCR may be the result of a higher representation of the G_{α_s} cDNA in the excised library and/or to a higher efficiency of amplification. The second round of PCR was carried out using 1% of the inverse PCR products as template with G-4 and

Benchmarks

reverse primers, or G-3 and T7 primers, which would be expected to amplify fragments 450 and 500 bp in length, respectively. Following the second round of PCR, specific bands of the expected size were observed for both the 5' and 3' flanking sequences (Figure 2C). In contrast to the P2Y-R sequence, amplification of the 5' flanking sequences of $G_{\alpha s}$ resulted in a much more homogeneous population of fragments, although some hybridization was observed both above and below the major hybridizing band. As seen with the P2Y-R, amplification of the 3' flanking sequence resulted in a homogeneous band of the expected size.

We originally attempted to amplify the flanking sequences of the P2Y-R and $G_{\alpha s}$ cDNAs from the same mass-excised cDNA library as above by two rounds of standard PCR. The first round of PCR contained the appropriate gene-specific and pBluescript vector primers, and the second round contained 1% of the first-round amplification products as template and the appropriate nested gene-specific and pBluescript vector primers. No specific amplification products were obtained (data not shown). We speculate that this approach does not work, because the vector primers anneal to all of the pBluescript clones in the template and are depleted in later rounds of PCR. Specific products were also not obtained after increasing the amount of vector primer in the amplification mixture or carrying out 40 rounds of amplification using only the gene-specific primer before the two rounds of PCR. Thus, it is crucial in this method to carry out an initial round of inverse PCR with two gene-specific primers, because the gene of interest becomes the dominant DNA template for the second round of standard PCR with a nested gene-specific primer and pBluescript vector primer. This permits specific amplification of the flanking sequences of the gene of interest.

In conclusion, we have developed a PCR strategy to obtain the flanking sequences of partial clones from a λ or plasmid cDNA library. Although we describe in this paper the method for inverse PCR from mass-excised λ ZAP cDNA libraries, any phage cDNA library that can be excised *in vivo* (such

as the λ ZIPLOX™ system; Life Technologies) or plasmid cDNA library can be used as a template for inverse PCR. There are several advantages to this method over previously used approaches. The method is rapid since *in vivo* excision requires 3–4 days, and the two rounds of PCR require an additional 2 days. Once a library is mass-excised, it is stable for years and can be used as template in inverse PCR to amplify any genes present in the excised library.

Some limitations to this approach also apply. The cDNA library must contain appropriate full-length clones, and the clones must be amplified equally during mass excision. Furthermore, some genes may be difficult to amplify (i.e., GC-rich regions). However, given these limitations, this method represents a straightforward and simple approach to determining the 5' and 3' flanking sequences of partial clones from mass-excised or plasmid cDNA libraries.

Figure 2. Amplification of the flanking sequences of turkey P2Y-R and $G_{\alpha s}$ from a turkey-blood cDNA library. (A) Diagram showing the predicted products after each round of PCR. (B) Aliquots of the amplification products from the two rounds of PCR with P2Y-R- and vector-specific primers were separated on a 1% agarose gel and either stained with ethidium bromide (left panel) or transferred to a nylon membrane and probed with the 32 P-labeled turkey P2Y-R gene (right panel). (C) Aliquots of the amplification products from PCR with $G_{\alpha s}$ - and vector-specific primers were treated as described in B, except that the blot was probed with the 32 P-labeled turkey $G_{\alpha s}$ gene. The blots were hybridized in 50% formamide, 6× standard saline citrate (SSC), 0.5% sodium dodecyl sulfate (SDS) at 42°C overnight, washed with 0.1× SSC, 0.1% SDS at 70°C for 2 h and exposed to film overnight. Lane 1 of B and C: amplification products from inverse PCR using the excised turkey-blood pBluescript cDNA library (1.5 μ g) as template with either Y-1 and Y-2 primers (B) or G-1 and G-2 primers (C); lane 2 of B and C: amplification products from the second round of PCR using 1% of the inverse PCR products as template with reverse primer and either the Y-4 (B) or G-4 (C) primer; lane 3 of B and C: same as lane 2, except with the T7 primer and either the Y-3 (B) or G3 (C) primer. MW: DNA molecular-weight markers (size in kb).

Benchmarks

REFERENCES

1. **Chowers, Y., W. Holtmerier, E. Morzycka-Wroblewska and M.F. Kagnoff.** 1995. Inverse PCR amplification of rare T cell receptor delta messages from mucosal biopsy specimens. *J. Immunol. Methods* 179:261-263.
2. **Dalby, B., A.J. Pereira and L.S.B. Goldstein.** 1995. An inverse PCR screen for the detection of P element insertions in cloned genomic intervals in *Drosophila melanogaster*. *Genetics* 139:757-766.
3. **Frohman, M.A., M.K. Dush and G.R. Martin.** 1988. Rapid production of full-length cDNAs from rare transcripts: Amplification using a single gene-specific oligonucleotide primer. *Proc. Natl. Acad. Sci. USA* 85:8998-9002.
4. **Huang, S.H.** 1994. Inverse polymerase chain reaction—an efficient approach to cloning cDNA ends. *Mol. Biotech.* 2:15-22.
5. **Ochman, H., A.S. Gerber and D.L. Hartl.** 1988. Genetic applications of an inverse polymerase reaction. *Genetics* 120:621-623.
6. **Sambrook, J., E.F. Fritsch and T. Maniatis.** 1989. *Molecular Cloning: A Laboratory Manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
7. **Varadaraj, K. and D.M. Skinner.** 1994. Denaturants or cosolvents improve the specificity of PCR amplification of a G+C-rich DNA using genetically engineered DNA polymerases. *Gene* 140:1-5.
8. **Waldo, G.L., A. Paterson, J.L. Boyer, R.A. Nicholas and T.K. Harden.** 1996. Molecular cloning, expression, and regulatory activity of G α_{11} - and $\beta\gamma$ -subunit-stimulated PLC- β from avian erythrocytes. *Biochem. J.* 316:559-568.
9. **Wetmur, J.G., D.M. Wong, B. Ortiz, J. Tong, F. Reichert and D.H. Gelfand.** 1994. Cloning, sequencing, and expression of RecA protein from three distantly related thermophilic eubacteria. *J. Biol. Chem.* 269:25928-25935.
10. **Zeiner, M. and U. Gehring.** 1994. Cloning of 5' cDNA regions by inverse PCR. *BioTechniques* 17:1050-1054.

We thank T. Kendall Harden, José Luis Boyer, Patricia A. Ropp, Joel B. Schachter, Fidel Del Toro and Christopher L. Herold for comments and suggestions, and Gary Waldo for providing the turkey-blood cDNA library. This work was supported by Grant-in-Aid No. 9301067 from the American

Heart Association and USPHS Grant No. GM38213. R.A.N. is an Established Investigator of the American Heart Association. Address correspondence to Robert A. Nicholas, Department of Pharmacology, CB# 7365 FLOB, School of Medicine, University of North Carolina, Chapel Hill, NC 27599-7365, USA. Internet: nicholas@med.unc.edu

Received 18 June 1996; accepted 4 December 1996.

Qing Li and Robert A. Nicholas
*University of North Carolina at Chapel Hill
Chapel Hill, NC, USA*