Internet On-Ramp

Searching ESTs

Stuart M. Brown, NYU Medical Center, New York, NY, USA. Internet: browns02@mccr6.med.nyu.edu

A top priority in the multidisciplinary effort to sequence the human genome is the identification of all of the genes. Since only 2%-3% of human genomic DNA actually encodes proteins, it makes sense to focus sequencing efforts on these "expressed" regions first. This is the basis for a number of expressed sequence tag (EST) projects.

ESTs are short (200–500 bp) "single-pass" sequencing reads taken from both the 3' and 5' ends of cDNA clones. The sequence data are then fed directly from automated sequencers into the data banks with no error correction and only automatic annotation. The EST sequences are not complete genes nor complete cDNAs, but are merely gene "Tags." GenBank® contains over 1.4 million of these EST sequences. The EST section (dbEST) now constitutes more than half of GenBank and is growing much faster than the other sections. It is likely that the vast amount of EST data contains at least part of virtually all expressed human genes. Exceptions most likely include genes expressed in tiny quantities, in rare specialized cell types or at restricted developmental periods. Various EST sequencing projects have addressed this issue by sequencing ESTs from a variety of tissues and developmental stages, but it is impossible to be absolutely comprehensive. Comparisons of EST data from various tissues can be used as a measure of tissue-specific expression patterns. There are some genes such as histones and non-coding RNA genes that are underrepresented in the EST databanks because their RNA transcripts lack poly(A) tails.

EST sequences are short and contain many errors, particularly frameshifts, which make the identification of open reading frames (protein coding regions) very difficult. The dataset is huge and highly redundant, containing thousands of copies

of many common genes. Yet the elusive value of ESTs is clear: every gene is probably in there somewhere. Searches of EST databanks are an essential component of the search for new disease genes.

The Basic Local Alignment Search Tool (BLAST) similarity search engine, at the Web site of the National Center for Biotechnology Information (NCBI; <http://www.ncbi.nlm.nih.gov/BLAST/>), can be used to search for EST sequences that are similar to a specific query sequence. Since EST sequences are likely to contain many incorrect bases and insertion/deletions, it is best to use the new Gapped BLAST (BLAST Version 2.0) tool (<http://www.ncbi.nlm.nih.gov/cgi-bin/BLAST/nph-newblast?Jform=0>). Also, since protein-protein comparisons are much more sensitive than DNA-DNA comparisons, use TBLASTN for protein query sequences (compares your protein to 6-frame translated ESTs) and TBLASTX for DNA query sequences (translates your query in six frames and compares it to 6-frame translated ESTs). In addition to BLAST searches of the entire dbEST database, NCBI offers subsets of mouse ESTs, human ESTs and "other ESTs" for searching. Do not be confused by other database options such as sequence tagged sites (dbSTS; short marker sequences for genome mapping projects), genome survey sequences (gss) or high-throughput genomic sequences (htgs); the data in these sections are not derived from cDNAs and are not directly comparable to ESTs.

When might a typical molecular biologist benefit from a search of EST data?

(i) If similarity searches of other databases have not yielded any significant matches, then the ESTs can be used as a databank of last resort. If a sequence matches nothing in the EST databank, then the chances are very good that it is not an expressed sequence.

(ii) The EST databanks contain new members of known protein families. Researchers who suspect that they have identified a new protein by function may discover candidates

news.group.news

News.group.news goes to cyberspace and back to gather Net news for you. We emphasize practical, methods-related issues that are discussed in many newsgroups and also feature issues of general interest to biological scientists. But be warned! On this page of *BioTechniques*, we don't *review* it — we just *report* it!

Netters note that a comprehensive database of **information on transgenic animals** and targeted mutations is available at TBASE at the Web address (<http://www.bis.med.jhmi.edu/Dan/tbase/tbase.html>).

To remove the silanizing compound from glass sequencing gel plates, netters recommend soaking the plates in 0.5 M NaOH, 0.5 M KOH or methanoic acid (5 kg KOH in 100 mL MeOH) for several hours.

Seeking a recent review on the effect of **rare arginine codons** AGA/AGG on heterologous gene expression in *E. coli*, a netter on **bionet.molbio.methds-reagents** was advised to consult C. Kurland and J. Gallant, 1996, Errors of heterologous protein expression, *Curr. Opin. Biotechnol.* 7:489-493.

A non-botanist seeking advice on **normalizing plant Northern**s was advised to probe his blot of RNAs expressed by red kidney beans with probes for 18S rRNA or cytosolic GAPDH.

BioTechniques Editorial Department

Internet On-Ramp

among ESTs that are similar but not identical to other known proteins.

(iii) EST sequences include many mutations and allelic variations of known genes. Investigators interested in gene diversity or seeking polymorphisms as genetic markers can obtain useful data from the EST databanks.

If an interesting EST sequence is identified in the databanks, the original cDNA clone from which that sequence was derived can be obtained from the IMAGE consortium for \$45 for the first clone and \$24 for each additional clone (<http://www.bio.lnl.gov/bbrp/image/>)

The redundancy of the EST databanks can be used to partially compensate for the high error rate and the fragmentary nature of the individual sequences. The reverse transcription reactions that are used to create cDNA libraries do not all run to completion. The pool of cDNA clones contains many partial sequences of varying lengths; therefore, ESTs databases will contain partially overlapping sequences from the same gene. Several groups are working on procedures to cluster all ESTs that are derived from a single gene. In this way, partial and fragmentary sequences can be assembled into contigs that correspond to whole gene transcripts. Thus, the vast, low-quality EST databanks can be collapsed into a single consensus set of nonredundant transcripts from all expressed human genes.

Investigators can conduct routine similarity searches against the EST clusters, which require only a tiny fraction of the disk storage space used by the complete EST databank; these searches can be computed much more quickly. Similarity matches to the EST clusters will be much more informative than matches to the current set of minimally annotated ESTs. Those wishing to examine tissue-specific gene frequencies or allelic sequence variations can extract the original EST sequences used to create the consensus from a central repository (i.e., dbEST at the NCBI Web site).

There are three different sets of clustered human ESTs available on the Web. Each set has been created with different software using different assumptions as to data quality, minimum overlap etc. The three sets of consensus sequences are available for download, so investigators can use their own searching software against these databases.

UniGene

The NCBI, home of GenBank and BLAST, has created UniGene, a collection of human EST sequences from GenBank that have been grouped into clusters by sequence similarity. Some of these clusters correspond to genes that have been identified by other means (i.e., functional cloning), while others are genes of unknown function. As of January 1998, UniGene contains 39 377 clusters. This is a much more manageable set of sequences to search than the 1 409 359 sequences in the complete EST section of GenBank (January 1998). The UniGene collection is also indexed by tissue of origin, so it is possible to search for genes that are expressed in a particular tissue. For more information, see the <http://www.ncbi.nlm.nih.gov/UniGene/index.html> address.

TIGR Human Gene Index

The Institute for Genomic Research (TIGR) has created a Human Gene Index (HGI) for their own collection of ESTs

BioBit: Fields for Searching the dbEST Database

The EST section of GenBank is directly searchable on the NCBI Web site (<http://www.ncbi.nlm.nih.gov/dbEST/index.html>), using the following fields:

DBID	EST id number
IDS	EST name, GenBank accession, gi number
CLIN	Clone information, source information
COM	Comments
LIB	Library name and organism
LIBX	Library description
SUB	Submitter information
CIT	Citation information
MAP	Mapping data
NBR	Homology (neighboring) information

(<http://www.tigr.org/tdb/hgi/hgi.html>). As of July 1997 (HGI Version 3.2), TIGR has identified 60 388 "tentative human consensus" (THC) sequences by assembling over 600 000 ESTs into complete "virtual transcripts". THCs are consensus sequences based on two or more ESTs that overlap for at least 40 bases with at least 95% sequence identity. THCs retain information about the source library and abundance of the ESTs from which they are composed. TIGR also provides a free Web-based similarity search service that can be used to compare query sequences against the HGI. Similarity results are returned by e-mail in less than 5 min. Clones corresponding to sequences in the TIGR EST database can be obtained from the American Type Culture Collection (ATCC, Rockville, MD, USA).

STACK

The South African National Bioinformatics Institute (SANBI) at the University of the Western Cape has created an EST consensus database known as Sequence Tag Alignment and Consensus Knowledgebase (STACK; <http://ziggy.sanbi.ac.za/stack/>). It consists of all available human EST sequences in GenBank, clustered, aligned and made into joined consensus records. STACK consensus sequences are often longer than TIGR or UniGene clusters. STACK creates separate consensus clusters for ESTs from each of 17 different tissue types.

SANBI provides a free Web-based BLAST search utility that can be used to compare query sequences against the STACK clusters. Similarity searches can be calculated as DNA-DNA (BLASTN), protein query against 6-frame translations of the STACK database (TBLASTN) or 6-frame translation of a DNA query against 6-frame translations of the STACK database (TBLASTX). There is also the option of returning search results by e-mail.

The EST databanks give researchers access to basic data from the genome project almost as soon as it is collected in the laboratory. This allows those investigators who are willing to sift through mountains of low-quality data to discover some hidden jewels. For most others, the pre-analyzed clustered EST databases will be much more useful. The promise of EST data continues to be "your gene is out there." You just have to have the patience and persistence to look for it and to be smart enough to know when you have found it.