Short Technical Reports

dominant form of integration in root-derived transgenics, whereas multiple insertions are found in leaf discs. *Plant Mol. Biol.* 23:847-860.

11. **Hecker, K.H. and K.H. Roux.** 1996. High and low annealing temperatures increase both specificity and yield in touchdown and step-down PCR. *BioTechniques* 20:478-485.
12. **Hobbs, S.L., T.D. Warkentin and C.M. De-Long.** 1993. Transgene copy number can be positively or negatively associated with transgene expression. *Plant Mol. Biol.* 21:17-26.
13. **Iglesias, V.A., E.A. Moscone, I. Papp, F. Neuhauser, S. Michalowski, T. Phelan, S. Spiker, M. Matzke and A.J. Matzke.** 1997. Molecular and cytogenetic analyses of stably and unstably expressed transgene loci in tobacco. *Plant Cell* 9:1251-1264.
14. **Mayerhofer, R., K.Z. Koncz, C. Nawrath, G. Bakkeren, A. Cramer, K. Angelis, G.P. Redei, J. Schell, B. Hohn and C. Koncz.** 1991. T-DNA integration: a mode of illegitimate recombination in plants. *EMBO J.* 10:697-704.
15. **McGarvey, P. and J.M. Kaper.** 1991. A simple and rapid method for screening transgenic plants using the PCR. *BioTechniques* 11:428-432.
16. **Nagata, T., Y. Nemoto and S. Hasegawa.** 1992. Tobacco BY-2 cell line as the "HeLa" cell in the cell biology of higher plants, p. 1-30. In K.W. Jeon and M. Friedlander (Eds.), *International Review of Cytology*. Academic Press, San Diego.
17. **Prod'homme, G., C. Guilhot, M.C. Gutierrez, A. Varnerot, B. Gicquel and V. Vincent.** 1997. Rapid discrimination of *Mycobacterium tuberculosis* complex strains by ligation-mediated PCR fingerprint analysis. *J. Clin. Microbiol.* 35:3331-3334.
18. **Sambrook, J., E.F. Fritsch and T. Maniatis.** 1989. *Molecular Cloning: A Laboratory Manual*, 2nd ed. CSH Laboratory Press, Cold Spring Harbor, NY.
19. **Siebert, P.D., A. Chenchik, D.E. Kellogg, K.A. Lukyanov and S.A. Lukyanov.** 1995. An improved PCR method for walking in uncloned genomic DNA. *Nucleic Acids Res.* 23:1087-1088.
20. **Swensen, J.** 1996. PCR with random primers to obtain sequence from yeast artificial chromosome insert ends or plasmids. *BioTechniques* 20:486-491.
21. **Triglia, T., M.G. Peterson and D.J. Kemp.** 1988. A procedure for in vitro amplification of DNA segments that lie outside the boundaries of known sequences. *Nucleic Acids Res.* 16:8186.
22. **Trueba, G.A. and R.C. Johnson.** 1996. Random primed gene walking PCR: a simple procedure to retrieve nucleotide fragments adjacent to known DNA sequences. *BioTechniques* 21:20.
23. **Valvekens, D., M. Van Montagu and M. Van Lijsebettens.** 1988. *Agrobacterium tumefaciens*-mediated transformation of *Arabidopsis thaliana* root explants by using kanamycin selection. *Proc. Natl. Acad. Sci. USA* 85:5536-5540.
24. **Zhou, Y., R.J. Newton and J.H. Gould.** 1997. A simple method for identifying plant/T-DNA junction sequences resulting

from *Agrobacterium*-mediated DNA transformation. *Plant Mol. Biol. Rep.* 15:246-254.

The authors wish to thank Colette Tremblay for careful revision of the manuscript. This work was supported by National Sciences and Engineering Research Council of Canada Operating Grant No. A6923, and a Strategic Grant to G.B. D.S. was supported by a fellowship from the National Sciences and Engineering Research Council of Canada (NSERC) and from the Fonds pour la formation de chercheurs et l'aide à la recherche (Fonds FCAR). Address correspondence to Guy Bellemare, Département de biochimie, Pavillon Charles-Eugène-Marchand, Université Laval, Québec, QC, Canada, G1K 7P4. Internet: bellemar@rsvs.ulaval.ca

Received 12 January 1999; accepted 10 May 1999.

Diego Spertini, Catherine Béliveau and Guy Bellemare
Université Laval
Québec, QC, Canada

Nonradioactive Labeling of Large DNA Fragments for Genome Walking, RFLP and Northern Blot Analysis

BioTechniques 27:314-320 (August 1999)

ABSTRACT

*In this report, we present a simple non-radioactive labeling procedure for DNA fragments of high specific labeling density that can be used for a variety of applications. The protocol is based on the universal mono-functional platinum reagent for chemical digoxigenin (DIG) labeling of nucleic acids. The labeling protocol was optimized for large DNA templates as complete bacterial artificial chromosomes (BAC). Variations of incubation time and temperature improved the labeling density such that about 30% of the nucleotides were DIG-modified within 30 min. Furthermore, the refined procedure generates in a single-tube reaction and without prior digestion-labeled DNA fragments of 0.5–4.0 kb from a 130-kb template. Hybridization experiments were performed on Southern and northern blots and allowed the detection of single copy genes in 2.5 µg genomic DNA from *Arabidopsis thaliana*, which has a haploid genome size of 0.13 pg (ca. 120 Mb) and medium expressed transcripts from 0.8 µg poly(A)⁺ RNA, respectively. The extremely high specific labeling density, the stability and the universal application of the probe generated with the platinum reagent makes this method a useful alternative to classical radioactive nuclei acids labeling techniques.*

INTRODUCTION

In recent years, nonradioactive labeling and detection systems of nucleic acids have overcome the safety, disposal, stability and cost problems that are associated with radioactive techniques. In contrast to traditional nonradioactive labeling methods (random priming, nick translation, cDNA synthesis, polymerase chain reaction [PCR], 3' end-labeling and 3' tailing), which are enzyme-mediated, the newly developed

Short Technical Reports

chemical labeling compounds no longer depend on the enzymatic incorporation of labeled nucleotides. The special *cis*-platinum Chem-Link Reagent (Roche Diagnostics GmbH) forms stable coordination bonds with the N7 atom of purine bases and therefore yields higher labeling densities (4,9).

In this report, we modified the available protocol for the chemical labeling method with digoxigenin (DIG) (4) in order to adapt it for the labeling of DNA probes from large DNA templates. We used a whole bacterial artificial chromosome (BAC) clone of 130 kb to optimize the DIG-Chem-Link Reagent method. Our experiments indicate that it is not necessary to cut the large DNA fragments before labeling. The resulting probe was sensitive enough to detect 0.3 pg of homologous DNA.

MATERIALS AND METHODS

DNA Isolation

For the labeling experiments, the BAC DNA was isolated using Nucleobond® AX-100 Cartridges (Macherey-Nagel, Duren, Germany). BAC DNA isolation for Southern blot analysis was prepared by a modified standard alkaline lysis method from 5 mL overnight cultures (11). Plant DNA was isolated from about 0.7 g frozen tissue according to the protocol of Dellaporta et al. (2).

Plant RNA Isolation

Total RNA was prepared from frozen tissue of *Arabidopsis thaliana* roots and shoots using TRIzol® LS Reagent (Life Technologies, Wien, Austria). Following the manufacturer's instructions, the yield of total RNA was about 0.3–0.5 µg/mg tissue. Poly(A)⁺ RNA was isolated from total RNA using an Oligotex™ mRNA Midi Kit (Qiagen GmbH, Hilden, Germany). The mRNA yield varied between 1% and 0.3% from total RNA of roots and shoots, respectively.

Northern Blot Analysis

Poly(A)⁺ RNA (0.8–1.3 µg) and 5 µg of total RNA were separated under de-

Table 1. Comparison of the Different DIG-Chem-Link Labeling Experiments

Labeling Conditions	Probe Size in kb	Efficiency		Sensitivity
		ng-Labeled Probe per µg Template	Labeling Density	
Undigested				
85°C/30 min	0.8–30	125 ng/µg	ca. 1 in 8 bases	ca. 1 pg
85°C/60 min	0.8–20	125 ng/µg	ca. 1 in 8 bases	ca. 1 pg
85°C/120 min	0.5–4	315 ng/µg	ca. 1 in 3 bases	ca. 0.3 pg
98°C/30 min	0.5–4	315 ng/µg	ca. 1 in 3 bases	ca. 0.3 pg
98°C/120 min	0.5–0.8	315 ng/µg	ca. 1 in 3 bases	ca. 0.3 pg
Digested				
50°C/30 min	n.d. ^a	<5 ng/µg	<1 in 200 bases	n.d.
85°C/30 min	<0.5–10	125 ng/µg	ca. 1 in 8 bases	ca. 1 pg
a n.d. = not determined.				

^an.d. = not determined.

naturing conditions on a 1% agarose gel, containing 6% formaldehyde in 1× MOPS buffer (20 mM 4-morpholinopropanesulfonic acid, 5 mM sodium acetate, 1 mM EDTA, pH 7.0). Transfer of RNA to a positively charged nylon membrane was performed using the downward alkaline transfer protocol of Münch (7). After 2 h, the transfer was

stopped, and the RNA was cross-linked to the membrane by 120 mJ of UV light.

Southern Blot Analysis

*Eco*RI-digested plant genomic DNA (2.5 µg) was separated on 1% agarose gels in 1× TAE (40 mM Tris, 5 mM sodium acetate, 1 mM EDTA, pH 7.8).

Figure 1. Effect of different reaction conditions on the integrity of the labeled 130-kb BAC clone. For Southern blot analysis, 1/25 of the different labeling reactions were separated, blotted and hybridized as described in Materials and Methods. By increasing the time and temperature of the labeling reaction, the average size of the DNA fragments becomes smaller. As molecular weight marker, 50 ng DIG-labeled λ HindIII DNA were used (M).

Figure 2. Dot blot quantification of the labeled BAC 6 probe. Dilutions of the labeled control DNA and the probe DNA were spotted, fixed and detected on a positively charged nylon membrane using chemiluminescent detection with anti-DIG-AP antibody and CSPD as substrate. The incorporation of digoxigenin was estimated by comparing the spot intensities of the control (5 ng/µL) to those of the diluted probes. Estimated concentrations are indicated on the bottom of the filter. Since for each labeling reaction the same amount of template DNA and DIG-Chem-Link was used, the dependency of temperature and time on labeling density can be quantified.

One-tenth of each BAC DNA was digested with *Sa*II and separated on 1% agarose gels with 0.5× TBE (89 mM Tris, 89 mM boric acid, 2 mM EDTA, pH 8.3) under pulsed-field electrophoresis conditions (CHEF Mapper™ System, autoalgorithm for 2–30 kb; Bio-Rad GmbH, Wien, Austria). After standard capillary transfer to a positively charged nylon membrane (Roche Diagnostics GmbH, Wien, Austria) with 10× SSPE buffer (150 mM NaCl, 10 mM NaH₂PO₄, 1 mM EDTA, pH 7.4), DNA fragments were cross-linked with 120 mJ of UV light.

Labeling of BAC DNA

The labeling reaction was performed at different temperatures (85° and 98°C) and for variable periods of time (30, 60 and 120 min) in a 20-μL reaction volume containing 2 μg DNA, 2 μg of the platinum-DIG complex (DIG-Chem-Link, C₃₉H₆₇N₇O₁₅Pt; Roche Diagnostics GmbH) and sterile water. Condensate at the tube lid was collected by centrifugation. The labeling reaction was stopped by adding 5 μL stop solution (Roche Diagnostics GmbH).

Labeling efficiency was determined by dot blot analysis and comparison with a labeled probe of known digoxigenin concentration. The labeling density was calculated from the dot blot results and normalized to the template concentration. Sensitivity of the probe was measured by dot blot analysis of a dilution series of homologous DNA of known concentration. The integrity of the labeled probe was determined by Southern blot analysis. The labeled probe (1/25) was separated on a 0.7% agarose gel, blotted as above, and the DNA fragments with incorporated DIG were determined by chemiluminescent detection.

Hybridization and Detection

Hybridization and detection was essentially done as described by the manufacturer's protocols for the DIG system (Roche Diagnostics GmbH). Hybridizations were performed in sealed plastic bags (autoclaving bags; Product No. 644200; Greiner, Kremsmünster, Austria) with the DIG Easy-Hyb Buffer (Roche Diagnostics GmbH)

Figure 3. Southern blot analysis for the construction of a the BAC contig. (A) The *Sa*II-digested BAC DNA separated on a 1% agarose gel stained with EtdBr. (B) The same gel was blotted onto a filter and hybridized with the labeled BAC clone 6 DNA. After 30-min exposure, all DNA fragments hybridizing with BAC clone 6 are visible. BAC clone 14 does not belong to the BAC contig, and therefore only the fragment of the vector lights up. BAC clone 7 has a smaller backbone compared with all other BAC clones. As DNA length marker, 50 ng DIG-labeled λ HindIII DNA were used (M).

Short Technical Reports

containing 20–30 ng labeled probe per mL. After incubation overnight at 42°C, the blots were washed with 2× SSC (150 mM NaCl, 15 mM sodium citrate, pH 7.0), 0.1% sodium dodecyl sulfate (SDS) at room temperature, followed by 0.1× SSC, 0.1% SDS at 55°C for plant DNA and 60°C for BAC DNA.

Hybridization of northern blots were done at 68°C using the same hybridization solution as for Southern blot analysis. The stringency wash for northern blots used 0.5× SSC, 0.1% SDS at 68°C.

For sensitive detection, the chemiluminescent detection procedure with CSPD®-ready-to-use (Roche Diagnostics GmbH) was utilized according to the manufacturer's instructions, and the light signal was recorded on X-ray films. Northern blots were stripped for 10 min with boiling 0.1% SDS solution and reprobbed with an abundantly expressed ubiquitin extension protein and a low expressed *Athb-14* probe. These DIG-labeled probes were prepared by amplification of appropriate sequences using PCR essentially as described by Finckh et al. (3). The templates were amplified from genomic DNA with the primers NUBQ (5'-GGTGCTAAGAA-GAGGAAGAAT-3') and CUBQ (5'-C-TCCTTCTTTCTGGTAAACGT-3') for ubiquitin, and AthB2F (5'-GTAGCG-ATGGTGCAGAGGATGTTA-3') and AthB2R (5'-CTGCGCAAGAGGAA-GAATGACT-3') for *Athb-14*. CUBQ and AthB2R was used for the synthesis of the antisense probes.

RESULTS AND DISCUSSION

Optimization of the Labeling Conditions

The efficiency of the labeling reaction with the DIG *cis*-platinum compound (DIG-Chem-Link) depends on the incubation time and temperature, the ratio of the reaction partners, the final reaction volume and the purity of the nucleic acid (4). As recommended by the manufacturer, the reaction should be performed with a 1:1 (μg DNA template:μg DIG-Chem-Link) ratio, in a small volume of 20 μL and at incubation conditions of 85°C for 30 min. Under these standard conditions, the DNA template becomes slightly

Figure 4. Southern blot analysis to detect allele-specific polymorphisms. Genomic *A. thaliana* DNA of wild-type Columbia (Col), Landsberg *erecta* (Ler), Wassilewskija (Ws), eight mutant alleles and the BAC 6 clone were digested with *Eco*RI, separated on a 1% agarose gel, blotted and hybridized with 30 ng/mL labeled BAC clone 6. As DNA length marker, 50 ng DIG-labeled λ HindIII DNA were used (M). Fragments larger than 10 kb are the result of partial digestion. An ecotype-specific polymorphism between Col and Ler/Ws is seen around fragment 5/5.2 kb, respectively (open circle). Allele-specific polymorphisms could be detected in alleles 1-3, 1-4, 1-5, 1-14 and 1-18 (see stars).

Figure 5. Northern blot analysis using a whole labeled BAC as probe. (A) Displays the EtdBr-stained denaturing agarose gel. (B) The same gel was blotted onto a membrane and hybridized with the labeled BAC 6 probe. (C) The same membrane was simultaneously hybridized with the ubiquitin extension protein and the *AthB-14* gene probe (see Materials and Methods). RNA from root tissue was loaded in lanes 1 and 3, and RNA from shoot tissue was loaded in lanes 2 and 4. Lanes 1 and 2 contained 1.3 μg poly(A)⁺ RNA and lanes 3 and 4 contained 5 μg total RNA. As RNA length marker, 50 ng DIG-Labeled RNA Length Standard II (Roche Diagnostics GmbH) were used (M).

fragmented (Figure 1). Under milder reaction conditions, such as 50°C and/or a short incubation time, the original fragment size is preserved, but labeling efficiency is very low (Table 1).

For optimal hybridization results, it is necessary that the labeled probe is not too large. Hence, we tested whether it is necessary to digest large DNA templates before labeling or whether undigested complete BACs of 130 kb result in labeled probes of satisfactory length and high incorporation of the DIG. The comparison was done between an *Eco*RI-digested BAC template with DNA fragments ranging from <0.5–10 kb and a complete, circular 130-kb BAC template. As summarized in Table 1, under standard reaction conditions of 85°C for 30 min, the digested BAC DNA had the same labeling efficiency as the undigested template. However, the average fragment size of the labeled probe was smaller for the digested tem-

plate. Nevertheless, this size range differences had no impact on the sensitivity of the probe (Table 1). Thus, it is not necessary to digest large DNA fragments before the labeling reaction.

As noted above, the labeling density is influenced by temperature and reaction time. Therefore, we investigated if the labeling efficiency and the probe sensitivity can be extended by increasing the reaction time and/or temperature. We tested different time and temperature conditions on their effect on probe fragment size, labeling efficiency and density as well as probe sensitivity (Table 1, Figures 1 and 2). Our results indicate that the optimal reaction conditions for labeling large DNA fragments were either 85°C for 120 min or 98°C for 30 min. Under these conditions, the labeled fragments ranged from about 0.5 to up to 4 kb. The fact that two different incubation conditions, 85°C for 120 min and 98°C for

30 min, result in the same fragment sizes indicates that both temperatures or times can be altered to obtain similar probe sizes. Moreover, whereas standard protocols yield one DIG label for every ten bases, these two labeling conditions lead to labeled probe with up to one DIG label in every three nucleotides (Table 1). To our knowledge, such a high specific labeling density is not achieved by current radioactive labeling techniques.

With these probes it was possible to detect <0.3 pg of complementary DNA within 30 min of exposure time (data not shown).

Application of the Nonradioactively Labeled BACs

Construction of BAC contigs.

While the present BAC or yeast artificial chromosome (YAC) physical maps are determined by fingerprint analysis

Short Technical Reports

of ethidium bromide (EtdBr)-stained gels, these analyses can be improved by hybridization of labeled BAC end probes or even complete BACs (10). To achieve clearly resolved hybridization signals, a high specific labeling density of the probe and a good resolution of the BAC restriction enzyme fragments are essential. Hence, we tested whether our nonradioactive labeled BAC meets the criteria to establish BAC contigs. For the optimal separation of the BAC DNA fragments, we used pulsed-field electrophoresis (see Materials and Methods). Figure 3 shows one example of a Southern blot analysis of *SalI*-digested BAC DNAs, which were hybridized to the DIG-Chem-Link-labeled BAC clone 6. The presence of similar fragments in multiple clones indicates potential overlaps. Furthermore, the cumulative sizes of these shared restriction fragments give an estimate of the extent of the overlap between BACs. This approach can facilitate contig assembly by fingerprinting large-range restriction mapping of genomic DNA and hence is suitable for positional cloning strategies.

Analysis of allele-specific polymorphisms. Once a BAC contig is established and a gene of interest is located on one of these BAC clones by conventional recombination analysis, the final isolation of the gene depends on the availability of polymorphic markers close by. One possibility to position the gene of interest on a sub-fragment of the BAC clone is to find allele-specific polymorphisms among mutant alleles (5,6). Here, we show that the specific labeling of the complete 130-kb BAC clone was dense enough to screen for restriction fragment length polymorphisms (RFLPs) in 2.5 µg genomic DNA derived from several mutant alleles of *A. thaliana* (Figure 4).

Northern blot analysis with the whole BAC DNA as probe. To detect the transcripts that are encoded by the BAC insert, we performed northern blot analysis by using the whole BAC DNA as a probe and compared different RNA preparations. With an average gene density of 4.8 kb (1), a 130-kb BAC clone could theoretically contain 27 genes in *A. thaliana*. In case many genes are expressed, the analysis might lead to diffuse and undiscernable

bands. Nevertheless, by using total RNA, we were able to detect at least six distinct transcripts ranging between 0.5 to 2.6 kb in shoots and two transcripts in roots (Figure 5B). From these transcripts, one was enriched in shoot- and two in root-derived poly(A)⁺ RNA, indicating that the RNA bands detected in total RNA preparations were most probably rRNA and not polyadenylated (Figure 5A). Thus, the entire BAC probe can cross-hybridize with rRNA, and only poly(A)⁺ RNA revealed insert-specific transcripts.

To quantify the sensitivity of the BAC probe, we reprobated the membrane against the abundantly expressed ubiquitin extension protein and the organ specifically expressed *AthB-14* transcript (Figure 5C). In mRNA preparations, the ubiquitin transcripts were highly enriched, and the *AthB-14* transcripts were detectable only in roots. These results are consistent with the previously published root-specific expression of *AthB-14* (8). Furthermore, the results also indicate that the DIG-labeled BAC probe is at least as sensitive as the radioactive probe used by Sessa et al. (8) and that the transcripts detected with the labeled BAC probe are medium-expressed in an organ-specific manner.

The advantages of this new single-tube labeling method are its simplicity, the avoidance of enzymes and its ability to generate labeled probes with a high specific labeling density. As with all nonradioactively labeled probes, they can be stored for up to one year and used for at least five times without significant loss in sensitivity. Our findings, that the probes can be used for DNA and RNA hybridization experiments, makes this technique a universal tool for the detection of nucleic acids.

REFERENCES

1. Bevan, M., I. Bancroft, E. Bent, K. Love, H. Goodman, C. Dean, R. Bergkamp, W. Dörk et al. 1998. Analysis of 1.9 Mb of contiguous sequence from chromosome 4 of *Arabidopsis thaliana*. *Nature* 391:485-488.
2. Dellaporta, S.L., J. Wood and J.B. Hicks. 1993. A plant DNA miniprep: version 2. *Plant Mol. Biol. Rep.* 1:19-22.
3. Finckh, U., P.A. Lingelfelter and D. Myerson. 1991. Producing single-stranded DNA probes with the *Taq* DNA polymerase: a high yield protocol. *BioTechniques* 10:35-39.
4. Holz, H., C. Goldstein, R. Rein and H.J. Höltke. 1997. A new member of labeling reagents, optimized for a variety of nucleic acids, independent of size or conformation. *Biochemica* 3:26-28.
5. Li, J. and J. Chory. 1997. A putative leucine-rich repeat receptor kinase involved in brassinosteroid signal transduction. *Cell* 90:929-938.
6. Lukowitz, W., U. Mayer and G. Jürgens. 1996. Cytokinesis in the *Arabidopsis* embryo involves the syntaxin-related KNOLLE gene product. *Cell* 84:61-71.
7. Münch, S. 1994. Nonradioactive Northern blots in 1.5 days and with substantially increased sensitivity through "Alkaline Blotting". *Biochemica* 3:30-32.
8. Sessa, G., C. Steindler, G. Morelli and I. Ruberti. 1998. The *Arabidopsis Athb-8, -9* and *-14* genes are members of a small gene family coding for highly related HD-ZIP proteins. *Plant Mol. Biol.* 38:609-622.
9. van Belkum, A., E. Linkels, T. Jelsma, F.M. van den Berg and W. Quint. 1994. Non-isotopic labeling of DNA by newly developed hapten-containing platinum compounds. *BioTechniques* 16:148-153.
10. Wang, M.L., L. Huang, D.K. Bongard-Pierce, S. Belmonte, E.A. Zachgo, J.W. Morris, M. Dolan and H.M. Goodman. 1997. Construction of an approximately 2 Mb contig in the region around 80 cM of *Arabidopsis thaliana* chromosome 2. *Plant J.* 12:711-730.
11. Woo, S.-S., J. Jiang, B.S. Gill, A.H. Patterson and R.A. Wing. 1994. Construction and characterization of a bacterial artificial chromosome library of *Sorghum bicolor*. *Nucleic Acids Res.* 22:4922-4931.

This work was supported by the Austrian Science Foundation (FWF, P11001-GEN) and with the EU-grant (PL-960217). We thank Eva Dornstauder for excellent technical assistance and Dr. Josef Glössl for providing the stimulating environment and also Christian Schlötterer and Christian Luschig for critical reading of the manuscript. Address correspondence to Dr. Marie-Theres Hauser, Center of Applied Genetics, University of Agricultural Sciences, Muthgasse 18, A-1190 Vienna, Austria. Internet: hauser@mail.boku.ac.at

Received 3 February 1999; accepted 3 May 1999.

Farzaneh Adhami, Sabine Müller and Marie-Theres Hauser
*University of Agricultural Sciences Vienna
Vienna, Austria*